


Drug Use

Percentage (%) of recent mothers in Illinois who reported taking prescription pain relievers during pregnancy

Survey Question D11: During your most recent pregnancy, did you use prescription pain relievers such as hydrocodone (Vicodin®), oxycodone (Percocet®), or codeine? Response: Yes



	Percentage <sup>1</sup>	95% CI <sup>2</sup>		Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>	
<b>All</b>	Overall	3.9	2.3	6.8	379	1,733
<b>Maternal Age</b>	<20 yrs	--	--	--	--	--
	20 to 24 yrs	7.1	2.5	18.4	60	520
	25 to 29 yrs	2.9	0.8	9.9	103	362
	30 to 34 yrs	6.4	2.9	13.9	119	794
	=>35 yrs	0.5	0.1	2.1	87	56
<b>Race/Ethnicity</b>	NH White	4.7	2.3	9.1	203	1,142
	NH Black	2.0	0.3	12.8	65	134
	Hispanic	2.7	0.5	13.6	70	228
	Other	6.1	1.8	19.1	35	228
<b>Education</b>	<12 yrs	6.1	1.7	19.9	52	418
	12 yrs	5.0	1.5	15.3	66	394
	>12 yrs	3.2	1.6	6.4	251	921
<b>Birth Weight</b>	Low	6.5	3.1	12.9	104	207
	Normal	3.7	2.0	6.9	275	1,526
<b>Marital Status</b>	Married	3.9	2.0	7.5	238	1,054
	Other	3.9	1.5	10.1	141	678
<b>Insurer Now</b>	Medicaid	3.5	1.1	11.2	107	457
	Other	4.1	2.2	7.5	272	1,276

<sup>1</sup>Percentage = Weighted percentage

<sup>2</sup>CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

<sup>3</sup>Respondents = The total number who responded to the survey question

<sup>4</sup>Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Data collected using Marijuana and Drug Use Supplement beginning December 2017. Data collection ended March 2019.