


Drug Use

Percentage (%) of recent mothers in Illinois who reported taking prescription pain relievers during pregnancy

Survey Question D11: During your most recent pregnancy, did you use prescription pain relievers such as hydrocodone (Vicodin®), oxycodone (Percocet®), or codeine? Response: Yes



	Percentage ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴		
All	Overall	5.0	3.8	6.5	1,291	6,765
Maternal Age	<20 yrs	--	--	--	--	--
	20 to 24 yrs	4.3	2.0	9.0	200	932
	25 to 29 yrs	6.4	4.0	10.0	349	2,428
	30 to 34 yrs	4.7	3.0	7.5	408	1,963
	=>35 yrs	3.4	1.8	6.3	281	974
Race/Ethnicity	NH White	4.2	2.8	6.1	697	3,049
	NH Black	6.0	3.3	10.6	245	1,361
	Hispanic	5.9	3.4	10.1	240	1,627
	Other	5.5	2.3	12.9	101	571
Education	<12 yrs	7.3	3.7	14.0	138	1,018
	12 yrs	6.8	4.2	10.9	287	2,159
	>12 yrs	4.0	2.8	5.6	845	3,432
Birth Weight	Low	6.4	4.3	9.6	334	657
	Normal	4.9	3.7	6.5	957	6,107
Marital Status	Married	4.4	3.1	6.2	800	3,644
	Other	6.0	4.0	8.9	491	3,120
Insurer Now	Medicaid	6.3	4.0	9.6	391	2,473
	Private	4.2	2.9	6.1	742	3,275
	Uninsured	6.8	3.3	13.5	132	1,017

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents or numerator <6 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition