

## Other Tobacco Products

## Percentage (%) of recent mothers in Illinois who reported using other tobacco products in the past two years

Survey Question 37: Have you used any of the following products in the past 2 years? Response: Yes

	Percentage <sup>1</sup>	95% CI <sup>2</sup>		Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>
E-cigarettes or other electronic nicotine products	4.6	3.5	6.0	1,362	6,624
Hookah	4.3	3.2	5.7	1,358	6,206
Cigars, cigarillos, or little filtered cigars	3.3	2.4	4.6	1,357	4,795
Chewing tobacco, snuff, snus, or dip	0.4	0.2	0.9	1,359	617

## Percentage (%) of recent mothers in Illinois who reported using other tobacco products in the three months before pregnancy or the last three months of pregnancy

## Among women who reported using other tobacco products in the past two years

Survey Question 38: During the 3 months before you got pregnant, on average, how often did you use e-cigarettes or other electronic nicotine products? Responses 'Yes/No' created using numeric response

Survey Question 39: During the last 3 months of your pregnancy, on average, how often did you use e-cigarettes or other electronic nicotine products? Responses 'Yes/No' created using numeric response

	Percentage <sup>1</sup>	95% CI <sup>2</sup>		Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>
E-cigarette use 3 months before pregnancy	2.0	1.3	2.9	1,365	2,857
E-cigarette use last 3 months of pregnancy	0.3	0.1	0.9	1,362	493

<sup>1</sup>Percentage = Weighted percentage<sup>2</sup>CI = confidence interval

-- Indicates denominator &lt;30 respondents

<sup>3</sup>Respondents = The total number who responded to the survey question<sup>4</sup>Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition