


Prescription Pain Reliever Use

Percentage (%) of recent mothers in Illinois who reported using prescription pain relievers during pregnancy

Survey Question O2: During your most recent pregnancy, did you use any of the following prescription pain relievers? Response: Yes



	Percentage <sup>1</sup>	95% CI <sup>2</sup>		Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>
Codeine	3.3	2.2	4.8	915	3,174
Hydrocodone	2.9	1.9	4.4	915	2,795
Morphine	1.0	0.5	2.1	912	963
Tramadol	1.0	0.5	2.0	912	960
Oxycodone	0.7	0.3	1.5	912	662

<sup>1</sup>Percentage = Weighted percentage

<sup>2</sup>CI = Confidence interval lower/upper bounds

<sup>3</sup>Respondents = The total number who responded to the survey question

<sup>4</sup>Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Data collection period April - December 2019

Hydrocodone (like Vicodin®, Norco®, Lortab®)

Codeine (like Tylenol® #3 #4, not regular Tylenol®)

Oxycodone (like Percocet®, Percodan®, OxyContin®, Roxicodone®)

Tramadol (like Ultram® or Ultracet®)

Hydromorphone or meperidine (like Demorol®, Exalgo®, Dilaudid®) - Not reported numerator <6

Oxymorphone (like Opana®) - Not reported numerator <6

Morphine (like MS Contin®, Avinza®, Kadian®)

Fentanyl (like Duragesic®, Fentora®, Actiq®) - Not reported numerator <6