

2017
Illinois Pregnancy Risk Assessment
Monitoring System (PRAMS)

Detailed Data Tables

Technical Notes

Collection of Data

The Division of Health Data and Policy, Illinois Department of Public Health collected the 2017 Illinois Pregnancy Risk Assessment Monitoring System (PRAMS) data. Illinois uses the standardized data collection methods developed by the U.S. Centers for Disease Control and Prevention (CDC). These methods also are used by the 47 states and New York City, Puerto Rico, the District of Columbia, and the Great Plains Tribal Chairmen's Health Board (GPTCHB) participating in PRAMS. Data are collected through mailed surveys with telephone follow-up for non-respondents. A systematic stratified random sample of approximately 200 mothers is selected monthly from a frame of eligible birth certificates. At two to six months after delivery, each sampled mother is mailed an introductory letter and survey. A reminder letter and two additional survey mailings are sent to those who do not initially respond. PRAMS interviewers telephone mothers who do not respond to any of the mailed surveys to administer the survey by telephone. Surveys are available in English and Spanish, and telephone interviews are conducted in both languages.

In 2017, a total of 2,235 women were selected to participate in the study; 1,224 women completed the survey. Data were weighted and are representative of the 12-month PRAMS-eligible population of 141,656 recent mothers. Data collection during 2017 marks the second year using the Phase 8 PRAMS survey.

Sampling Design

A systematic stratified random sampling design is used to oversample low birthweight (less than 2,500 grams) births. The sample is stratified by low birthweight (less than 2,500 grams) and normal birthweight (2,500 grams or greater). The 2017 overall weighted response rate was 55%. The weighted responses rates by birth weight are 55% for both low birth weight and normal birth weight.

Weighting and Interpretation of Results

Statistics are based on weighted data. The weights were developed by CDC to adjust for sample design, non-response patterns, and omissions from the sampling frame. The final sampling weight used in the analysis of the survey data is the product of these three elements. Weighting is necessary to give unbiased estimates of population parameters.

The percentages, 95 percent confidence intervals, and total estimated population affected were calculated using Survey Data Analysis (SUDAAN®*) software. Estimates for response categories with small sample size (i.e., fewer than 30 respondents in the denominator) are not reported due to possible imprecision and bias.

PRAMS data are representative of Illinois resident women, age 14 years or older, who gave birth in Illinois to live infants. The sampling design is valid at the state level and not intended or developed to represent sub-state geographies or their populations without further evaluation. The data are not applicable to all pregnant women. At the direction of IDPH legal counsel, women younger than 18 years of age were not asked questions about physical abuse.

Acknowledgements

The Illinois PRAMS project would like to thank all the mothers who took the time to participate in our survey. These women provided valuable information to improve the understanding of why some babies are born healthy and some are not. The Illinois PRAMS project also would like to thank the CDC PRAMS team for their on-going operational and technical support.

This publication was supported by Cooperative Agreement Number 6U01DP006192-02-04 from the CDC.

Please call 1-866-643-7194 with any questions about Illinois PRAMS or email us at dph.pramtrac@illinois.gov.

Suggested citation: 2017 Illinois PRAMS Annual Report, Division of Health Data and Policy, Illinois Department of Public Health, 2019.

*SUDAAN® is a registered trademark of the Research Triangle Institute. © 2005 Research Triangle Institute. All rights reserved



Illinois PRAMS FACTS – 2017

Pregnancy Risk Assessment Monitoring System



The Illinois Pregnancy Risk Assessment Monitoring System (PRAMS) is an ongoing, population-based survey of women who delivered live born infants in Illinois. It is part of a national initiative of the U.S. Centers for Disease Control and Prevention (CDC) to reduce infant morbidity and mortality. More Illinois PRAMS information and detailed data can be found at <http://www.dph.illinois.gov/data-statistics/prams>.

SELECTED INDICATORS	PERCENTAGE% ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴
Before Pregnancy				
During the 12 months before pregnancy				
• Exercised three or more days per week	40.6	37.6-43.7	502	57,425
• Dieted to lose weight	27.7	25.0-30.5	332	39,104
• Had teeth cleaned by dentist or dental hygienist	63.0	59.4-66.5	551	62,718
• Had a health care visit with OB/GYN	59.0	55.3-62.5	538	58,724
• Had health care visit for illness or chronic condition	16.1	13.6-19.0	147	16,070
• Had health care visit for family planning or birth control	13.3	11.1-15.9	133	13,244
• Had health care visit for depression or anxiety	9.1	7.1-11.4	77	9,016
• Checked for diabetes by health care worker	20.2	17.8-22.9	243	28,536
• Taking regular prescription medications, other than birth control	19.0	16.7-21.6	242	26,857
• Talked with a health care worker about their family history	37.9	34.9-40.9	482	53,544
During the 3 months before pregnancy				
• Diagnosed with diabetes	1.6	1.0-2.5	25	2,199
• Diagnosed with high blood pressure	3.1	2.2-4.3	44	4,377
• Diagnosed with depression	9.7	8.0-11.8	115	13,704
Took multivitamins, prenatal vitamins, or folic acid vitamins every day during the month before pregnancy	36.9	34.0-39.9	475	52,092
Pregnancy unintended – want to be pregnant later, not at all, or not sure	42.8	39.7-45.9	513	60,241
Smoked during the three months before pregnancy	16.1	13.9-18.6	203	22,355
Consumed alcoholic drinks during the three months before pregnancy	62.4	59.3-65.4	741	87,557
Were physically abused by current or ex-husbands/partners, a family member, or someone else during 12 months before pregnancy	4.0	2.9-5.6	44	5,654
Enrolled in Medicaid the month before pregnancy	27.3	24.5-30.2	336	38,685

SELECTED INDICATORS	PERCENTAGE% ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴
During Pregnancy				
Received prenatal care during the first trimester	88.4	86.2-90.3	1,066	123,636
Used WIC services during pregnancy	33.9	30.9-37.1	385	46,460
Had teeth cleaned by dentist or dental hygienist	44.5	41.4-47.6	554	62,607
Diagnosed with depression	9.9	8.2-12.0	116	13,903
Received influenza immunization 12 months before or during pregnancy	58.3	55.2-61.4	714	81,740
Diagnosed with high blood pressure (starting in pregnancy) or (pre-)eclampsia	11.7	9.9-13.8	176	16,369
Diagnosed with gestational diabetes	8.8	7.2-10.7	125	12,395
Were physically abused by current or ex-husbands/partners, a family member, or someone else during pregnancy	2.5	1.7-3.8	30	3,505
Smoked last three months of pregnancy	6.8	5.3-8.6	89	9,414
Felt unsafe in the neighborhood where living	10.5	8.8-12.6	133	14,866
Enrolled in Medicaid for prenatal care	45.4	42.3-48.6	539	63,488
Postpartum Experiences and Infant Health				
Smoked after delivery	11.0	9.2-13.2	148	15,319
Diagnosed with depression	9.6	7.9-11.7	112	13,459
Had a postpartum checkup	92.4	90.5-94.0	1,121	129,243
Used postpartum birth control	80.9	78.3-83.2	965	113,393
Breastfed at least once after birth	88.2	86.0-90.1	1,048	121,861
Breastfed for at least 12 weeks after birth	60.7	57.5-63.9	690	80,147
Infant usual sleeping				
• Infant put to sleep on their backs most of the time	81.5	78.9-83.8	946	111,812
• Infant slept in crib/bassinet/play yard	88.9	86.7-90.8	1,046	121,522
• Infant slept with a blanket	44.0	40.9-47.2	518	59,524
Enrolled in Medicaid after delivery	42.5	39.2-45.8	456	52,900

¹Percentage = Weighted percentage of Illinois mothers giving birth to live infants in 2017 who responded Yes or as indicated to the survey question for the specified indicator

²CI = confidence interval

³Respondents (N) = Total number of mothers who responded Yes to the survey question for the specified indicator; <30 respondents results may be unreliable

⁴Estimated population affected = Estimated number of Illinois mothers giving birth in 2017 who would respond Yes or as indicated to the survey question for the specified indicator

The 2017 Illinois PRAMS survey was sent to 2,235 recent mothers in Illinois who delivered live born infants. The number of mothers responding to the survey was 1,224 for an unweighted response rate of 55%. The overall weighted response rate was 55%. The weighted response rates were 55% for mothers of low birth weight infants and for mothers of normal birth weight infants. (Source: Illinois 2017 Analysis Weights report.)

Illinois Department of Public Health
2017 Pregnancy Risk Monitoring System (PRAMS)

Detailed Data Tables Contents

Description	Table Number
2017 Illinois PRAMS Sample	Table 1
2017 Illinois PRAMS Detailed Tables	
Before Pregnancy	
Health Behaviors	
• Health activities 12 months before pregnancy	Table 2
• Dieting before pregnancy	Table 3
• Exercise before pregnancy	Table 4
• Health conditions	Table 5
• Vitamin usage everyday	Table 6
• Health care visit before pregnancy	Table 7
• Type of health care visit before pregnancy	Table 8
• Teeth cleaning 12 months before pregnancy	Table 9
• Content of health care visit before pregnancy	Table 10
• Pregnancy preparation discussed during healthcare visit	Table 11
• Pregnancy preparation topics discussed during healthcare visit	Table 12
• Pregnancy preparation discussed dental care	Table 13
• Pregnancy preparation discussed depression	Table 14
Family Planning	
• Feelings about being pregnant	Table 15
• Unintended pregnancy	Table 16
• Intended pregnancy	Table 17
• Unintended pregnancy and birth control use	Table 18
• Reasons for not using birth control	Table 19
During Pregnancy	
Prenatal Care	
• Prenatal care started in first trimester	Table 20
• Reasons early prenatal care was not received	Table 21
• Adequacy of prenatal care	Table 22
• Use of WIC during pregnancy	Table 23
• Health care topics discussed during prenatal care visits	Table 24
• Health care topics asked about during prenatal care visits	Table 25
• Folic acid awareness	Table 26
• Teeth cleaning during pregnancy	Table 27
• Health conditions during pregnancy	Table 28
Immunizations	
• Healthcare worker advice to get an influenza vaccination	Table 29
• Influenza vaccination 12 months before delivery	Table 30
Smoking	
• Cigarette smoking in past two years	Table 31
• Smoking during three months before pregnancy	Table 32

Description	Table Number
<ul style="list-style-type: none"> Smoking during last three months of pregnancy 	Table 33
<ul style="list-style-type: none"> Smoking now (at time of survey completion) 	Table 34
<ul style="list-style-type: none"> Number of cigarettes smoked 	Table 35
<ul style="list-style-type: none"> During prenatal care visit advised to quit smoking 	Table 36
Other Tobacco Products	
<ul style="list-style-type: none"> Other tobacco product use in past two years 	Table 37
Alcohol Consumption	
<ul style="list-style-type: none"> Alcohol use in past two years 	Table 38
<ul style="list-style-type: none"> Alcohol use during three months before pregnancy 	Table 39
<ul style="list-style-type: none"> Number of alcoholic drinks consumed 	Table 40
Stressors During Pregnancy	
<ul style="list-style-type: none"> Types of stressors 	Table 41
<ul style="list-style-type: none"> Number of stressors 	Table 42
<ul style="list-style-type: none"> Neighborhood safety 	Table 43
<ul style="list-style-type: none"> Unfair treatment during health care visit 	Table 44
<ul style="list-style-type: none"> Physical abuse before, during, or after pregnancy 	Table 45
After Pregnancy	
Breastfeeding	
<ul style="list-style-type: none"> Sources of breastfeeding information 	Table 46
<ul style="list-style-type: none"> Initiation of breastfeeding 	Table 47
<ul style="list-style-type: none"> Reasons for not breastfeeding 	Table 48
<ul style="list-style-type: none"> Breastfeeding continuation 1 month and 2 months after delivery 	Table 49
<ul style="list-style-type: none"> Exclusive breastfeeding continuation 12 weeks after delivery 	Table 50
Infant Sleep Habits	
<ul style="list-style-type: none"> Infant sleeps on back 	Table 51
<ul style="list-style-type: none"> Infant sleeps in own crib 	Table 52
<ul style="list-style-type: none"> Infant sleeps alone in same room with mother 	Table 53
<ul style="list-style-type: none"> Infant sleep environment 	Table 54
<ul style="list-style-type: none"> Infant sleep position topics discussed with health care worker 	Table 55
Birth Control Use after Delivery	
<ul style="list-style-type: none"> Use of birth control 	Table 56
<ul style="list-style-type: none"> Types of birth control used 	Table 57
<ul style="list-style-type: none"> Use of most or moderately effective birth control 	Table 58
<ul style="list-style-type: none"> Reasons for not using birth control 	Table 59
Postpartum Care	
<ul style="list-style-type: none"> Received a postpartum visit 	Table 60
<ul style="list-style-type: none"> Postpartum visit barriers 	Table 61
<ul style="list-style-type: none"> Postpartum visit location 	Table 62
<ul style="list-style-type: none"> Postpartum visit content 	Table 63
Mental Health	
<ul style="list-style-type: none"> Self-reported symptoms of depression after delivery 	Table 64
<ul style="list-style-type: none"> Self-reported depression after delivery 	Table 65
<ul style="list-style-type: none"> Self-reported lack of interest after delivery 	Table 66
<ul style="list-style-type: none"> Reported diagnosis of postpartum depression 	Table 67
<ul style="list-style-type: none"> Medication or counseling for postpartum depression treatment 	Table 68

Description	Table Number
Support	
<ul style="list-style-type: none"> • Support systems available to help mother 	Table 69
<ul style="list-style-type: none"> • Husband/partner engagement and emotional support 	Table 70
<ul style="list-style-type: none"> • Husband/partner engagement with baby 	Table 71
<ul style="list-style-type: none"> • Husband/partner emotionally supports mother 	Table 72
Health Insurance	
<ul style="list-style-type: none"> • Health insurance before pregnancy 	Table 73
<ul style="list-style-type: none"> • Health insurance for prenatal care 	Table 74
<ul style="list-style-type: none"> • Current health insurance 	Table 75
<ul style="list-style-type: none"> • Medicaid before pregnancy by demographics 	Table 76
<ul style="list-style-type: none"> • Medicaid for prenatal care by demographics 	Table 77
<ul style="list-style-type: none"> • Medicaid current health insurance by demographics 	Table 78
Drug Use	
<ul style="list-style-type: none"> • Marijuana use before or during pregnancy 	Table 79
<ul style="list-style-type: none"> • Prenatal care visit addresses marijuana use or drug treatment 	Table 80
<ul style="list-style-type: none"> • Marijuana use after delivery 	Table 81
<ul style="list-style-type: none"> • Marijuana use and breastfeeding 	Table 82
<ul style="list-style-type: none"> • Prescription antidepressant use during pregnancy 	Table 83
<ul style="list-style-type: none"> • Prescription pain reliever use during pregnancy 	Table 84

Table 1
Sample Numbers, Response Rates, and Estimated Populations by Selected Characteristics
2017 Illinois PRAMS


Characteristic	Number Sampled	Number of Respondents	Unweighted Percent Responding	Estimated Population*
All	2,235	1,224	54.7	141,656
Age				
< 20 years	117	44	37.6	4,974
20-24 years	423	208	49.2	24,816
25-29 years	578	317	54.8	37,795
30-34 years	677	396	58.5	43,979
=>35 years	440	259	58.9	30,124
Race				
NH White	1,073	639	59.6	76,068
NH Black	470	228	48.5	22,893
Hispanic	476	241	50.6	29,731
Other	194	103	53.1	11,241
Education				
< High School	265	144	54.3	17,479
High School	522	236	45.2	29,218
> High School	1,391	815	58.6	91,789
Marital Status				
Married	1,270	761	59.9	84,308
Other	965	463	48.0	57,381
Birthweight				
Low (<2,500 grams)	618	342	55.3	10,477
Normal (=>2,500 grams)	1,617	882	54.5	131,212

*Estimated population is the PRAMS eligible population of Illinois resident women, age 14 years or older, who gave live birth in Illinois. Total estimated population for some characteristics may not equal the total estimated population for the "All" population due to incomplete data for those characteristics.

Health Behaviors

Percentage (%) of recent mothers in Illinois who reported doing the following activities 12 months before pregnancy

Survey Question 4: At any time during the 12 months before you got pregnant with your new baby, did you do any of the following things? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
I was exercising 3 or more days of the week for fitness outside of my regular job.	40.6	37.6	43.7	1,223	57,426
I talked to a health care worker about my family medical history.	37.9	34.9	40.9	1,221	53,545
I was dieting (changing my eating habits) to lose weight.	27.7	25.0	30.5	1,221	39,105
A health care worker checked me for diabetes.	20.2	17.8	22.9	1,220	28,537
I was regularly taking prescription medicines other than birth control.	19.0	16.7	21.6	1,221	26,857

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds -- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated population affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage (%) of recent mothers in Illinois who reported dieting 12 months before pregnancy

Survey Question 4: At any time during the 12 months before you got pregnant with your new baby, did you do any of the following things?
Response: Yes A. I was dieting (changing my eating habits) to lose weight.



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	27.7	25.0	30.5	1,221	39,105
Maternal Age	<20 yrs	21.4	10.2	39.4	44	1,064
	20 to 24 yrs	23.2	17.3	30.3	207	5,712
	25 to 29 yrs	26.2	21.2	31.8	317	9,884
	30 to 34 yrs	32.4	27.5	37.6	395	14,187
	=>35 yrs	27.6	22.2	33.8	258	8,259
Race/Ethnicity	NH White	31.0	27.3	35.0	637	23,493
	NH Black	19.4	14.1	26.1	228	4,442
	Hispanic	26.8	21.1	33.3	240	7,901
	Other	25.5	17.3	35.9	103	2,868
Education	<12 yrs	15.8	10.2	23.8	143	2,731
	12 yrs	19.3	14.3	25.5	236	5,644
	>12 yrs	33.0	29.6	36.6	813	30,163
Birth Weight	Low	21.2	17.1	25.8	342	2,217
	Normal	28.2	25.4	31.3	879	36,888
Marital Status	Married	34.2	30.7	37.9	760	28,778
	Other	18.1	14.4	22.5	461	10,326
Insurer before Pregnant	Medicaid	22.0	17.4	27.4	335	8,465
	Other	29.8	26.7	33.2	886	30,640

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage (%) of recent mothers in Illinois who reported exercising three or more times per week during the 12 months before pregnancy

Survey Question 4: At any time during the 12 months before you got pregnant with your new baby, did you do any of the following things?

Response: Yes B. I was exercising three or more days of the week for fitness outside of my regular job.



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	40.6	37.6	43.7	1,223	57,426
Maternal Age	<20 yrs	27.9	15.0	45.9	44	1,388
	20 to 24 yrs	34.4	27.5	42.0	207	8,477
	25 to 29 yrs	36.5	30.9	42.5	317	13,789
	30 to 34 yrs	45.7	40.4	51.2	396	20,114
	=>35 yrs	45.3	38.8	52.0	259	13,658
Race/Ethnicity	NH White	44.3	40.2	48.4	638	33,597
	NH Black	26.3	20.2	33.6	228	6,031
	Hispanic	43.0	36.2	50.1	241	12,788
	Other	37.1	27.6	47.8	103	4,175
Education	<12 yrs	32.0	23.9	41.4	144	5,599
	12 yrs	29.3	23.2	36.3	236	8,566
	>12 yrs	45.9	42.2	49.6	814	42,033
Birth Weight	Low	37.8	32.9	43.1	342	3,964
	Normal	40.8	37.6	44.1	881	53,462
Marital Status	Married	47.2	43.4	51.0	761	39,806
	Other	30.8	26.2	35.9	462	17,620
Insurer before Pregnant	Medicaid	28.5	23.3	34.4	336	11,041
	Other	45.1	41.6	48.7	887	46,385

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Conditions

Percentage (%) of recent mothers in Illinois who reported a having diabetes, high blood pressure, or diabetes.

Survey Question 5: During the 3 months before you got pregnant with your new baby, did you have any of the following health conditions?

Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Depression	9.7	8.0	11.8	1,221	13,705
High blood pressure or hypertension	3.1	2.2	4.3	1,222	4,378
Type 1 or Type 2 diabetes	1.6	1.0	2.5	1,223	2,200

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage (%) of recent mothers in Illinois who reported taking a daily multivitamin with folic acid

Survey Question 6: During the month before you got pregnant with your new baby, how many times a week did you take a multivitamin, a prenatal vitamin, or a folic acid vitamin? Response: Every day of the week



	Percentage ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴		
All	Overall	36.9	34.0	39.9	1,220	52,093
Maternal Age	<20 yrs	7.0	2.1	20.6	44	348
	20 to 24 yrs	19.8	14.4	26.7	207	4,884
	25 to 29 yrs	31.5	26.2	37.3	317	11,901
	30 to 34 yrs	48.2	42.8	53.6	395	21,127
	=>35 yrs	46.2	39.7	52.8	257	13,833
Race/Ethnicity	NH White	47.0	42.8	51.2	637	35,591
	NH Black	24.5	18.4	31.8	227	5,594
	Hispanic	21.8	16.6	28.1	240	6,457
	Other	36.4	27.0	46.9	103	4,089
Education	<12 yrs	22.7	15.9	31.2	144	3,959
	12 yrs	21.3	15.9	27.9	235	6,190
	>12 yrs	44.7	41.0	48.4	812	40,851
Birth Weight	Low	37.1	32.2	42.3	341	3,869
	Normal	36.9	33.8	40.1	879	48,223
Marital Status	Married	48.8	45.0	52.6	759	41,019
	Other	19.4	15.6	23.9	461	11,074
Insurer before Pregnant	Medicaid	21.7	17.0	27.2	336	8,381
	Other	42.6	39.2	46.2	884	43,712

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage (%) of recent mothers in Illinois who had a visit with a health care worker in the 12 months before pregnancy

Survey Question 7: In the 12 months before you got pregnant with your new baby, did you have any health care visits with a doctor, nurse, or other health care worker, including a dental or mental health worker? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	70.3	67.3	73.1	1,207	98,245
Maternal Age	<20 yrs	57.2	39.5	73.3	44	2,847
	20 to 24 yrs	55.0	47.2	62.7	205	13,480
	25 to 29 yrs	67.1	61.0	72.7	313	24,990
	30 to 34 yrs	78.3	73.4	82.5	390	33,972
	=>35 yrs	77.4	71.2	82.6	255	22,956
Race/Ethnicity	NH White	79.7	75.9	83.0	632	59,981
	NH Black	67.7	59.8	74.6	227	15,380
	Hispanic	53.8	46.7	60.8	238	15,896
	Other	58.7	47.6	69.0	97	6,130
Education	<12 yrs	51.0	41.4	60.5	139	8,676
	12 yrs	53.4	46.0	60.6	234	15,507
	>12 yrs	79.6	76.4	82.5	805	72,016
Birth Weight	Low	73.2	68.2	77.6	338	7,579
	Normal	70.1	66.9	73.1	869	90,666
Marital Status	Married	77.0	73.6	80.1	749	63,839
	Other	60.5	55.2	65.6	458	34,406
Insurer before Pregnant	Medicaid	64.4	58.2	70.2	332	24,612
	Other	72.5	69.1	75.7	875	73,633

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition


Health Behaviors

Percentage (%) of recent mothers in Illinois who had the following types of health care worker visits during the 12 months before pregnancy

Among women reporting they had a visit with a health care worker in the 12 months before pregnancy

Survey Question 8: What type of health care worker visit did you have in the 12 months before you got pregnant with your new baby?

Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Teeth cleaned by dentist/dental hygienist	63.0	59.4	66.5	884	62,718
OB/GYN	59.0	55.3	62.5	884	58,724
Family doctor	51.6	47.9	55.3	884	51,383
Illness/chronic condition visit	16.1	13.6	19.0	884	16,070
Family planning visit	13.3	11.1	15.9	884	13,245
Depression/anxiety visit	9.1	7.1	11.4	884	9,016
Injury visit	3.8	2.6	5.5	884	3,790

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage (%) of recent mothers in Illinois who had a visit to have their teeth cleaned in the 12 months before pregnancy

Among women reporting they had a visit with a health care worker in the 12 months before pregnancy

Survey Question 8: In the 12 months before you got pregnant with your new baby, did you have any health care visits with a doctor, nurse, or other health care worker, including a dental or mental health worker? Response: G. Visit to have my teeth cleaned by a dentist or dental hygienist



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	63.0	59.4	66.5	884	62,718
Maternal Age	<20 yrs	--	--	--	--	--
	20 to 24 yrs	45.2	35.2	55.5	123	6,151
	25 to 29 yrs	62.2	54.9	68.9	219	15,790
	30 to 34 yrs	68.4	62.4	73.9	310	23,470
	=>35 yrs	67.3	60.0	73.9	206	15,761
Race/Ethnicity	NH White	68.3	63.8	72.5	524	41,271
	NH Black	52.6	43.1	61.9	159	8,090
	Hispanic	56.8	47.2	65.8	132	9,134
	Other	55.3	41.8	68.1	64	3,760
Education	<12 yrs	44.0	31.8	57.0	82	4,020
	12 yrs	38.7	29.6	48.6	130	6,049
	>12 yrs	71.0	67.1	74.7	656	51,657
Birth Weight	Low	56.3	50.1	62.3	253	4,314
	Normal	63.6	59.7	67.3	631	58,405
Marital Status	Married	68.1	64.0	72.0	594	44,256
	Other	53.4	46.6	60.0	290	18,462
Insurer before Pregnant	Medicaid	41.4	34.1	49.1	225	10,253
	Other	70.1	66.2	73.8	659	52,465

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage of recent mothers in Illinois who reported that during a pre-pregnancy health care visit a health care worker did any of the following things

Among women reporting they had a visit with a health care worker in the 12 months before pregnancy

Survey Question 9: During any of your health care visits in the 12 months before you got pregnant, did a doctor, nurse, or other health care worker do any of the following things? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Ask me if I was smoking cigarettes	76.2	73.0	79.2	870	74,629
Ask me about the kind of work I do	66.3	62.8	69.7	870	65,040
Ask me if I was feeling down or depressed	54.2	50.5	57.8	871	53,154
Talk to me about using birth control to prevent pregnancy	49.9	46.2	53.5	871	48,903
Ask me if someone was hurting me emotionally or physically	49.4	45.7	53.1	871	48,417
Talk to me about my desire to have or not have children	48.0	44.3	51.6	871	47,038
Talk to me about maintaining a healthy weight	45.4	41.7	49.1	408	44,295
Tell me to take a vitamin with folic acid	45.0	41.3	48.7	868	43,967
Test me for HIV (the virus that causes AIDS)	33.6	30.2	37.3	860	32,522
Talk to me about sexually transmitted infections such as chlamydia, gonorrhea, or syphilis	32.8	29.4	36.4	871	32,147
Talk to me about how I could improve my health before a pregnancy	31.6	28.2	35.1	867	30,840
Talk to me about controlling any medical conditions such as diabetes or high blood pressure	20.8	18.0	24.0	869	20,372

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage (%) of recent mothers in Illinois who had a health care worker talk to them about preparing for a pregnancy

Survey Question 10: Before you got pregnant with your new baby, did a doctor, nurse, or other health care worker talk to you about preparing for a pregnancy? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	28.7	26.0	31.6	1,204	40,154
Maternal Age	<20 yrs	14.9	6.1	32.2	44	741
	20 to 24 yrs	21.0	15.3	28.0	205	5,138
	25 to 29 yrs	21.8	17.2	27.1	313	8,139
	30 to 34 yrs	37.0	32.0	42.4	391	16,158
	=>35 yrs	34.0	28.0	40.5	251	9,977
Race/Ethnicity	NH White	33.9	30.1	38.0	630	25,536
	NH Black	17.4	12.3	24.2	227	3,969
	Hispanic	28.5	22.5	35.3	234	8,316
	Other	21.1	13.9	30.9	100	2,293
Education	<12 yrs	24.1	16.8	33.2	138	4,087
	12 yrs	18.1	13.2	24.4	234	5,252
	>12 yrs	33.2	29.8	36.8	803	30,145
Birth Weight	Low	28.1	23.6	33.1	334	2,878
	Normal	28.8	25.9	31.9	870	37,275
Marital Status	Married	37.3	33.6	41.0	746	30,862
	Other	16.3	12.7	20.7	458	9,292
Insurer before Pregnant	Medicaid	21.0	16.3	26.5	335	8,079
	Other	31.7	28.5	35.1	869	32,075

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage (%) of recent mothers in Illinois who reported that during a health care visit a health care worker talked to them about the following things in preparation for a pregnancy

Among women reporting that before pregnancy a health care worker talked to them about preparing for pregnancy

Survey Question 11: Before you got pregnant with your new baby, did a doctor, nurse, or other health care worker talk with you about any of the things listed below about preparing for a pregnancy? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
How drinking alcohol during pregnancy can affect a baby	58.7	53.1	64.1	368	23,856
The safety of using prescription or over-the-counter medicines during pregnancy	61.1	55.5	66.4	367	24,790
Getting my vaccines updated before pregnancy	51.7	46.0	57.3	366	20,988
How smoking during pregnancy can affect a baby	57.3	51.7	62.8	367	23,179
How using illegal drugs during pregnancy can affect a baby	54.3	48.7	59.8	368	22,059
Visiting a dentist or dental hygienist before pregnancy	41.0	35.5	46.6	366	16,609
Getting counseling for any genetic diseases that run in my family	39.0	33.5	44.6	365	15,746
Getting counseling or treatment for depression or anxiety	24.8	20.1	30.1	366	10,040

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage (%) of recent mothers in Illinois who reported that during a health care visit a health care worker talked to them about visiting a dentist or dental hygienist in preparation for a pregnancy
 Among women reporting that before pregnancy a health care worker talked to them about preparing for pregnancy

Survey Question 11: Before you got pregnant with your new baby, did a doctor, nurse, or other health care worker talk with you about any of the things listed below about preparing for a pregnancy? 11b. Visiting a dentist or dental hygienist before pregnancy. Response: Yes



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	41.0	35.5	46.6	366	16,609
Maternal Age	<20 yrs	--	--	--	--	--
	20 to 24 yrs	55.5	39.1	70.7	44	2,995
	25 to 29 yrs	43.7	31.7	56.5	73	3,564
	30 to 34 yrs	37.4	29.3	46.3	148	6,106
	=>35 yrs	36.4	26.5	47.7	95	3,636
Race/Ethnicity	NH White	33.7	27.4	40.6	231	8,724
	NH Black	57.2	38.9	73.8	42	2,329
	Hispanic	51.2	38.1	64.1	66	4,177
	Other	--	--	--	--	--
Education	<12 yrs	63.9	44.0	79.9	31	2,631
	12 yrs	52.5	35.8	68.8	40	2,747
	>12 yrs	36.2	30.4	42.5	289	11,067
Birth Weight	Low	37.8	28.9	47.7	101	1,104
	Normal	41.2	35.4	47.3	265	15,505
Marital Status	Married	39.0	33.1	45.2	295	12,220
	Other	47.5	34.7	60.6	71	4,389
Insurer before Pregnant	Medicaid	48.2	35.0	61.6	66	3,939
	Other	39.1	33.2	45.4	300	12,670

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Health Behaviors

Percentage (%) of recent mothers in Illinois who reported that during a health care visit a health care worker talked to them about getting counseling or treatment for depression or anxiety in preparation for a pregnancy
 Among women reporting that before pregnancy a health care worker talked to them about preparing for pregnancy

Survey Question 11: Before you got pregnant with your new baby, did a doctor, nurse, or other health care worker talk with you about any of the things listed below about preparing for a pregnancy? 11d. Getting counseling or treatment for depression or anxiety. Response: Yes



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	24.8	20.1	30.1	366	10,040
Maternal Age	<20 yrs	--	--	--	--	--
	20 to 24 yrs	30.3	17.4	47.3	44	1,638
	25 to 29 yrs	28.3	18.0	41.4	73	2,310
	30 to 34 yrs	18.5	12.5	26.5	148	3,014
	=>35 yrs	24.2	15.9	35.1	95	2,416
Race/Ethnicity	NH White	17.4	12.7	23.5	231	4,505
	NH Black	46.7	29.4	64.9	42	1,903
	Hispanic	39.9	27.8	53.5	66	3,259
	Other	--	--	--	--	--
Education	<12 yrs	54.0	35.1	71.7	31	2,222
	12 yrs	44.3	28.5	61.2	40	2,314
	>12 yrs	16.8	12.6	22.2	289	5,141
Birth Weight	Low	23.1	15.8	32.3	101	673
	Normal	24.9	19.9	30.6	265	9,366
Marital Status	Married	18.6	14.2	23.9	295	5,817
	Other	45.7	33.0	59.0	71	4,223
Insurer before Pregnant	Medicaid	39.7	27.3	53.6	66	3,250
	Other	21.0	16.3	26.6	300	6,790

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Family Planning

Percentage (%) of recent mothers in Illinois who reported the following feelings about being pregnant

Survey Question 15: Thinking back to just before you got pregnant with your new baby, how did you feel about becoming pregnant?

Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
I wanted to be pregnant then	42.2	39.1	45.3	1,216	59,384
I wanted to be pregnant later	18.7	16.3	21.3	1,216	26,272
I wanted to be pregnant sooner	15.0	13.0	17.3	1,216	21,127
I wasn't sure what I wanted	15.7	13.5	18.2	1,216	22,152
I didn't want to be pregnant then or at any time in the future	8.4	6.8	10.4	1,216	11,818

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Family Planning

Percentage (%) of recent mothers in Illinois who reported that pregnancy was unintended

Survey Question 15: Thinking back to just before you got pregnant with your new baby, how did you feel about becoming pregnant? A. I want to be pregnant later; B. I wanted to be pregnant sooner; C. I want to be pregnant then; D. I didn't want to be pregnant then or at any time in the future; E. I wasn't sure what I wanted. Responses B and C were grouped as 'Intended Pregnancy'; Responses A, D, and E were grouped as 'Unintended Pregnancy'. Chart shows unintended pregnancy rates.



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	42.8	39.7	45.9	1,216	60,241
Maternal Age	<20 yrs	83.0	65.5	92.7	44	4,131
	20 to 24 yrs	55.8	48.0	63.3	208	13,835
	25 to 29 yrs	43.8	37.9	50.0	314	16,481
	30 to 34 yrs	33.4	28.3	38.8	392	14,471
	=>35 yrs	37.8	31.5	44.4	258	11,323
Race/Ethnicity	NH White	34.0	30.1	38.2	636	25,723
	NH Black	69.4	61.7	76.1	228	15,881
	Hispanic	48.4	41.4	55.4	237	14,185
	Other	31.7	22.6	42.4	102	3,523
Education	<12 yrs	60.5	50.9	69.3	140	10,267
	12 yrs	56.8	49.5	63.8	236	16,586
	>12 yrs	35.1	31.6	38.8	811	32,042
Birth Weight	Low	50.6	45.3	55.8	340	5,260
	Normal	42.2	38.9	45.5	876	54,981
Marital Status	Married	27.7	24.4	31.2	756	23,173
	Other	64.9	59.7	69.8	460	37,069
Insurer before Pregnant	Medicaid	62.6	56.4	68.4	335	24,187
	Other	35.3	31.9	38.9	881	36,054

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Family Planning

Percentage (%) of recent mothers in Illinois who reported that they were trying to get pregnant

Survey Question 16: When you got pregnant with your new baby, were you trying to get pregnant? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	56.0	52.9	59.1	1,216	78,882
Maternal Age	<20 yrs	20.9	9.8	39.1	44	1,040
	20 to 24 yrs	40.0	32.7	47.8	207	9,874
	25 to 29 yrs	56.4	50.3	62.4	316	21,308
	30 to 34 yrs	65.8	60.4	70.9	391	28,576
	=>35 yrs	60.3	53.6	66.6	258	18,084
Race/Ethnicity	NH White	64.5	60.3	68.4	636	48,775
	NH Black	27.6	21.1	35.1	228	6,308
	Hispanic	55.2	48.1	62.1	240	16,338
	Other	62.3	51.3	72.1	99	6,792
Education	<12 yrs	42.0	33.0	51.6	142	7,229
	12 yrs	41.6	34.7	48.9	236	12,159
	>12 yrs	63.9	60.2	67.5	810	58,393
Birth Weight	Low	46.9	41.7	52.2	340	4,887
	Normal	56.7	53.4	60.0	876	73,995
Marital Status	Married	71.8	68.3	75.1	753	59,938
	Other	33.0	28.2	38.2	463	18,944
Insurer before Pregnant	Medicaid	34.9	29.2	41.1	335	13,454
	Other	64.0	60.4	67.4	881	65,428

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Family Planning

Percentage (%) of recent mothers in Illinois who were not trying to get pregnant and who reported using birth control

Among women who were not trying to get pregnant

Survey Question 17: When you got pregnant with your new baby, were you or your husband or partner doing anything to keep from getting pregnant? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	41.7	37.0	46.5	526	25,499
Maternal Age	<20 yrs	64.5	44.5	80.5	35	2,513
	20 to 24 yrs	40.7	31.2	50.9	128	6,022
	25 to 29 yrs	41.3	32.3	50.8	139	6,791
	30 to 34 yrs	37.7	28.9	47.5	124	5,446
	=>35 yrs	40.7	30.6	51.7	100	4,726
Race/Ethnicity	NH White	44.3	37.2	51.6	215	11,728
	NH Black	33.9	25.6	43.3	164	5,502
	Hispanic	46.2	35.8	56.9	104	6,126
	Other	42.4	26.3	60.4	36	1748.2
Education	<12 yrs	40.4	28.6	53.4	82	3,978
	12 yrs	39.1	30.2	48.8	139	6,654
	>12 yrs	44.6	38.4	51.0	287	14,440
Birth Weight	Low	35.3	28.5	42.6	175	1,935
	Normal	42.3	37.2	47.6	351	23,563
Marital Status	Married	42.3	35.3	49.6	208	9,718
	Other	41.3	35.1	47.7	318	15,781
Insurer before Pregnant	Medicaid	40.7	33.4	48.5	228	10,200
	Other	42.3	36.3	48.6	298	15,299

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition


Family Planning

Percentage (%) of recent mothers in Illinois who reported the following reasons for not using birth control

Among women who were not trying to get pregnant

Survey Question 18: What were your reasons or your husband's or partner's reasons for not doing anything to keep from getting pregnant?

Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
I didn't mind if I got pregnant	39.2	33.4	45.3	329	14,560
I thought I could not get pregnant at that time	19.4	14.9	24.8	330	7,210
I had side effects from the birth control method I was using	16.4	12.2	21.6	330	6,085
My husband or partner didn't want to use anything	10.8	7.5	15.4	330	4,023
I forgot to use a birth control method	6.0	3.6	9.6	329	2,218
I thought my husband or partner or I was sterile	5.5	3.3	9.1	330	2,062
I had problems getting birth control when I needed it	3.6	2.0	6.5	330	1,333

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Prenatal Care

Percentage (%) of recent mothers in Illinois who reported beginning prenatal care in the first trimester

Survey Question 19: How many weeks or months pregnant were you when you had your first visit for prenatal care? Response: First trimester care



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	88.4	86.2	90.3	1,205	123,636
Maternal Age	<20 yrs	52.9	35.4	69.7	43	2,613
	20 to 24 yrs	90.6	85.2	94.1	203	22,183
	25 to 29 yrs	85.7	80.8	89.5	315	32,106
	30 to 34 yrs	91.6	87.7	94.3	389	39,490
	=>35 yrs	91.4	86.8	94.5	255	27,245
Race/Ethnicity	NH White	92.8	90.2	94.8	631	69,867
	NH Black	81.1	74.2	86.6	223	18,232
	Hispanic	80.9	74.6	86.0	237	23,690
	Other	92.8	84.6	96.8	101	10,287
Education	<12 yrs	77.7	68.6	84.8	137	13,045
	12 yrs	81.4	75.1	86.5	231	23,445
	>12 yrs	92.9	90.7	94.6	808	84,565
Birth Weight	Low	84.8	80.4	88.3	334	8,682
	Normal	88.7	86.3	90.7	871	114,954
Marital Status	Married	92.6	90.4	94.4	752	77,080
	Other	82.2	77.8	86.0	453	46,557
Insurer Prenatal Care	Medicaid	83.3	79.3	86.6	527	51,911
	Other	93.0	90.5	94.9	661	70,248

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Prenatal Care

Percentage (%) of recent mothers in Illinois who reported they did not receive prenatal care as early as desired

Among women who received prenatal care

Survey Question 20: Did you get prenatal care as early in your pregnancy as you wanted?

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
No	10.2	8.5	12.3	1,198	14,192
Yes	89.8	87.7	91.5	1,198	124,492

Reported reasons for not getting early prenatal care among recent mothers who reported not getting prenatal care as early as desired

Among women who received prenatal care

Survey Question 21: Did any of these things keep you from getting prenatal care when you wanted it? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
I didn't know that I was pregnant	32.8	24.4	42.6	140	4,805
I couldn't get an appointment when I wanted one	33.7	25.1	43.4	140	4,928
I didn't have my Medicaid card	18.0	11.6	27.0	140	2,639
I didn't have enough money or insurance to pay for the visit	21.5	14.6	30.4	139	3,116
The doctor or health plan would not start care as early as I wanted	15.2	9.7	23.0	140	2,228
I didn't want anyone else to know I was pregnant	8.0	4.0	15.2	139	1,155
I had too many other things going on	11.8	6.6	20.1	139	1,712
I didn't have any transportation to get to the clinic or doctor's office	4.9	1.9	12.0	139	717
I couldn't take off from work or school	3.1	1.0	9.2	138	446
I didn't have anyone to take care of my children	0.4	0.1	1.7	138	63
I didn't want prenatal care	3.7	1.2	10.9	138	527

¹Percentage = Weighted percentage²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Prenatal Care

Percentage (%) of recent mothers in Illinois who received adequate prenatal care based on the Kotelchuck Index

Among women who received prenatal care

Inadequate <50% of expected visits, Intermediate 50%-79%, Adequate 80%-109%, Adequate Plus 110% or more. Adequate is a combination of Adequate and Adequate Plus.



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	78.2	75.4	80.7	1,159	105,350
	Maternal Age					
	<20 yrs	79.1	61.4	90.0	43	3,901
	20 to 24 yrs	75.2	67.7	81.4	196	17,889
	25 to 29 yrs	72.5	66.6	77.8	306	26,342
	30 to 34 yrs	83.8	79.2	87.6	377	35,310
	=>35 yrs	79.4	73.0	84.5	237	21,909
Race/Ethnicity	NH White	84.5	81.1	87.4	615	62,046
	NH Black	62.2	54.1	69.8	216	13,588
	Hispanic	74.3	67.3	80.3	221	20,377
	Other	77.3	66.8	85.2	94	7,985
Education	<12 yrs	62.4	52.6	71.3	138	10,559
	12 yrs	77.0	69.9	82.8	217	20,782
	>12 yrs	81.7	78.5	84.5	775	71,633
Birth Weight	Low	76.6	71.5	81.0	319	7,434
	Normal	78.3	75.3	81.0	840	97,916
Marital Status	Married	84.2	81.1	86.8	717	67,037
	Other	69.5	64.3	74.2	442	38,313
Insurer Prenatal Care	Medicaid	69.8	65.0	74.2	508	41,988
	Other	85.2	81.9	88.0	634	61,924

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds


-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Prenatal Care

Percentage (%) of recent mothers in Illinois who reported using WIC services during pregnancy. Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	33.9	30.9	37.1	1,186	46,461
Maternal Age	<20 yrs	63.4	45.5	78.3	43	3,129
	20 to 24 yrs	55.8	47.9	63.4	202	13,252
	25 to 29 yrs	37.3	31.4	43.5	310	13,676
	30 to 34 yrs	21.8	17.5	27.0	381	9,302
	=>35 yrs	24.5	19.1	30.9	250	7,101
Race/Ethnicity	NH White	20.4	17.0	24.2	628	15,212
	NH Black	58.9	50.9	66.4	220	13,087
	Hispanic	54.2	47.0	61.3	229	15,120
	Other	21.8	14.0	32.4	98	2,348
Education	<12 yrs	69.3	60.1	77.3	139	11,677
	12 yrs	58.7	51.2	65.7	229	16,504
	>12 yrs	19.4	16.6	22.7	793	17,391
Birth Weight	Low	39.9	34.8	45.3	333	4,073
	Normal	33.5	30.2	36.8	853	42,388
Marital Status	Married	15.7	13.1	18.7	740	12,865
	Other	61.2	55.9	66.3	446	33,596
Insurer Prenatal Care	Medicaid	65.9	61.1	70.4	517	40,115
	Other	7.6	5.6	10.3	652	5,653

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Prenatal Care

Percentage (%) of recent mothers in Illinois who reported the following topics were discussed during prenatal care visits

Among women who received prenatal care

Survey Question 23: During any of your prenatal care visits, did a doctor, nurse, or other health care worker talk with you about any of the things listed below? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Doing tests to screen for birth defects or diseases that run in my family	91.9	90.0	93.4	1,203	128,179
Medicines that are safe to take during my pregnancy	89.9	87.8	91.6	1,205	125,490
Breastfeeding my baby	83.6	81.2	85.7	1,204	116,593
The signs and symptoms of preterm labor (labor more than 3 week sbefore the baby is due)	83.2	80.7	85.3	1,204	116,023
What to do if I feel depressed during my pregnancy or after my baby is born	75.0	72.2	77.6	1,201	104,542
How drinking alcohol during pregnancy could affect my baby	75.0	72.3	77.6	1,203	104,600
How smoking during pregnancy could affect my baby	73.2	70.4	75.9	1,203	102,171
How using illegal drugs could affect my baby	66.2	63.2	69.0	1,204	92,342
Using a seal belt during my pregnancy	59.6	56.5	62.7	1,201	83,032
Physical abuse to women by their husbands or partners	53.7	50.6	56.8	1,200	74,855

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Prenatal Care

Percentage (%) of recent mothers in Illinois who reported a health care worker asked about the following topics during prenatal care visits

Among women who received prenatal care

Survey Question 24: During any of your prenatal care visits, did a doctor, nurse, or other health care worker ask you any of the things listed below?

Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
If I was taking any prescription medication	94.7	93.0	95.9	1,206	132,172
If I was smoking cigarettes	93.5	91.7	94.9	1,205	130,483
If I was drinking alcohol	92.5	90.7	94.0	1,206	129,208
If I planned to breastfeed my new baby	90.9	88.9	92.5	1,205	126,748
If I planned to use birth control after my baby was born	86.9	84.7	88.8	1,205	121,183
If I was feeling down or depressed	81.9	79.4	84.2	1,205	114,277
If I was using drugs such as marijuana, cocaine, crack, or meth	75.0	72.3	77.6	1,206	104,768
If someone was hurting me emotionally or physically	69.1	66.1	71.9	1,204	96,237
If I knew how much weight I should gain during pregnancy	64.3	61.2	67.2	1,202	89,446
If I wanted to be tested for HIV (the virus that causes AIDS)	60.4	57.3	63.4	1,197	83,785

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Prenatal Care

Percentage (%) of recent mothers in Illinois who reported they have heard or read about folic acid preventing birth defects

Survey Question 25: Have you ever heard or read that taking a vitamin with folic acid can help prevent some birth defects? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
No	25.2	22.5	28.0	1,215	35,307
Yes	74.8	72.0	77.5	1,215	104,990

Reported sources of folic acid information

Among women who reported hearing or reading about folic acid

Survey Question 26: Have you ever heard about folic acid from any of the following? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Doctor, nurse, or other health care worker	81.7	78.7	84.3	919	86,488
Family or friends	39.6	36.1	43.1	919	41,896
Magazine or newspaper article	30.0	26.8	33.3	919	31,764
Book	29.9	26.7	33.2	919	31,639
Radio or television	12.0	9.8	14.6	919	12,717

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Prenatal Care

Percentage (%) of recent mothers in Illinois who reported having their teeth cleaned during their pregnancy

Survey Question 29: During your most recent pregnancy, did you have your teeth cleaned by a dentist or dental hygienist? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	44.5	41.4	47.6	1,216	62,608
Maternal Age	<20 yrs	20.8	10.2	37.7	43	1,026
	20 to 24 yrs	31.5	24.8	39.0	205	7,692
	25 to 29 yrs	38.3	32.6	44.3	317	14,464
	30 to 34 yrs	53.8	48.3	59.1	393	23,426
	=>35 yrs	53.4	46.7	60.0	258	15,999
Race/Ethnicity	NH White	52.7	48.4	56.8	636	39,796
	NH Black	24.7	18.6	31.9	226	5,626
	Hispanic	42.3	35.6	49.4	239	12,477
	Other	37.6	28.1	48.3	102	4,182
Education	<12 yrs	27.9	20.3	37.0	141	4,779
	12 yrs	28.7	22.6	35.6	234	8,331
	>12 yrs	53.2	49.4	56.9	812	48,576
Birth Weight	Low	40.6	35.5	45.9	340	4,224
	Normal	44.8	41.5	48.1	876	58,384
Marital Status	Married	57.2	53.3	60.9	757	47,879
	Other	25.9	21.5	30.7	459	14,729
Insurer Prenatal Care	Medicaid	31.1	26.8	35.7	535	19,602
	Other	56.0	51.8	60.0	664	42,338

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Prenatal Care

Percentage (%) of recent mothers in Illinois who reported health conditions during pregnancy

Survey Question 30: During your most recent pregnancy, did you have any of the following health conditions? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
High blood pressure (that started during this pregnancy), pre-eclampsia or eclampsia	11.7	9.9	13.8	1,210	16,369
Depression	9.9	8.2	12.0	1,210	13,904
Gestational diabetes (diabetes that started during this pregnancy)	8.8	7.2	10.7	1,214	12,395

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Immunizations

Percentage (%) of recent mothers in Illinois who reported being informed to get an influenza vaccination

Survey Question 27: During the 12 months before the delivery of your new baby, did a doctor, nurse, or other health care worker offer you a flu shot or tell you to get one? Response: Yes



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	86.3	84.0	88.3	1,203	120,371
Maternal Age	<20 yrs	73.0	55.1	85.6	43	3,606
	20 to 24 yrs	79.9	72.9	85.5	202	19,189
	25 to 29 yrs	85.4	80.6	89.2	312	31,925
	30 to 34 yrs	92.4	89.2	94.8	388	39,763
	=>35 yrs	86.0	80.6	90.1	258	25,889
Race/Ethnicity	NH White	88.8	85.8	91.2	635	66,939
	NH Black	80.1	73.2	85.6	225	18,156
	Hispanic	84.1	78.1	88.7	232	24,343
	Other	89.0	80.6	94.1	100	9,651
Education	<12 yrs	76.1	67.0	83.4	138	12,878
	12 yrs	80.6	74.2	85.6	232	23,136
	>12 yrs	90.1	87.6	92.1	806	81,784
Birth Weight	Low	76.5	71.7	80.8	335	7,843
	Normal	87.1	84.6	89.2	868	112,528
Marital Status	Married	89.6	87.1	91.7	745	74,108
	Other	81.5	77.1	85.2	458	46,264
Insurer Prenatal Care	Medicaid	81.6	77.6	85.0	526	50,657
	Other	90.7	88.0	92.8	660	68,325

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents= The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Immunizations

Percentage (%) of recent mothers in Illinois who reported receiving an influenza vaccination

Survey Question 28: During the 12 months before the delivery of your new baby, did you get a flu shot? Responses: 'Yes, before my pregnancy' and 'Yes, during my pregnancy' are combined and shown in chart



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	58.3	55.2	61.4	1,213	81,740
Maternal Age	<20 yrs	38.9	23.5	56.9	44	1,935
	20 to 24 yrs	55.9	48.0	63.5	203	13,478
	25 to 29 yrs	56.2	50.0	62.1	315	21,098
	30 to 34 yrs	62.5	57.1	67.7	392	27,172
	=>35 yrs	59.9	53.3	66.2	259	18,057
Race/Ethnicity	NH White	62.6	58.5	66.6	636	47,347
	NH Black	40.2	32.7	48.1	226	9,173
	Hispanic	56.7	49.5	63.6	237	16,473
	Other	72.8	62.4	81.2	102	8,084
Education	<12 yrs	51.1	41.5	60.6	139	8,625
	12 yrs	52.5	45.2	59.7	234	15,164
	>12 yrs	62.2	58.5	65.8	812	56,857
Birth Weight	Low	55.8	50.5	61.0	340	5,805
	Normal	58.5	55.1	61.8	873	75,935
Marital Status	Married	65.8	62.1	69.3	755	54,966
	Other	47.2	42.0	52.5	458	26,774
Insurer Prenatal Care	Medicaid	48.4	43.5	53.2	532	30,278
	Other	66.7	62.7	70.5	664	50,449

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Smoking

Percentage (%) of recent mothers in Illinois who reported smoking cigarettes in the past two years

Survey Question 31: Have you smoked any cigarettes in the past 2 years? Response: Yes



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	19.0	16.7	21.7	1,204	26,547
Maternal Age	<20 yrs	25.3	12.9	43.7	44	1,261
	20 to 24 yrs	28.0	21.5	35.5	205	6,890
	25 to 29 yrs	22.8	17.9	28.5	311	8,432
	30 to 34 yrs	14.3	10.8	18.6	388	6,142
	=>35 yrs	12.8	9.0	17.8	256	3,822
Race/Ethnicity	NH White	24.4	20.9	28.3	630	18,354
	NH Black	17.6	12.5	24.1	226	3,989
	Hispanic	11.0	7.3	16.3	236	3,213
	Other	9.3	4.4	18.5	99	991
Education	<12 yrs	25.3	17.9	34.5	140	4,342
	12 yrs	27.5	21.5	34.5	235	7,996
	>12 yrs	15.6	13.1	18.6	800	14,082
Birth Weight	Low	26.2	21.8	31.2	337	2,709
	Normal	18.5	15.9	21.3	867	23,837
Marital Status	Married	12.5	10.2	15.2	744	10,288
	Other	28.5	24.0	33.5	460	16,259
Insurer before Pregnant	Medicaid	31.5	26.0	37.5	335	12,172
	Other	14.3	11.9	17.0	869	14,374

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Smoking

Percentage (%) of recent mothers in Illinois who reported smoking cigarettes in the three months before pregnancy

Among women who reported cigarette smoking in the past two years

Survey Question 32: In the 3 months before you got pregnant, how many cigarettes did you smoke on an average day? Responses 'Yes/No' created using numeric response



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	16.1	13.9	18.6	1,201	22,355
Maternal Age	<20 yrs	22.3	10.7	40.7	43	1,064
	20 to 24 yrs	25.1	18.9	32.4	205	6,170
	25 to 29 yrs	19.1	14.6	24.6	309	7,011
	30 to 34 yrs	12.4	9.1	16.5	387	5,315
	=>35 yrs	9.3	6.2	13.8	257	2,795
Race/Ethnicity	NH White	21.3	17.9	25.1	627	15,899
	NH Black	15.8	11.1	22.1	226	3,598
	Hispanic	7.0	4.1	11.6	236	2,032
	Other	7.8	3.3	17.0	98	826
Education	<12 yrs	21.8	15.0	30.7	139	3,688
	12 yrs	22.8	17.2	29.6	233	6,573
	>12 yrs	13.3	10.9	16.1	800	11,968
Birth Weight	Low	23.2	19.0	28.0	336	2,392
	Normal	15.5	13.2	18.2	865	19,963
Marital Status	Married	9.4	7.4	11.9	742	7,757
	Other	25.7	21.4	30.6	459	14,599
Insurer before Pregnant	Medicaid	29.1	23.8	35.1	332	11,118
	Other	11.1	9.0	13.7	869	11,238

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Smoking


Percentage (%) of recent mothers in Illinois who reported smoking cigarettes in the last three months of pregnancy

Among women who reported cigarette smoking in the past two years

Survey Question 33: In the last 3 months of your pregnancy, how many cigarettes did you smoke on an average day?

Responses

'Yes/No' created using numeric response



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	6.8	5.3	8.6	1,202	9,414
	Maternal Age					
	<20 yrs	12.4	4.1	31.7	43	591
	20 to 24 yrs	8.2	4.7	13.8	205	2,014
	25 to 29 yrs	9.5	6.4	13.9	309	3,483
	30 to 34 yrs	5.4	3.3	8.5	388	2,310
	=>35 yrs	3.4	1.8	6.4	257	1,016
Race/Ethnicity	NH White	9.7	7.4	12.7	627	7,275
	NH Black	6.4	3.6	11.1	226	1,454
	Hispanic	0.5	0.1	2.9	237	158
	Other	4.9	1.6	14.3	99	527
Education	<12 yrs	9.2	5.2	15.9	139	1,560
	12 yrs	12.2	8.0	18.1	233	3,515
	>12 yrs	4.7	3.3	6.6	801	4,245
Birth Weight	Low	12.0	9.0	16.0	337	1,244
	Normal	6.3	4.8	8.3	865	8,171
Marital Status	Married	2.8	1.8	4.3	743	2,265
	Other	12.6	9.5	16.5	459	7,150
Insurer Prenatal Care	Medicaid	12.8	9.9	16.4	533	8,028
	Other	1.9	1.0	3.5	653	1,386

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Smoking

Percentage (%) of recent mothers in Illinois who reported smoking cigarettes now

Among women who reported cigarette smoking in the past two years

Survey Question 36: How many cigarettes do you smoke on an average day now? Responses 'Yes/No' created using numeric response



	Percentage ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴		
All	Overall	11.0	9.2	13.2	1,202	15,319
Maternal Age	<20 yrs	17.3	7.1	36.6	43	827
	20 to 24 yrs	18.0	12.8	24.7	204	4,384
	25 to 29 yrs	12.9	9.2	17.7	310	4,729
	30 to 34 yrs	8.3	5.7	11.9	388	3,575
	=>35 yrs	6.0	3.6	9.8	257	1,804
Race/Ethnicity	NH White	14.5	11.7	17.9	627	10,818
	NH Black	11.8	7.9	17.5	226	2,691
	Hispanic	3.9	2.0	7.7	237	1,149
	Other	6.2	2.3	15.5	99	661
Education	<12 yrs	12.1	7.4	19.2	139	2,043
	12 yrs	20.6	15.2	27.2	233	5,917
	>12 yrs	8.0	6.2	10.4	803	7,233
Birth Weight	Low	19.3	15.5	23.9	337	1,994
	Normal	10.4	8.4	12.7	865	13,325
Marital Status	Married	6.3	4.7	8.4	744	5,207
	Other	17.9	14.2	22.2	458	10,112
Insurer Now	Medicaid	21.2	17.2	25.8	452	11,100
	Other	4.9	3.4	6.9	750	4,219

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Smoking

Reported number of cigarettes smoked daily during specified time periods

Among women who reported cigarette smoking in the past two years



Note: Confidence intervals (CI) and other details below

Smoking

Reported number of cigarettes smoked daily during specified time periods

Among women who reported cigarette smoking in the past two years

	Percentage ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴
Three Months Before Pregnancy, cigarettes/day*				
<1	6.0	3.3 10.8	203	1,346
1-5	36.4	29.0 44.4	203	8,127
6-10	34.5	27.3 42.6	203	7,722
11-20	19.1	13.6 26.1	203	4,266
>20	4.0	1.8 8.5	203	893
Last Three Months of Pregnancy, cigarettes/day**				
<1	5.6	1.9 15.2	89	524
1-5	54.9	42.5 66.7	89	5,169
6-10	22.5	14.3 33.6	89	2,119
11-20	15.0	7.9 26.7	89	1,413
>20	2.0	0.5 8.1	89	190
Now (at time of survey completion), cigarettes/day***				
<1	12.8	7.7 20.6	148	1,968
1 - 5	37.4	28.6 47.0	148	5,726
6 - 10	30.2	22.1 39.7	148	4,625
11 -20	16.3	10.4 24.5	148	2,492
>20	3.3	1.1 9.4	148	507

¹Percentage = Weighted percentage²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

*Survey Question 32: In the three months before you got pregnant, how many cigarettes did you smoke on an average day?

**Survey Question 33: In the last 3 months of your pregnancy, how many cigarettes did you smoke on an average day?

***Survey Question 36: How many cigarettes do you smoke on an average day now?

Smoking

Percentage (%) of recent mothers in Illinois who reported a health care worker advised them to quit smoking

Among women who reported cigarette smoking in the three months before pregnancy and who had a prenatal visit

Survey Question 34: During any of your prenatal care visits, did a doctor, nurse, or other health care worker advise you to quit smoking?

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
No	28.3	21.4	36.4	189	5,866
Yes	71.7	63.6	78.6	189	14,880

Percentage (%) of recent mothers in Illinois reporting that a health care worker did any of the following things to address quitting smoking

Among women who reported cigarette smoking in the three months before pregnancy and who had a prenatal visit

Survey Question 35: Listed below are some things about quitting smoking that a doctor, nurse, or other health care worker might have done during any of your prenatal care visits. Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Spend time with me discussing how to quit smoking	42.6	34.7	51.0	189	8,888
Provide me with booklets, videos, or other materials to help me quit smoking on my own	32.5	25.2	40.8	188	6,768
Ask if a family member or friend would support my decision to quit	20.0	14.1	27.4	188	4,157
Suggest that I set a specific date to stop smoking	19.3	13.6	26.6	188	4,012
Recommend using a nicotine patch	14.4	9.5	21.1	188	2,991
Recommend using nicotine gum	11.9	7.6	18.3	188	2,478
Suggest I attend a class or program to stop smoking	9.8	5.8	16.0	189	2,046
Refer me to a national or state quit line	9.6	5.8	15.6	188	2,007
Refer me to counseling for help with quitting	5.1	2.5	9.9	188	1,054
Prescribe a nicotine nasal spray or nicotine inhaler	1.7	0.5	5.4	188	358
Prescribe a pill like Zyban® (also known as Wellbutrin® or bupropion) to help me quit	1.1	0.2	5.8	187	237
Prescribe a pill like Chantix® (also known as varenicline) to help me quit	0.2	0.0	1.0	187	32

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Other Tobacco Products

Percentage (%) of recent mothers in Illinois who reported using other tobacco products in the past two years

Survey Question 37: Have you used any of the following products in the past 2 years? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
E-cigarettes or other electronic nicotine products	6.0	4.6	7.8	1,209	8,426
Hookah	3.5	2.5	4.8	1,206	4,823
Cigars, cigarillos, or little filtered cigars	3.0	2.0	4.3	1,209	4,138
Chewing tobacco, snuff, snus, or dip	0.5	0.2	1.3	1,206	753

Percentage (%) of recent mothers in Illinois who reported using other tobacco products in the three months before pregnancy or the last three months of pregnancy

Among women who reported using other tobacco products in the past two years

Survey Question 38: During the 3 months before you got pregnant, on average, how often did you use e-cigarettes or other electronic nicotine products?

Responses 'Yes/No' created using numeric response

Survey Question 39: During the last 3 months of your pregnancy, on average, how often did you use e-cigarettes or other electronic nicotine products?

Responses 'Yes/No' created using numeric response

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
E-cigarette use 3 months before pregnancy	3.1	2.2	4.5	1,213	4,381
E-cigarette use last 3 months of pregnancy	1.3	0.7	2.3	1,212	1,800

¹Percentage = Weighted percentage²CI = Confidence interval lower/upper bounds


-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Alcohol Consumption

Percentage (%) of recent mothers in Illinois who reported drinking alcohol in the past two years

Survey Question 40: Have you had any alcoholic drinks in the past 2 years? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	72.4	69.5	75.2	1,213	101,610
Maternal Age	<20 yrs	36.8	21.7	55.1	44	1,833
	20 to 24 yrs	64.3	56.5	71.4	206	15,844
	25 to 29 yrs	73.7	68.0	78.7	313	27,349
	30 to 34 yrs	82.2	77.6	86.0	395	36,030
	=>35 yrs	69.2	62.7	75.1	255	20,555
Race/Ethnicity	NH White	85.1	81.8	87.9	635	64,250
	NH Black	61.1	53.2	68.4	227	13,963
	Hispanic	53.7	46.6	60.7	236	15,612
	Other	57.6	47.0	67.6	102	6,400
Education	<12 yrs	42.7	33.5	52.4	140	7,288
	12 yrs	59.7	52.4	66.6	234	17,248
	>12 yrs	82.1	79.1	84.8	810	74,830
Birth Weight	Low	63.3	58.0	68.3	340	6,584
	Normal	73.2	70.1	76.1	873	95,026
Marital Status	Married	75.8	72.4	78.9	754	63,280
	Other	67.5	62.3	72.3	459	38,329
Insurer before Pregnant	Medicaid	59.9	53.6	65.8	334	23,036
	Other	77.2	74.0	80.1	879	78,574

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition


Alcohol Consumption

Percentage (%) of recent mothers in Illinois who reported drinking alcohol in the three months before pregnancy

Among women who reported drinking alcohol in the past two years

Survey Question 41: During the 3 months before you got pregnant, how many alcoholic drinks did you have in an average week?

Responses 'Yes/No' created using numeric response



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	62.4	59.3	65.4	1,213	87,557
	Maternal Age					
	<20 yrs	24.2	12.1	42.4	44	1,202
	20 to 24 yrs	48.8	41.0	56.6	205	11,938
	25 to 29 yrs	63.3	57.2	69.0	313	23,522
	30 to 34 yrs	72.5	67.4	77.1	395	31,799
	=>35 yrs	64.0	57.4	70.2	256	19,096
Race/Ethnicity	NH White	76.5	72.7	79.9	637	57,915
	NH Black	48.3	40.5	56.2	226	10,953
	Hispanic	41.4	34.6	48.5	235	12,008
	Other	47.7	37.3	58.3	102	5,295
Education	<12 yrs	28.6	20.6	38.1	141	4,933
	12 yrs	47.3	40.1	54.7	233	13,609
	>12 yrs	74.0	70.6	77.1	810	67,363
Birth Weight	Low	51.1	45.8	56.4	340	5,322
	Normal	63.3	60.0	66.5	873	82,235
Marital Status	Married	68.3	64.6	71.7	754	56,998
	Other	53.8	48.5	59.0	459	30,559
Insurer before Pregnant	Medicaid	45.7	39.6	51.9	333	17,506
	Other	68.7	65.2	71.9	880	70,051

¹Percentage = Weighted percentage

²CI = confidence interval

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Alcohol Consumption

Number of drinks per week during the three months before pregnancy

Among women who reported drinking alcohol in the past two years

Survey Question 41: During the 3 months before you got pregnant, how many alcoholic drinks did you have in an average week?



Number of drinks/week	Percentage ¹		95% CI ²		Respondents ³	Estimated Population Affected ⁴
	Weighted	Unweighted	Lower	Upper		
<1	41.7	37.8	35.6	45.6	741	36,469
1-3	39.3	35.5	32.3	43.2	741	34,374
4 or more	19.1	16.2	12.3	22.3	741	16,714

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Stressors

Percentage (%) of recent mothers in Illinois who reported that the following events happened during the 12 months before delivery

Survey Question 42: This question is about things that may have happened during the 12 months before you new baby was born. For each item, check No if it did not happen to you or Yes if it did. Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
I moved to a new address	31.7	28.9	34.7	1,218	44,733
A close family member was very sick and had to go into the hospital	25.0	22.4	27.8	1,215	35,138
My husband, partner, or I had a cut in work hours or pay	17.5	15.2	20.0	1,219	24,707
Someone very close to me died	16.9	14.7	19.4	1,218	23,842
I had problems paying the rent, mortgage, or other bills	16.9	14.7	19.4	1,218	23,840
I argued with my husband or partner more than usual	16.1	13.9	18.6	1,218	22,690
Someone very close to me had a problem with drinking or drugs	10.7	8.9	12.8	1,219	15,094
My husband or partner lost their job	9.8	8.1	11.9	1,218	13,864
I got separated or divorced from my husband or partner	9.0	7.3	11.1	1,218	12,680
I lost my job even though I wanted to go on working	8.8	7.2	10.8	1,218	12,459
My husband or partner said they didn't want me to be pregnant	5.7	4.4	7.4	1,219	8,086
I was apart from my husband or partner due to military deployment or extended work-related travel	3.5	2.5	4.9	1,218	4,973
My husband, partner, or I went to jail	3.1	2.2	4.5	1,219	4,432
I was homeless or had to sleep outside, in a car, or in a shelter	2.5	1.6	3.7	1,219	3,477

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Stressors

Percentage (%) of recent mothers in Illinois who reported the number of stressors indicated by category

Survey Question 42: This question is about things that may have happened during the 12 months before your new baby was born. For each item, check No if it did not happen to you or Yes if it did. Response: Yes count total

	No Stressors			1-2 Stressors			3-5 Stressors			More Than 5 Stressors		
	Percentage ¹	95% Confidence Interval		Percentage ¹	95% Confidence Interval		Percentage ¹	95% Confidence Interval		Percentage ¹	95% Confidence Interval	
Total	31.3	28.5	34.2	44.6	41.5	47.7	17.9	15.6	20.4	6.3	4.9	8.1
Age												
<20 yrs	5.6	1.3	20.8	38.2	23.0	56.2	42.8	26.8	60.4	13.5	5.0	31.4
20 to 24 yrs	24.3	18.2	31.5	39.3	32.0	47.1	26.4	20.1	33.9	10.0	6.2	15.7
25 to 29 yrs	28.5	23.4	34.3	43.5	37.5	49.6	19.2	14.8	24.5	8.8	5.8	13.0
30 to 34 yrs	37.7	32.7	43.1	46.3	41.0	51.8	12.2	9.0	16.3	3.7	2.1	6.6
=>35 yrs	35.2	29.1	41.7	48.6	42.1	55.3	13.5	9.7	18.5	2.7	1.2	6.2
Race/Ethnicity												
NH White	30.2	26.5	34.2	47.7	43.6	51.9	14.5	11.7	17.8	7.5	5.5	10.2
NH Black	25.5	19.2	32.9	38.6	31.3	46.6	26.7	20.4	34.1	9.2	5.5	15.0
Hispanic	33.3	27.0	40.3	42.1	35.3	49.2	22.0	16.7	28.5	2.6	1.1	5.7
Other	40.2	30.4	50.7	42.4	32.3	53.0	14.6	8.5	24.0	2.9	0.7	10.7
Education												
<12 yrs	36.5	27.9	46.1	27.6	19.9	36.8	24.6	17.3	33.6	11.3	6.4	19.2
12 yrs	29.7	23.5	36.7	41.7	34.7	49.1	22.3	16.8	29.0	6.3	3.5	10.9
>12 yrs	30.4	27.1	33.9	48.3	44.6	52.0	15.8	13.2	18.8	5.5	4.0	7.6
Birth Weight												
LBW	32.3	27.6	37.4	33.4	28.6	38.5	26.2	21.8	31.1	8.2	5.7	11.6
NBW	31.2	28.2	34.3	45.5	42.2	48.8	17.2	14.8	20.0	6.1	4.7	8.0
Marital Status												
Married	37.1	33.5	40.8	49.4	45.6	53.2	11.1	9.0	13.7	2.4	1.5	3.9
Other/Unk.	22.7	18.5	27.5	37.5	32.5	42.8	27.8	23.4	32.7	12.0	9.0	15.8
Insurer before Pregnant												
Medicaid	24.9	19.8	30.7	35.8	30.0	42.0	27.3	22.1	33.2	12.0	8.5	16.8
Other	33.6	30.4	37.1	47.8	44.3	51.4	14.4	12.0	17.1	4.1	2.9	5.9


¹Percentage = Weighted percentage

-- Indicates denominator <30 respondents

Stressors

Percentage (%) of recent mothers in Illinois reporting how safe they felt in their neighborhood

Survey Question 43: During the past 12 months before your new baby was born, how often did you feel unsafe in the neighborhood where you lived? Response: Responses Always, Often, or Sometimes combined and shown in chart



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	10.5	8.8	12.6	1,219	14,867
	Maternal Age					
Maternal Age	<20 yrs	9.4	2.9	26.4	44	465
	20 to 24 yrs	10.8	6.8	16.8	205	2,640
	25 to 29 yrs	11.4	8.0	15.8	315	4,253
	30 to 34 yrs	9.6	6.8	13.4	396	4,232
	=>35 yrs	10.9	7.3	15.9	259	3,278
	Race/Ethnicity					
Race/Ethnicity	NH White	6.5	4.7	8.9	637	4,915
	NH Black	20.1	14.6	27.1	226	4,556
	Hispanic	13.2	9.0	18.8	240	3,900
	Other	10.4	5.4	19.0	103	1,165
Education						
Education	<12 yrs	12.2	7.2	19.9	143	2,120
	12 yrs	12.2	8.1	17.8	233	3,489
	>12 yrs	9.7	7.7	12.2	814	8,927
Birth Weight						
Birth Weight	Low	14.3	10.9	18.5	341	1,492
	Normal	10.2	8.4	12.5	878	13,374
Marital Status						
Marital Status	Married	7.4	5.7	9.7	759	6,251
	Other	15.1	11.8	19.3	460	8,615
Insurer before Pregnant						
Insurer before Pregnant	Medicaid	16.6	12.4	21.8	332	6,321
	Other	8.3	6.5	10.5	887	8,545

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Stressors

Percentage (%) of recent mothers in Illinois who reported being treated unfairly during a health care visit

Survey Question 76: During pregnancy, you probably had to get different kinds of health-related services. These may have included clinic visits, doctor's or nurse's office visits, applying for health insurance, applying for Medicaid, or getting help for a family problem. Did you ever feel you were treated unfairly in getting these kinds of services because of any of the following? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
My insurance or Medicaid status	3.7	2.7	5.1	1,206	5,197
I felt unfairly treated for other reasons	2.4	1.6	3.7	1,172	3,317
My age	2.0	1.3	3.2	1,206	2,857
My race, ethnicity, or culture	1.2	0.7	2.1	1,208	1,681
The language I speak	1.0	0.5	1.9	1,206	1,389
My citizenship	0.7	0.3	1.5	1,204	953

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

⁵Any Abuse = Combines 'Yes' responses reported for each person category

-- Indicates denominator <30 respondents

Physical Abuse

Percentage (%) of recent mothers in Illinois who reported being physically abused before pregnancy

Survey Question 44: In the 12 months before you got pregnant with your new baby, did any of the following people push, hit, slap, kick, choke, or physically hurt you in any other way? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Any abuse ⁵	4.0	2.9	5.5	1,224	5,655
My husband or partner	2.3	1.5	3.5	1,206	3,271
Someone else	1.5	0.9	2.6	1,201	2,060
Another family member	0.6	0.2	1.4	1,202	795
My ex-husband or ex-partner	0.5	0.2	1.3	1,203	752

Percentage (%) of recent mothers in Illinois who reported being physically abused during pregnancy

Survey Question 45: During your most recent pregnancy, did any of the following people push, hit, slap, kick, choke, or physically hurt you in any other way? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Any abuse ⁵	2.5	1.6	3.7	1,224	3,506
My husband or partner	0.8	0.4	1.6	1,207	1,126
My ex-husband or ex-partner	0.6	0.3	1.5	1,203	882
Another family member	0.3	0.1	0.9	1,202	401
Someone else	1.1	0.6	2.0	1,202	1,497

Percentage (%) of recent mothers in Illinois who reported being threatened by husband or partner during the time periods indicated

Survey Question 75: During any of the following time periods, did your husband or partner threaten you, limit your activities against your will, or make you feel unsafe in any other way? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
During the 12 months before I got pregnant	1.8	1.1	2.9	1,196	2,512
During my most recent pregnancy	1.4	0.9	2.4	1,194	1,991
Since my new baby was born	2.4	1.6	3.6	1,195	3,285

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

⁵Any Abuse = Combines 'Yes' responses reported for each person category

-- Indicates denominator <30 respondents


Breastfeeding

Percentage (%) of recent mothers in Illinois who reported receiving information about breastfeeding

Among women who reported their baby is living and with them now

Survey Question 50: Before or after your new baby was born, did you receive information about breastfeeding from any of the following sources?

Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Nurse, midwife, doula	81.6	79.0	83.9	1,168	110,854
My doctor	81.2	78.6	83.5	1,171	110,808
Breastfeeding or lactation specialist	76.0	73.1	78.6	1,168	103,496
My baby's doctor or health care provider	70.3	67.3	73.1	1,151	94,318
Family or friends	63.2	60.0	66.2	1,152	84,666
Breastfeeding support group	22.8	20.2	25.7	1,133	30,114
Breastfeeding hotline/toll-free number	5.4	4.1	7.0	1,135	7,081

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Breastfeeding

Percentage (%) of recent mothers in Illinois who reported initiation of breastfeeding

Among women who reported their baby is living and with them now

Survey Question 51: Did you ever breastfeed or pump breast milk to feed your new baby, even for a short period of time? Response: Yes



	Percentage ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴		
All	Overall	88.2	86.0	90.1	1,188	121,862
Maternal Age	<20 yrs	57.5	38.6	74.5	44	2,504
	20 to 24 yrs	87.2	80.9	91.6	198	20,761
	25 to 29 yrs	91.4	87.1	94.3	307	33,556
	30 to 34 yrs	90.7	87.0	93.5	387	39,207
	=>35 yrs	86.0	80.6	90.1	256	25,833
Race/Ethnicity	NH White	88.1	85.0	90.6	626	65,897
	NH Black	81.9	74.9	87.3	212	17,445
	Hispanic	90.3	85.1	93.9	235	26,240
	Other	95.6	87.7	98.5	102	10,721
Education	<12 yrs	78.4	68.8	85.6	130	12,539
	12 yrs	85.7	79.9	90.0	225	24,065
	>12 yrs	90.7	88.3	92.7	806	82,574
Birth Weight	Low	85.0	80.7	88.5	324	8,454
	Normal	88.5	86.1	90.5	864	113,407
Marital Status	Married	92.8	90.5	94.5	750	77,357
	Other	81.3	76.7	85.1	438	44,504
Insurer Now	Medicaid	81.5	76.8	85.4	430	40,921
	Other	92.1	89.7	93.9	758	80,941

¹Percentage = Weighted percentage²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Breastfeeding

Percentage (%) of recent mothers in Illinois who reported the following reasons for not breastfeeding

Among women who did not initiate breastfeeding, even for a short period of time

Survey Question 52: What were your reasons for not breastfeeding your new baby? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
I didn't like breastfeeding	35.6	27.1	45.2	140	5,836
I didn't want to	29.6	21.8	38.7	140	4,843
I had other children to take care of	20.1	13.6	28.6	140	3,287
I went back to work	19.8	13.3	28.3	140	3,241
I was sick or on medicine	12.6	7.6	20.2	140	2,067
I had too many household duties	9.4	5.1	16.9	140	1,543
I tried but it was too hard	5.6	2.5	12.2	140	922
I went back to school	4.4	1.7	10.7	140	714

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds -- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Breastfeeding

Percentage (%) of recent mothers in Illinois who reported continuing breastfeeding at one month and two months after delivery¹

Among women who reported initiating breastfeeding



Breastfeeding

Percentage (%) of recent mothers in Illinois who reported continuing breastfeeding at one month and two months after delivery¹

Among women who reported initiating breastfeeding

	Still breastfeeding at 1 month					Still breastfeeding at 2 months				
	Percentage ²	95% Confidence Interval		Respondents ³	Estimated Population Affected ⁴	Percentage ²	95% Confidence Interval		Respondents ³	Estimated Population Affected ⁴
All	77.0	74.2	79.6	1,173	104,683	68.3	65.2	71.2	1,173	92,866
Age										
< 20 years	43.9	27.0	62.4	40	1,912	34.6	19.4	53.7	40	1,505
20-24 years	70.5	62.6	77.2	193	16,222	56.0	47.9	63.7	193	12,882
25-29 years	79.9	74.3	84.5	304	29,082	71.6	65.6	77.0	304	26,080
30-34 years	83.4	78.9	87.2	384	35,664	74.3	69.2	78.9	384	31,769
=> 35 years	74.1	67.6	79.6	252	21,804	70.1	63.5	75.9	252	20,630
Race/Ethnicity										
NH White	77.1	73.3	80.5	620	57,126	68.7	64.6	72.5	620	50,890
NH Black	65.6	57.3	73.0	208	13,561	55.8	47.5	63.8	208	11,529
Hispanic	78.6	72.0	84.0	230	22,181	67.7	60.6	74.1	230	19,117
Other	93.2	85.0	97.1	102	10,454	88.9	80.0	94.1	102	9,969
Education										
< High School	66.8	56.8	75.5	130	10,685	55.7	45.7	65.2	130	8,905
High School	67.1	59.6	73.8	219	18,095	55.2	47.6	62.5	219	14,874
> High School	81.8	78.6	84.5	797	73,550	74.2	70.7	77.4	797	66,734
Birth Weight										
Low	74.8	69.8	79.3	322	7,393	63.5	58.1	68.6	322	6,271
Normal	77.2	74.1	80.0	851	97,290	68.7	65.4	71.8	851	86,595
Insurer Now										
Medicaid	66.7	61.3	71.7	423	32,697	55.7	50.2	61.1	423	27,312
Other	82.8	79.6	85.6	750	71,987	75.4	71.8	78.6	750	65,555

¹Question 53: Are you currently breastfeeding or feeding pumped milk to your new baby?


-- Indicates denominator <30 respondents

¹Question 54: How many weeks or months did you breastfeed or feed pumped milk to your baby?²Percentage = Weighted percentage³Respondents = The total number who responded to the survey question⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Breastfeeding

Percentage (%) of recent mothers in Illinois who reported continuing breastfeeding exclusively at 12 weeks after delivery¹

Among women who reported initiating breastfeeding



Breastfeeding

Percentage (%) of recent mothers in Illinois who reported continuing breastfeeding exclusively at 12 weeks after delivery¹

Among women who reported initiating breastfeeding

	Still breastfeeding at 12 weeks				Exclusive Breastfeeding at 12 Weeks					
	Percentage ²	95% Confidence Interval		Respondents ³	Estimated Population Affected ⁴	Percentage ²	95% Confidence Interval		Respondents ³	Estimated Population Affected ⁴
All	60.7	57.5	63.9	1,141	80,147	34.2	31.2	37.2	1,154	45,679
Age										
< 20 years	20.8	9.6	39.6	40	907	15.4	6.2	33.4	40	670
20-24 years	43.6	35.8	51.7	191	9,887	19.5	14.0	26.6	194	4,508
25-29 years	65.2	58.9	71.0	290	22,609	36.1	30.3	42.2	296	12,763
30-34 years	68.5	63.1	73.4	376	28,648	41.3	36.0	46.8	379	17,501
=> 35 years	63.8	56.9	70.1	244	18,097	35.9	29.7	42.6	245	10,238
Race/Ethnicity										
NH White	61.9	57.6	66.0	597	44,073	43.0	38.8	47.3	604	31,038
NH Black	45.3	37.2	53.6	204	9,152	18.9	13.5	25.9	209	3,953
Hispanic	59.9	52.6	66.8	228	16,671	25.6	19.8	32.3	229	7,176
Other	82.6	72.6	89.5	99	9,023	27.3	18.9	37.8	99	2,985
Education										
< High School	47.9	38.2	57.7	130	7,661	19.2	12.6	28.1	129	3,031
High School	46.0	38.5	53.7	215	12,179	20.2	14.8	26.8	217	5,429
> High School	67.4	63.6	70.9	771	58,449	41.3	37.6	45.1	782	36,361
Birth Weight										
Low	52.6	47.1	58.1	317	5,119	23.6	19.3	28.5	319	2,308
Normal	61.4	58.0	64.7	824	75,028	35.0	31.8	38.3	835	43,371
Insurer Now										
Medicaid	46.2	40.7	51.7	418	22,283	20.8	16.7	25.6	423	10,253
Other	69.2	65.4	72.7	723	57,864	41.9	38.1	45.8	731	35,427

¹Question 53: Are you currently breastfeeding or feeding pumped milk to your new baby?


-- Indicates denominator <30 respondents

¹Question 54: How many weeks or months did you breastfeed or feed pumped milk to your baby?¹Question 55: How old was your new baby the first time he or she had liquids other than breast milk (such as formula, water, juice, or cow's milk)?¹Question 56: How old was your new baby the first time he or she ate food (such as baby cereal, baby food, or any other food)?²Percentage = Weighted percentage³Respondents = The total number who responded to the survey question⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Infant Sleep Habits

Percentage (%) of recent mothers in Illinois who reported most often placing infants on their back to sleep

Survey Question 57: In which one position do you most often lay your baby down to sleep now? Response: On his or her back



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	81.5	78.9	83.8	1,175	111,812
Maternal Age	<20 yrs	72.6	53.6	85.8	39	3,130
	20 to 24 yrs	72.1	64.5	78.7	195	16,943
	25 to 29 yrs	83.0	78.0	87.1	304	30,190
	30 to 34 yrs	87.6	83.4	90.9	384	37,788
	=>35 yrs	79.4	73.4	84.3	253	23,761
Race/Ethnicity	NH White	87.9	84.8	90.4	622	65,401
	NH Black	62.0	53.8	69.6	205	13,009
	Hispanic	81.7	75.5	86.6	234	23,599
	Other	76.7	66.7	84.5	101	8,584
Education	<12 yrs	73.9	64.5	81.6	130	11,904
	12 yrs	79.4	72.9	84.7	223	22,111
	>12 yrs	83.8	80.9	86.4	796	75,688
Birth Weight	Low	75.4	70.3	79.8	315	7,286
	Normal	81.9	79.2	84.4	860	104,526
Marital Status	Married	84.1	81.1	86.7	745	69,829
	Other	77.4	72.6	81.6	430	41,983
Insurer Now	Medicaid	72.6	67.6	77.2	423	36,093
	Other	86.5	83.6	88.9	752	75,719

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Infant Sleep Habits

Percentage (%) of recent mothers in Illinois who reported their baby slept in their own crib

Survey Question 58: In the past 2 weeks, how often has your new baby slept alone in his or her own crib or bed? Response: Always/Often/Sometimes combined



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	86.1	83.7	88.2	1,175	118,213
Maternal Age	<20 yrs	81.7	63.2	92.1	38	3,492
	20 to 24 yrs	77.0	69.5	83.1	195	18,094
	25 to 29 yrs	83.9	78.8	88.0	305	30,620
	30 to 34 yrs	89.8	86.0	92.6	384	38,726
	=>35 yrs	91.1	86.4	94.3	253	27,281
Race/Ethnicity	NH White	88.9	85.9	91.4	622	66,181
	NH Black	72.9	64.9	79.7	205	15,358
	Hispanic	91.7	87.0	94.7	235	26,599
	Other	78.3	68.2	85.8	100	8,649
Education	<12 yrs	84.5	75.7	90.5	129	13,487
	12 yrs	83.9	77.6	88.6	222	23,317
	>12 yrs	87.0	84.2	89.3	798	78,764
Birth Weight	Low	87.2	83.0	90.4	314	8,391
	Normal	86.0	83.5	88.2	861	109,821
Marital Status	Married	88.8	86.2	91.0	746	73,862
	Other	81.9	77.3	85.7	429	44,350
Insurer Now	Medicaid	80.8	76.1	84.8	421	40,024
	Other	89.0	86.4	91.2	754	78,188

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Infant Sleep Habits

Percentage (%) of recent mothers in Illinois whose baby slept in their own crib in the same room with the mother

Among women who reported their baby sleeps alone in a crib or bed

Survey Question 59: When your new baby sleeps alone, is his or her crib or bed in the same room where you sleep? Response: Always/Often/Sometimes combined



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	73.8	70.8	76.6	1,071	92,353
Maternal Age	<20 yrs	88.8	67.5	96.8	34	3,429
	20 to 24 yrs	83.8	76.2	89.3	170	16,848
	25 to 29 yrs	72.3	66.1	77.8	268	23,221
	30 to 34 yrs	70.9	65.5	75.7	356	28,404
	=>35 yrs	70.6	64.1	76.4	243	20,451
Race/Ethnicity	NH White	64.2	59.9	68.3	583	44,479
	NH Black	84.9	77.2	90.3	171	14,803
	Hispanic	90.9	85.6	94.3	219	24,893
	Other	77.9	66.6	86.2	86	7,311
Education	<12 yrs	90.0	81.2	94.9	116	12,698
	12 yrs	81.9	74.9	87.3	199	20,528
	>12 yrs	68.7	65.0	72.2	732	57,111
Birth Weight	Low	80.0	75.0	84.2	284	6,960
	Normal	73.4	70.2	76.3	787	85,393
Marital Status	Married	69.1	65.3	72.6	692	53,299
	Other	81.4	76.4	85.6	379	39,054
Insurer Now	Medicaid	85.6	80.9	89.4	370	37,167
	Other	67.5	63.7	71.2	474	55,186

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Infant Sleep Habits

Percentage (%) of recent mothers in Illinois who reported the following related to their baby's sleep environment

Survey Question 60: Listed below are some more things about how babies sleep. How did your new baby usually sleep in the past 2 weeks? For each item, check No if your baby did not usually sleep like this or Yes if he or she did. Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
In a crib, bassinet, or pack and play	88.9	86.7	90.8	1,169	121,523
With a blanket	44.0	40.9	47.2	1,156	59,525
In an infant car seat or swing	39.5	36.5	42.7	1,162	53,760
In a sleeping sack or wearable blanket	37.3	34.3	40.4	1,157	50,376
On a twin or larger mattress or bed	32.0	29.1	35.1	1,157	43,348
With crib bumper pads (mesh or non-mesh)	15.6	13.4	18.1	1,156	21,096
With toys, cushions, or pillows, including nursing pillows	9.1	7.4	11.2	1,158	12,359
On a couch, sofa, or armchair	7.6	6.1	9.4	1,158	10,283

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds -- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Infant Sleep Habits

Percentage (%) of recent mothers in Illinois who reported receiving information about baby sleep positions from a health care worker

Survey Question 61: Did a doctor, nurse, or other health care worker tell you any of the following things? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Place my baby on his or her back to sleep	93.0	91.2	94.5	1,172	127,582
What things should and should not go in bed with my baby	87.7	85.5	89.7	1,169	119,871
Place my baby to sleep in a crib, bassinet, or pack and play	87.6	85.4	89.6	1,170	119,834
Place my baby's crib or bed in my room	56.4	53.2	59.5	1,169	77,105

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported using birth control now

Survey Question 62: Are you or your husband or partner doing anything now to keep from getting pregnant? Response: Yes



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	80.9	78.3	83.2	1,212	113,394
Maternal Age	<20 yrs	88.8	73.4	95.8	42	4,205
	20 to 24 yrs	78.8	71.7	84.6	201	18,966
	25 to 29 yrs	80.5	75.3	84.9	315	30,123
	30 to 34 yrs	81.5	76.9	85.4	395	35,735
	=>35 yrs	80.9	75.2	85.5	259	24,365
Race/Ethnicity	NH White	84.0	80.7	86.8	637	63,721
	NH Black	78.6	71.5	84.3	221	17,313
	Hispanic	77.6	71.1	83.1	238	22,753
	Other	72.5	62.3	80.8	103	8,150
Education	<12 yrs	73.1	63.8	80.7	141	12,502
	12 yrs	82.6	76.3	87.5	230	23,473
	>12 yrs	81.8	78.8	84.5	813	74,844
Birth Weight	Low	75.7	70.9	80.0	338	7,827
	Normal	81.3	78.6	83.8	874	105,567
Marital Status	Married	80.8	77.7	83.6	759	67,923
	Other	81.0	76.5	84.8	453	45,471
Insurer Now	Medicaid	80.6	76.0	84.5	447	41,647
	Other	81.1	77.9	83.9	765	71,746

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported using the following birth control methods

Among women reporting using birth control now

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response:

Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Condoms	34.8	31.6	38.2	965	39,456
Birth control pills	22.8	20.0	25.9	965	25,868
IUD (including Mirena®, ParaGard®, Liletta®, or Skyla®)	14.1	11.9	16.8	965	16,042
Withdrawal (pulling out)	14.1	11.9	16.7	965	16,037
Tubes tied or blocked (female sterilization or Essure®)	7.1	5.5	9.1	965	8,060
Shots or injections (Depo-Provera®)	6.3	4.8	8.3	965	7,151
Not having sex (abstinence)	9.8	7.9	12.1	965	11,103
Natural family planning (including rhythm method)	7.2	5.6	9.1	965	8,124
Contraceptive implant in the arm (Nexplanon® or Implanon®)	4.4	3.1	6.1	965	4,988
Vasectomy (male sterilization)	3.0	2.1	4.4	965	3,431
Contraceptive patch (OrthoEvra®) or vaginal ring (NuvaRing®)	2.2	1.4	3.5	965	2,504

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported using a most or moderately effective* form of birth control

Among women who were not trying to get pregnant and who were not currently pregnant

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response: Yes to a most or moderately effective method



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	50.9	47.7	54.1	1,159	68,448
Maternal Age	<20 yrs	66.2	48.1	80.6	41	3,008
	20 to 24 yrs	55.3	47.2	63.0	194	12,927
	25 to 29 yrs	46.5	40.3	52.7	304	16,837
	30 to 34 yrs	48.7	43.2	54.2	377	20,427
	=>35 yrs	53.9	47.0	60.6	243	15,248
Race/Ethnicity	NH White	51.2	46.9	55.5	600	36,706
	NH Black	47.1	39.2	55.2	215	10,187
	Hispanic	60.3	53.2	67.1	232	17,323
	Other	30.3	21.4	41.1	99	3238.6
Education	<12 yrs	57.1	47.3	66.4	136	9,471
	12 yrs	56.8	49.4	64.0	226	15,907
	>12 yrs	47.7	43.9	51.5	771	41,455
Birth Weight	Low	52.1	46.7	57.5	324	5,176
	Normal	50.8	47.4	54.2	835	63,271
Marital Status	Married	46.5	42.7	50.5	713	36,746
	Other	57.1	51.8	62.3	446	31,701
Insurer Now	Medicaid	55.5	50.0	60.8	436	28,032
	Other	48.2	44.2	52.1	723	40,415

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

*Most Moderately Effective Method includes male/female sterilization, birth control pills, implant (e.g., Nexplanon®, Implanon®), IUD (e.g., Mirena®, ParaGard®, Liletta®, Skyla®), contraceptive patch (e.g. OrthoEvra®) or shot (Depo-Provera®), or vaginal ring (e.g., NuvaRing®).

Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported the following reasons for not using birth control after delivery

Among women reporting they are not using birth control now

Survey Question 63: What are your reasons or your husband's or partner's reasons for not doing anything to keep from getting pregnant now? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
I don't want to use birth control	29.4	23.4	36.2	247	7,894
I am not having sex	28.4	22.3	35.3	247	7,622
I am worried about side effects from birth control	23.1	17.7	29.6	247	6,210
I want to get pregnant	15.8	11.4	21.4	247	4,237
My husband or partner doesn't want to use anything	7.1	4.2	11.8	247	1,905
I had my tubes tied or blocked	6.8	4.0	11.4	247	1,835
I have problems paying for birth control	2.2	0.9	5.3	247	591
I am pregnant now	1.5	0.4	4.8	247	394

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Postpartum Care

Percentage (%) of recent mothers in Illinois who received a postpartum visit

Survey Question 65: Since your new baby was born, have you had a postpartum checkup for yourself? Response: Yes



	Percentage ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴		
All	Overall	92.4	90.5	94.0	1,209	129,243
Maternal Age	<20 yrs	71.6	52.6	85.1	42	3,391
	20 to 24 yrs	89.3	83.0	93.4	198	21,213
	25 to 29 yrs	91.7	87.6	94.6	314	34,192
	30 to 34 yrs	94.4	91.0	96.5	396	41,507
	=>35 yrs	96.1	92.8	97.9	259	28,940
Race/Ethnicity	NH White	94.2	91.8	96.0	637	71,443
	NH Black	87.2	80.7	91.8	221	19,219
	Hispanic	90.2	84.9	93.7	238	26,520
	Other	95.4	87.3	98.4	100	10,345
Education	<12 yrs	82.0	73.1	88.3	141	14,022
	12 yrs	87.8	82.0	91.9	229	24,933
	>12 yrs	95.7	93.9	97.0	811	87,322
Birth Weight	Low	90.8	87.1	93.5	337	9,360
	Normal	92.5	90.5	94.2	872	119,883
Marital Status	Married	95.7	93.8	97.0	757	80,132
	Other	87.5	83.5	90.7	452	49,111
Insurer Now	Medicaid	89.4	85.6	92.4	445	46,081
	Other	94.1	91.9	95.8	764	83,162

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Postpartum Care

Percentage (%) of recent mothers in Illinois who reported the following reasons for not having a postpartum visit

Among women reporting they did not have a postpartum visit

Survey Question 66: Did any of these things keep you from having a postpartum checkup? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
I had too many things going on	39.1	28.5	50.8	91	4,246
I felt fine and did not think I needed to have a visit	26.5	17.6	37.9	91	2,880
I didn't have any transportation to get to the clinic or doctor's office	15.1	8.4	25.7	91	1,643
I couldn't get an appointment when I wanted one	14.7	8.4	24.5	91	1,597
I didn't have health insurance to cover the cost of the visit	10.6	5.5	19.6	91	1,153
I couldn't take time off from work	6.3	2.4	15.8	91	689

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Postpartum Care

Percentage (%) of recent mothers in Illinois who reported the following location for their postpartum visit

Among women reporting they had a postpartum visit

Survey Question 67: Where did you go for your postpartum checkup? Response: Yes - Categories summed total 100%

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
My OB/GYN's office	67.2	64.0	70.2	1,119	86,796
Hospital clinic	10.0	8.2	12.3	1,119	12,977
Community health clinic	7.5	5.9	9.5	1,119	9,724
Other*	7.4	5.9	9.2	1,119	9,521
My family doctor's office	5.1	3.8	6.7	1,119	6,529
Health department clinic	2.8	1.9	4.2	1,119	3,626

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

* Most frequent "Other" narrative response was a midwife's office.

Postpartum Care

Percentage (%) of recent mothers in Illinois who reported a health care worker did any of these things during the postpartum visit

Among women reporting they had a postpartum visit

Survey Question 68: During your postpartum checkup, did a doctor, nurse, or other health care worker do any of the following things?

Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Ask me if I was feeling down or depressed	90.8	88.8	92.5	1,121	117,231
Talk to me about birth control methods I can use after giving birth	89.7	87.6	91.6	1,121	115,838
Tell me to take a vitamin with folic acid	64.2	61.0	67.2	1,121	82,943
Talk to me about healthy eating, exercise, and losing weight gained during pregnancy	63.2	60.0	66.3	1,120	81,590
Ask me if someone was hurting me emotionally or physically	56.1	52.9	59.3	1,122	72,583
Ask me if I was smoking cigarettes	56.0	52.7	59.2	1,121	72,262
Talk to me about how long to wait before getting pregnant again	53.0	49.7	56.2	1,118	68,125
Give or prescribe me a contraceptive method such as the pill, patch, shot (Depo-Provera®), NuvaRing®, or condoms	40.0	36.9	43.3	1,118	51,511
Test me for diabetes	25.3	22.6	28.3	1,117	32,561
Insert an IUD (Mirena®, ParaGard®, Liletta®, or Skyla®) or a contraceptive implant (Nexplanon®, or Implanon®)	16.0	13.7	18.5	1,113	20,458

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Mental Health

Percentage (%) of recent mothers in Illinois who reported symptoms of depression after delivery

Survey Question 69: Since your new baby was born, how often have you felt down, depressed or hopeless? Response: Yes - Categories summed total 100%

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Always	0.7	0.30	1.50	1,209	999
Often	4.1	3.00	5.50	1,209	5,716
Sometimes	18.1	15.90	20.70	1,209	25,350
Rarely	29.3	26.60	32.20	1,209	40,967
Never	47.8	44.60	50.90	1,209	66,781

Survey Question 70: Since your new baby was born, how often have you had little interest or little pleasure in doing things you usually enjoyed? Response: Yes - Categories summed total 100%

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Always	2.6	1.7	3.7	1,207	3,563
Often	3.8	2.8	5.1	1,207	5,270
Sometimes	14.8	12.7	17.1	1,207	20,604
Rarely	23.3	20.8	26.0	1,207	32,518
Never	55.6	52.5	58.7	1,207	77,693

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Mental Health

Percentage (%) of recent mothers in Illinois who reported symptoms of depression after delivery

Question 69: Since your new baby was born, how often have you felt down, depressed, or hopeless? A. Always, B. Often, C. Sometimes, D. Rarely, E. Never. Responses A, B, and C were combined to identify those mothers with self-reported depression after pregnancy.



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	22.9	20.4	25.7	1,209	32,065
Maternal Age	<20 yrs	35.1	20.1	53.7	42	1,663
	20 to 24 yrs	24.7	18.5	32.1	200	5,937
	25 to 29 yrs	19.9	15.6	25.1	315	7,451
	30 to 34 yrs	21.7	17.6	26.6	395	9,519
	=>35 yrs	25.1	19.8	31.3	257	7,497
Race/Ethnicity	NH White	24.4	21.0	28.2	637	18,538
	NH Black	24.3	18.1	31.8	220	5,305
	Hispanic	17.4	12.7	23.4	236	5,080
	Other	22.7	15.0	32.9	103	2,552
Education	<12 yrs	24.6	17.3	33.8	141	4,201
	12 yrs	19.0	13.9	25.4	227	5,347
	>12 yrs	23.2	20.2	26.6	813	21,265
Birth Weight	Low	29.7	25.0	34.8	337	3,060
	Normal	22.4	19.7	25.3	872	29,006
Marital Status	Married	21.1	18.1	24.3	758	17,668
	Other	25.8	21.4	30.7	451	14,398
Insurer Now	Medicaid	26.8	22.3	31.8	444	13,741
	Other	20.7	17.8	23.9	765	18,324

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Mental Health

Percentage (%) of recent mothers in Illinois who reported lack of interest in doing things after delivery

Question 70: Since your new baby was born, how often have you had little interest or little pleasure in doing things you usually enjoyed? A. Always, B. Often, C. Sometimes, D. Rarely, E. Never. Responses A, B, and C were combined to identify those mothers with lack of interest after pregnancy.



	Percentage ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴		
All	Overall	21.1	18.6	23.7	1,207	29,436
Maternal Age	<20 yrs	35.4	20.2	54.3	42	1,678
	20 to 24 yrs	21.7	15.9	28.8	198	5,152
	25 to 29 yrs	23.8	19.1	29.3	315	8,905
	30 to 34 yrs	18.4	14.5	23.0	393	8,026
	=>35 yrs	18.8	14.2	24.6	259	5,676
Race/Ethnicity	NH White	18.9	15.8	22.4	637	14,334
	NH Black	25.2	18.9	32.8	218	5,468
	Hispanic	20.2	15.1	26.4	237	5,900
	Other	30.9	22.1	41.5	102	3,435
Education	<12 yrs	28.4	20.6	37.8	140	4,830
	12 yrs	19.7	14.6	26.0	228	5,572
	>12 yrs	19.8	17.0	23.0	811	18,043
Birth Weight	Low	22.0	17.9	26.7	336	2,259
	Normal	21.0	18.4	23.9	871	27,177
Marital Status	Married	20.6	17.7	23.9	757	17,259
	Other	21.8	17.7	26.5	450	12,178
Insurer Now	Medicaid	26.5	22.0	31.5	443	13,577
	Other	17.9	15.2	21.1	764	15,860

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Mental Health

Percentage (%) of recent mothers in Illinois who reported a diagnosis of postpartum depression

Survey Question 71: Since your new baby was born, has a doctor, nurse, or other health care worker told you that you had depression?

Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	9.6	7.9	11.7	1,209	13,460
Maternal Age	<20 yrs	7.8	2.2	24.3	42	370
	20 to 24 yrs	12.3	8.0	18.6	198	2,929
	25 to 29 yrs	12.1	8.6	16.7	314	4,496
	30 to 34 yrs	7.8	5.3	11.3	396	3,425
	=>35 yrs	7.4	4.6	11.8	259	2,239
Race/Ethnicity	NH White	12.1	9.6	15.2	637	9,201
	NH Black	6.8	3.7	12.0	220	1,484
	Hispanic	5.9	3.4	9.9	237	1,716
	Other	7.8	3.6	15.8	102	861
Education	<12 yrs	9.6	5.3	16.7	140	1,617
	12 yrs	9.2	5.8	14.2	229	2,614
	>12 yrs	9.6	7.5	12.1	812	8,733
Birth Weight	Low	8.3	5.8	11.8	337	859
	Normal	9.7	7.9	11.9	872	12,601
Marital Status	Married	8.3	6.4	10.6	758	6,934
	Other	11.7	8.7	15.5	451	6,526
Insurer Now	Medicaid	13.3	10.1	17.4	445	6,867
	Other	7.5	5.7	9.8	764	6,593

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Mental Health

Percentage (%) of recent mothers in Illinois who reported taking medication for depression

Among women who reported a diagnosis of postpartum depression

Survey Question 72: Since your new baby was born, have you taken prescription medicine for your depression?

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
No	38.0	28.7	48.3	110	4,986
Yes	62.0	51.7	71.3	110	8,143

Percentage (%) of recent mothers in Illinois who reported receiving counseling for depression

Among women who reported a diagnosis of postpartum depression

Survey Question 73: Since your new baby was born, have you gotten counseling for your depression?

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
No	63.6	53.3	72.8	111	8,477
Yes	36.4	27.2	46.7	111	4,849

¹Percentage = Weighted percentage²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Support

Percentage (%) of recent mothers in Illinois who reported they have various kinds of support

Survey Question 74: Since you delivered your new baby, would you have the kinds of help listed below if you needed them? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Someone to help me if I were tired and feeling frustrated with my new baby	88.9	86.7	90.8	1,175	122,193
Someone to talk with about my problems	88.3	86.0	90.2	1,176	121,225
Someone to take care of my baby	88.1	85.9	90.1	1,175	121,158
Someone to help me if I were sick and needed to be in bed	86.8	84.5	88.9	1,176	119,426
Someone to loan me \$50	80.3	77.7	82.7	1,177	110,495

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Support

Percentage (%) of recent mothers in Illinois reporting the engagement and support of their husband/partner

Survey Question 77: When you new baby's father is with your baby, how often does he hug, kiss, hold, or play with the baby?

Response: Yes - Categories summed total 100%

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Always	66.5	63.40	69.40	1,178	91,338
Often	21.8	19.30	24.50	1,178	29,895
Sometimes	5.1	3.80	6.70	1,178	7,000
Rarely	1.3	0.80	2.30	1,178	1,813
Never	2.9	2.00	4.30	1,178	4,006
My new baby's father doesn't regularly spend time with the baby	2.5	1.60	3.80	1,178	3,387

Survey Question 78: Since your new baby was born, how often does your husband or partner provide you with encouragement and emotional support? Response: Yes - Categories summed total 100%

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Always	59.3	56.2	62.4	1,204	82,667
Often	19.1	16.8	21.7	1,204	26,623
Sometimes	12.2	10.2	14.4	1,204	16,961
Rarely	3.5	2.5	4.8	1,204	4,815
Never	6.0	4.6	7.8	1,204	8,328

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Support

Percentage (%) of recent mothers in Illinois who reported the baby's father spends time with the baby in play

Survey Question 77: When your new baby's father is with your baby, how often does he hug, kiss, hold, or play with the baby? A. Always, B. Often, C. Sometimes, D. Rarely, E. Never, F. My new baby's father doesn't regularly spend time with my baby. Responses A, B, and C were combined and reported in chart.



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	93.3	91.4	94.8	1,178	128,233
Maternal Age	<20 yrs	76.6	57.8	88.7	41	3,606
	20 to 24 yrs	89.1	82.7	93.3	194	20,898
	25 to 29 yrs	95.7	92.2	97.6	306	35,161
	30 to 34 yrs	95.2	92.0	97.2	385	40,904
	=>35 yrs	93.5	89.2	96.2	252	27,664
Race/Ethnicity	NH White	94.6	92.1	96.3	623	70,559
	NH Black	83.6	76.5	88.8	209	17,651
	Hispanic	94.6	90.0	97.1	233	27,466
	Other	98.8	91.9	99.8	100	10,801
Education	<12 yrs	86.1	77.6	91.7	134	14,348
	12 yrs	90.9	85.3	94.5	221	25,166
	>12 yrs	95.2	93.2	96.6	797	85,712
Birth Weight	Low	90.5	86.7	93.2	318	8,820
	Normal	93.5	91.5	95.1	860	119,412
Marital Status	Married	98.5	97.2	99.1	746	81,552
	Other	85.5	81.2	88.9	432	46,681
Insurer Now	Medicaid	88.5	84.4	91.6	425	43,893
	Other	96.0	94.1	97.4	753	84,339

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Support

Percentage (%) of recent mothers in Illinois who reported their husband or partner supports them emotionally

Survey Question 78: Since your new baby was born, how often does your husband or partner provide you with encouragement and emotional support? A. Always, B. Often, C. Sometimes, D. Rarely, E. Never. Responses A, B, and C were combined and reported in chart.



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	90.6	88.5	92.3	1,204	126,251
Maternal Age	<20 yrs	78.5	61.2	89.4	42	3,720
	20 to 24 yrs	86.4	79.9	91.1	198	20,500
	25 to 29 yrs	91.3	87.1	94.2	312	33,806
	30 to 34 yrs	93.1	89.6	95.5	395	40,835
	=>35 yrs	91.1	86.3	94.3	257	27,390
Race/Ethnicity	NH White	92.5	89.9	94.5	635	69,998
	NH Black	79.9	72.7	85.6	219	17,473
	Hispanic	90.9	85.5	94.4	235	26,374
	Other	97.1	89.2	99.3	102	10,784
Education	<12 yrs	81.8	73.0	88.2	139	13,899
	12 yrs	91.1	85.8	94.5	230	25,893
	>12 yrs	91.9	89.5	93.8	807	83,452
Birth Weight	Low	86.8	82.7	90.1	334	8,870
	Normal	90.9	88.6	92.7	870	117,380
Marital Status	Married	96.3	94.5	97.5	758	80,786
	Other	81.9	77.5	85.7	446	45,465
Insurer Now	Medicaid	84.3	80.0	87.8	445	43,304
	Other	94.2	92.0	95.9	759	82,947

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Before Pregnancy **Table 73** **2017 Phase 8 Illinois PRAMS Data**

Health Insurance Sources

Percentage (%) of recent mothers in Illinois who reported health insurance sources before pregnancy

Survey Question 12: During the month before you got pregnant with your new baby, what kind of health insurance did you have? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Private health insurance from my job or job of husband/partner	48.8	45.7	51.9	1,224	69,142
Medicaid ⁵	27.3	24.5	30.2	1,224	38,686
I did not have health insurance during the month before I got pregnant	13.7	11.7	16.1	1,224	19,446
Private health insurance from my parents	5.4	4.1	7.0	1,224	7,580
Private health insurance from the Illinois Health Insurance Marketplace or GetcoveredIllinois.gov or HealthCare.gov	3.5	2.6	4.9	1,224	5,023
Other health insurance	1.9	1.2	2.9	1,224	2,639
TRICARE or other military health care	1.3	0.8	2.3	1,224	1,891

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

⁵Medicaid combines response categories "Medicaid" and "CHIP or All Kids"

During Pregnancy **Table 74** **2017 Phase 8 Illinois PRAMS Data**

Health Insurance Sources

Percentage (%) of recent mothers in Illinois who reported health insurance sources during pregnancy for prenatal care

Among women who went for prenatal care

Survey Question 13: During your most recent pregnancy, what kind of health insurance did you have for your prenatal care? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Private health insurance from my job or the job of my husband or partner	48.3	45.2	51.5	1,207	67,527
Medicaid ⁵	45.4	42.3	48.6	1,207	63,489
Private health insurance from my parents	4.4	3.2	6.0	1,207	6,175
Private health insurance from the Illinois Health Insurance Marketplace or GetcoveredIllinois.gov or HealthCare.gov	3.3	2.4	4.6	1,207	4,621
Other health insurance	1.8	1.1	2.9	1,207	2,536
I did not have health insurance for my prenatal care	1.3	0.7	2.2	1,207	1,773
TRICARE or other military health care	1.3	0.8	2.3	1,207	1,863

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

⁵Medicaid combines response categories "Medicaid" and "CHIP, All Kids, or Moms & Babies"

After Delivery **Table 75** **2017 Phase 8 Illinois PRAMS Data**
Health Insurance Sources
Percentage (%) of recent mothers in Illinois who reported health insurance sources now

Survey Question 14: What kind of health insurance do you have now? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Private health insurance from my job or job of my husband or partner	47.7	44.6	50.8	1,224	67,567
Medicaid ⁵	37.3	34.3	40.4	1,224	52,900
I do not have health insurance now	9.3	7.6	11.4	1,224	13,225
Private health insurance from my parents	3.1	2.2	4.5	1,224	4,421
Private health insurance from the Illinois Health Insurance Marketplace or GetcoveredIllinois.gov or HealthCare.gov	3.1	2.2	4.3	1,224	4,355
Other health insurance	1.5	0.9	2.5	1,224	2,180
TRICARE or other military health care	1.1	0.6	1.9	1,224	1,529

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition


⁵Medicaid combines response categories "Medicaid" and "CHIP, All Kids, or Moms & Babies"

Health Insurance Sources

Percentage (%) of recent mothers in Illinois who reported Medicaid* coverage before pregnancy

Survey Question 12: During the month before you got pregnant with your new baby, what kind of health insurance did you have?

Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	27.3	24.5	30.2	1,224	38,686
Maternal Age	<20 yrs	59.6	41.9	75.1	44	2,964
	20 to 24 yrs	47.9	40.2	55.6	208	11,880
	25 to 29 yrs	25.3	20.3	31.1	317	9,570
	30 to 34 yrs	18.4	14.5	23.2	396	8,114
	=>35 yrs	20.4	15.5	26.5	259	6,158
Race/Ethnicity	NH White	18.4	15.3	22.1	639	14,018
	NH Black	64.2	56.5	71.3	228	14,700
	Hispanic	25.8	20.0	32.5	241	7,661
	Other	13.1	7.2	22.7	103	1,473
Education	<12 yrs	46.2	37.0	55.7	144	8,074
	12 yrs	50.8	43.6	58.1	236	14,857
	>12 yrs	15.6	13.0	18.5	815	14,298
Birth Weight	Low	39.2	34.1	44.5	342	4,103
	Normal	26.4	23.4	29.5	882	34,583
Marital Status	Married	11.5	9.3	14.1	761	9,679
	Other	50.6	45.3	55.8	463	29,006

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

*Medicaid combines response categories "Medicaid" and "CHIP or All Kids"

Health Insurance Sources

Percentage (%) of recent mothers in Illinois who reported Medicaid* coverage for prenatal care

Survey Question 13: During your most recent pregnancy, what kind of health insurance did you have for your prenatal care? Response: Yes



		Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
All	Overall	45.4	42.3	48.6	1,207	63,489
Maternal Age	<20 yrs	71.5	52.7	84.9	43	3,415
	20 to 24 yrs	73.4	66.2	79.6	206	18,180
	25 to 29 yrs	47.7	41.6	53.8	312	17,732
	30 to 34 yrs	32.6	27.6	38.0	392	14,169
	=>35 yrs	33.9	27.8	40.6	254	9,993
Race/Ethnicity	NH White	27.9	24.2	32.0	633	21,066
	NH Black	76.1	68.9	82.1	227	17,399
	Hispanic	72.3	65.7	78.0	236	20,951
	Other	29.5	20.5	40.5	98	3,176
Education	<12 yrs	81.3	72.9	87.5	142	13,887
	12 yrs	75.3	68.7	81.0	234	21,833
	>12 yrs	28.5	25.2	32.1	803	25,792
Birth Weight	Low	57.0	51.7	62.1	337	5,877
	Normal	44.5	41.2	47.9	870	57,611
Marital Status	Married	24.3	21.1	27.7	750	20,177
	Other	76.6	71.8	80.8	457	43,312

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

*Medicaid combines response categories "Medicaid" and "CHIP, All Kids, or Moms & Babies"

Health Insurance Sources

Percentage (%) of recent mothers in Illinois who reported Medicaid* coverage now

Survey Question 14: What kind of health insurance do you have now? Response: Yes



	Percentage ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴		
All	Overall	37.3	34.3	40.4	1,224	52,900
Maternal Age	<20 yrs	78.9	61.6	89.7	44	3,926
	20 to 24 yrs	67.1	59.6	73.8	208	16,660
	25 to 29 yrs	37.6	31.8	43.7	317	14,203
	30 to 34 yrs	24.9	20.4	30.0	396	10,941
	=>35 yrs	23.8	18.5	30.1	259	7,171
Race/Ethnicity	NH White	25.7	22.0	29.7	639	19,522
	NH Black	74.5	67.2	80.6	228	17,049
	Hispanic	42.4	35.6	49.5	241	12,615
	Other	23.9	15.8	34.3	103	2,682
Education	<12 yrs	56.3	46.8	65.4	144	9,841
	12 yrs	64.0	56.9	70.6	236	18,709
	>12 yrs	24.6	21.5	28.0	815	22,602
Birth Weight	Low	50.9	45.6	56.1	342	5,331
	Normal	36.3	33.0	39.6	882	47,570
Marital Status	Married	16.4	13.8	19.4	761	13,839
	Other	68.1	62.9	72.8	463	39,061

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question


⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

*Medicaid combines response categories "Medicaid" and "CHIP, All Kids, or Moms & Babies"

Drug Use

Percentage (%) of recent mothers in Illinois who reported using marijuana or hash before or during pregnancy

Survey Question D1: At any time during the 3 months before you got pregnant or during your most recent pregnancy, did you use marijuana or hash in any form? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	10.2	7.2	14.2	379	4,499
Maternal Age	<20 yrs	--	--	--	--	--
	20 to 24 yrs	15.2	7.4	28.6	61	1,132
	25 to 29 yrs	13.8	7.8	23.2	103	1,733
	30 to 34 yrs	5.8	2.6	12.3	118	706
	=>35 yrs	5.7	2.4	13.1	87	598
Race/Ethnicity	NH White	8.1	4.9	13.3	203	1,997
	NH Black	24.6	14.2	39.2	65	1,648
	Hispanic	6.1	2.1	16.4	71	524
	Other	9.1	2.3	30.2	34	331
Education	<12 yrs	12.0	5.1	25.6	52	837
	12 yrs	17.2	9.1	30.3	66	1,364
	>12 yrs	8.0	5.0	12.6	251	2,267
Birth Weight	Low	15.3	9.7	23.5	103	484
	Normal	9.8	6.7	14.2	276	4,015
Marital Status	Married	5.6	3.2	9.7	238	1,501
	Other	17.4	11.3	25.7	141	2,998
Insurer Now	Medicaid	21.4	13.8	31.6	108	2,787
	Other	5.5	3.2	9.3	271	1,712

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Data collected using Marijuana and Drug Use Supplement beginning December 2017. Data collection ended March 2019.

Drug Use

Percentage (%) of recent mothers in Illinois who reported a health care worker did any of these things during the prenatal visit

Among women reporting they had a prenatal visit

Survey Question D6: During any of your prenatal care visits, did a doctor, nurse, or other health care worker do any of the following things?

Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
Ask me if I was using marijuana	58.1	52.4	63.5	372	25,023
Advise me not to use marijuana	37.8	32.5	43.5	371	16,364
Advise me not to breastfeed my baby if I was using marijuana	23.8	19.2	29.0	369	10,208
Recommend that I use marijuana for any reason	1.2	0.4	3.3	373	520


Survey Question D7: During any of your prenatal care visits, did a doctor, nurse, or other health care worker refer you to treatment because of drug use (prescribed or non-prescribed drugs)? Response: Yes & I did not use any drugs (or only used over-the-counter pain relievers) during my pregnancy are combined

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
No	99.8	99.2	99.9	267	30,763
Yes	0.2	0.1	0.8	267	63.11

Drug Use

Percentage (%) of recent mothers in Illinois who reported using marijuana or hash since delivery

Survey Question D8: Since your new baby was born, have you used marijuana or hash in any form? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	4.4	2.6	7.5	381	1,965
Maternal Age	<20 yrs	--	--	--	--	--
	20 to 24 yrs	11.3	4.8	24.5	61	840
	25 to 29 yrs	6.0	2.4	14.0	103	752
	30 to 34 yrs	0.7	0.2	2.0	119	81
	=>35 yrs	2.8	0.8	9.4	88	292
Race/Ethnicity	NH White	3.2	1.3	7.3	203	773
	NH Black	11.6	5.0	24.4	65	773
	Hispanic	2.6	0.4	13.4	72	221
	Other	5.3	0.8	28.9	35	197
Education	<12 yrs	4.3	1.1	14.9	53	302
	12 yrs	12.3	5.5	25.5	66	977
	>12 yrs	2.4	1.0	5.6	252	685
Birth Weight	Low	8.0	4.2	14.7	104	256
	Normal	4.2	2.2	7.6	277	1,709
Marital Status	Married	1.6	0.5	4.5	240	425
	Other	8.9	4.8	16.1	141	1,539
Insurer Now	Medicaid	10.7	5.5	19.6	108	1,388
	Other	1.8	0.7	4.7	273	576

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Data collected using Marijuana and Drug Use Supplement beginning December 2017. Data collection ended March 2019.

Drug Use

Percentage (%) of recent mothers in Illinois who reported how long they believe women should wait after using marijuana to breastfeed


Survey Question D9: How long do you think it is necessary for a woman to wait after using marijuana to breastfeed her baby? Response: Yes

	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴
I don't think it is safe for breastfeeding women to use marijuana at all	94.2	90.9	96.3	366	39,626
I think it is best to wait at least 2-3 hours after she is no longer high	3.3	1.8	6.1	366	1,392
I don't think she needs to wait at all	1.6	0.6	3.8	366	654
I think it is best to wait until she is no longer high	0.9	0.3	2.7	366	398

Drug Use

Percentage (%) of recent mothers in Illinois who reported taking prescription antidepressants during pregnancy

Survey Question D10: During your most recent pregnancy, did you take prescription antidepressants or selective serotonin reuptake inhibitors (SSRIs) such as Prozac, Zoloft, or Lexapro? Response: Yes



	Percentage ¹	95% CI ²	Respondents ³	Estimated Population Affected ⁴		
All	Overall	5.2	3.2	8.4	379	2,288
Maternal Age	<20 yrs	--	--	--	--	--
	20 to 24 yrs	6.2	1.8	18.7	60	450
	25 to 29 yrs	5.8	2.4	13.2	102	724
	30 to 34 yrs	4.0	1.5	10.4	119	492
	=>35 yrs	4.0	1.4	11.1	88	425
Race/Ethnicity	NH White	8.6	5.1	14.0	202	2,092
	NH Black	0.0	.	.	65	0
	Hispanic	2.3	0.6	8.9	71	197
	Other	0.0	.	.	35	0
Education	<12 yrs	8.8	3.0	23.0	53	615
	12 yrs	2.5	0.6	9.6	66	197
	>12 yrs	5.2	2.9	9.3	250	1,476
Birth Weight	Low	4.5	1.9	10.3	104	144
	Normal	5.3	3.1	8.7	275	2,145
Marital Status	Married	4.2	2.2	7.9	238	1,125
	Other	6.7	3.2	13.5	141	1,164
Insurer Now	Medicaid	7.1	3.0	15.5	108	918
	Other	4.4	2.4	7.9	271	1,371

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents


³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Drug Use

Percentage (%) of recent mothers in Illinois who reported taking prescription pain relievers during pregnancy

Survey Question D11: During your most recent pregnancy, did you use prescription pain relievers such as hydrocodone (Vicodin®), oxycodone (Percocet®), or codeine? Response: Yes



	Percentage ¹	95% CI ²		Respondents ³	Estimated Population Affected ⁴	
All	Overall	3.9	2.3	6.8	379	1,733
Maternal Age	<20 yrs	--	--	--	--	--
	20 to 24 yrs	7.1	2.5	18.4	60	520
	25 to 29 yrs	2.9	0.8	9.9	103	362
	30 to 34 yrs	6.4	2.9	13.9	119	794
	=>35 yrs	0.5	0.1	2.1	87	56
Race/Ethnicity	NH White	4.7	2.3	9.1	203	1,142
	NH Black	2.0	0.3	12.8	65	134
	Hispanic	2.7	0.5	13.6	70	228
	Other	6.1	1.8	19.1	35	228
Education	<12 yrs	6.1	1.7	19.9	52	418
	12 yrs	5.0	1.5	15.3	66	394
	>12 yrs	3.2	1.6	6.4	251	921
Birth Weight	Low	6.5	3.1	12.9	104	207
	Normal	3.7	2.0	6.9	275	1,526
Marital Status	Married	3.9	2.0	7.5	238	1,054
	Other	3.9	1.5	10.1	141	678
Insurer Now	Medicaid	3.5	1.1	11.2	107	457
	Other	4.1	2.2	7.5	272	1,276

¹Percentage = Weighted percentage

²CI = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents

³Respondents = The total number who responded to the survey question

⁴Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition

Data collected using Marijuana and Drug Use Supplement beginning December 2017. Data collection ended March 2019.