

INFLUENZA SURVEILLANCE UPDATE

ILLINOIS DEPARTMENT OF PUBLIC HEALTH Division of Infectious Disease Week 50: Week Ending Saturday, December 17, 2016


All data in this report are provisional and may change as additional reports are received. For questions, please contact the IDPH CD Section at 217-782-2016 or <u>dph.influenza@illinois.gov</u>. Additional data on influenza in Chicago can be found on the <u>City of Chicago Influenza Website</u>

| Current Week Quick Stats | | | | | |
|---|---|--|--|--|--|
| Illinois Influenza Geographic Spread | Sporadic | | | | |
| Percent of Outpatient Visits for ILI ¹ | 1.33% (baseline 1.9%) | | | | |
| Percent/Number of Influenza Positive Tests ² | Current Week: 2.70% (14/517); Season: 1.08% (50/4617) | | | | |
| Influenza-Associated ICU Admissions ³ | Current Week: 6; Season: 29 | | | | |
| Influenza Outbreaks | Current Week: 2; Season: 6 | | | | |
| Influenza-Associated Pediatric Deaths (Season Total) | 0 | | | | |

Become an Illinois Sentinel Provider

Illinois outpatient health care providers are encouraged to join the Illinois influenza surveillance program. Each week, providers report data to CDC on the number of patients seen and the number with influenza-like illness. These reporters are critical to determining when and where influenza activity is occurring and who it is affecting. For more information on how to participate, contact <u>DPH.INFLUENZA@ILLINOIS.GOV</u>.

Illinois Influenza-Like Illness (ILI) Surveillance




¹ ILI "Influenza like Illness" is defined as fever $\geq 100^{\circ}$ F and cough and/or sore throat.

² Specimens tested by WHO/NREVSS collaborating laboratories and IDPH laboratories.


³ For the purpose of diagnosis, influenza can be diagnosed by using the following test: reverse transcription polymerase chain reaction RT-PCR], viral culture, Immunofluorescence [Direct Fluorescent Antibody (DFA) or Indirect Fluorescent Antibody (IFA) Staining], Enzyme Immuno Assay (EIA) or any rapid diagnostic test. Sensitivities of rapid diagnostic tests are approximately 50-70% when compared with viral culture or reverse transcription polymerase chain reaction (RT-PCR), and specificities of rapid diagnostic tests for influenza are approximately 90-95%. False-positive (and true-negative) results are more likely to occur when disease prevalence in the community is low, which is generally at the beginning and end of the influenza seasons. False-negative (and true-positive) results are more likely to occur when disease prevalence is high in the community, which is typically at the height of the influenza season.

Regional Sentinel Reports




Percent of ILI Reported From Sentinel Providers by Region, Illinois, Influenza Season 2016-2017

Number of Office Visits by Age Group




Week Ending



Illinois Influenza-associated Intensive Care Unit (ICU) Admissions


| Age | Current Week | Season Total (10/02/2016 – Current Week) |
|-------|--------------|--|
| 0-4 | 0 | 0 |
| 5-24 | 0 | 1 |
| 25-49 | 2 | 8 |
| 50-64 | 2 | 7 |
| >64 | 2 | 13 |
| Total | 6 | 29 |



| | illinois Laboratory Surveillance (IDPH, NREV55 & ACL Laboratories) | | | | | | |
|-----------------|--|-----------|------------------------------------|----------------------------|----------------------------|------------------------------------|--|
| | A (H1N1) pdm09 | A (H3) | A (Sub typing not performed) | B (Victoria Lineage) | B (Yamagata Lineage) | B (Lineage not performed) | |
| Current Week | 0 | 8 | 3 | 0 | 0 | 3 | |
| Season Total | 0 | 18 | 14 | 6 | 0 | 18 | |

NDEVCC

Influenza Isolates Reported by WHO/NREVSS Collaborating Laboratories, IDPH Lab & ACL Labs, Ilinois, Influenza Season 2016-2017




Vaccination is the best way to protect against influenza infection. All Illinois residents aged six months and older should be vaccinated annually.

- Flu vaccination can keep you from getting sick with flu.
- Flu vaccination can reduce the risk of flu-associated hospitalization, including among children and older adults.
- Flu vaccination is an important preventive tool for people with chronic health conditions.
- Vaccination helps protect women during and after pregnancy. Getting vaccinated also protects the developing baby during pregnancy and for several months after the baby is born.
- Flu vaccination also may make your illness milder if you do get sick.
- Getting vaccinated yourself also protects people around you, including those who are more vulnerable to serious flu illness, like babies and young children, older people, and people with certain chronic health

| Influenza Outbreaks | | | | | | |
|---------------------|-------|--------------|---|--|--|--|
| Region | | Current Week | Season Total (10/02/2016 – Current Week) | | | |
| Rockford | | 0 | 0 | | | |
| Peoria | | 0 | 1 | | | |
| Edwardsville | | 0 | 1 | | | |
| Marion | | 0 | 1 | | | |
| Champaign | | 0 | 0 | | | |
| West Chicago | | 1 | 1 | | | |
| Chicago/Cook | | 1 | 2 | | | |
| | Total | 2 | 6 | | | |

Influenza Outbreaks by First Onset Date, Illinois, 2016-2017



Week Ending

Resources

- **IDPH Seasonal Influenza** •
- **IDPH Immunization** •
- **CDC** Influenza
- **CDC** Influenza Vaccine
- **Cook County Influenza Report** •
- **DuPage County Influenza Report**
- Kane County Influenza Report •

- Vaccine Finder •
- **Immunization Action Coalition** •
- National Respiratory and Enteric Virus Surveillance System (NREVSS), CDC
- CDC FluView Activity Report and Maps
- International influenza surveillance (WHO)
- St. Louis Children's Hospital Data •