## Table 36

Smoking

Percentage (%) of recent mothers in Illinois who reported a health care worker advised them to quit smoking

## Among women who reported cigarette smoking in the three months before pregnancy and who had a prenatal visit

Survey Question 34: During any of your prental care visits, did a doctor, nurse, or other health care worker advise you to quit smoking?

	Percentage <sup>1</sup>	95% CI <sup>2</sup>		Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>
No	20.4	14.6	27.8	179	3,711
Yes	79.6	72.2	85.4	179	14,474

Percentage (%) of recent mothers in Illinois reporting that a health care worker did any of the following things to address quitting smoking

## Among women who reported cigarette smoking in the three months before pregnancy and who had a prenatal visit

Survey Question 35: Listed below are some things about quitting smoking that a doctor, nurse, or other health care worker might have done during any of your prenatal care visits. Response: Yes

your prenatal care visits. Response. Tes	Percentage <sup>1</sup> 49.1	95% CI <sup>2</sup>		Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>
Spend time with me discussing how to quit smoking		40.9	57.3	177	8,789
Provide me with booklets, videos, or other materials to help me quit smoking on my own	37.6	30.0	45.9	176	6,693
Suggest that I set a specific date to stop smoking	29.9	22.8	38.0	176	5,317
Ask if a family member or friend would support my decision to quit	28.6	21.7	36.6	176	5,093
Suggest I attend a class or program to stop smoking	16.3	11.0	23.6	176	2,906
Recommend using a nicotine patch	14.6	9.6	21.5	175	2,594
Refer me to a national or state quit line	14.1	9.2	21.0	176	2,508
Refer me to counseling for help with quitting	13.4	8.6	20.2	176	2,387
Recommend using nicotine gum	12.2	7.6	19.0	175	2,162
Prescribe a nicotine nasal spray or nicotine inhaler					
Prescribe a pill like Zyban <sup>®</sup> (also known as Wellbutrin <sup>®</sup> or bupropion) to help me quit					
Prescribe a pill like Chantix <sup>®</sup> (also known as varenicline) to help me quit					

<sup>1</sup>**Percentage** = Weighted percentage

 $^{2}\mathbf{CI}$  = Confidence interval lower/upper bounds

-- Indicates denominator <30 respondents or numerator <6 respondents

<sup>3</sup>**Respondents** = The total number who responded to the survey question

<sup>4</sup>Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition