

## Birth Control Use After Delivery

Percentage (%) of recent mothers in Illinois who reported using the following birth control methods

Among women reporting using birth control now

Survey Question 64: What kind of birth control are you or your husband or partner using now to keep from getting pregnant? Response: Yes

	Percentage <sup>1</sup>	95% CI <sup>2</sup>		Respondents <sup>3</sup>	Estimated Population Affected <sup>4</sup>
Condoms	34.8	31.6	38.2	965	39,456
Birth control pills	22.8	20.0	25.9	965	25,868
IUD (including Mirena®, ParaGard®, Liletta®, or Skyla®)	14.1	11.9	16.8	965	16,042
Withdrawal (pulling out)	14.1	11.9	16.7	965	16,037
Tubes tied or blocked (female sterilization or Essure®)	7.1	5.5	9.1	965	8,060
Shots or injections (Depo-Provera®)	6.3	4.8	8.3	965	7,151
Not having sex (abstinence)	9.8	7.9	12.1	965	11,103
Natural family planning (including rhythm method)	7.2	5.6	9.1	965	8,124
Contraceptive implant in the arm (Nexplanon® or Implanon®)	4.4	3.1	6.1	965	4,988
Vasectomy (male sterilization)	3.0	2.1	4.4	965	3,431
Contraceptive patch (OrthoEvra®) or vaginal ring (NuvaRing®)	2.2	1.4	3.5	965	2,504

<sup>1</sup>Percentage = Weighted percentage<sup>2</sup>CI = Confidence interval lower/upper bounds

-- Indicates denominator &lt;30 respondents

<sup>3</sup>Respondents = The total number who responded to the survey question<sup>4</sup>Estimated Population Affected = Estimated number of Illinois mothers/infants with characteristic or condition