

Having Sex? Learn more about the Zika Virus

In some places, Zika virus is primarily spread by infected mosquitoes. However, it can also be spread by having sex (vaginal, oral, anal) with a person who has Zika virus.

If a woman is pregnant and gets Zika virus, her baby can be born with serious birth defects.



To avoid Zika virus and/or a Zika virus-affected pregnancy, talk to your health care provider about your family planning options.



For more information about Zika virus, contraceptive services, or to plan a healthy pregnancy, call the Women's Health Line: 888-522-1282.



www.dph.illinois.gov/topics-services/diseases-and-conditions/zika for more information.




Ou pral fè sèks? Aprann plis sou viris Zika a

Nan kèk kote, viris Zika a pwopaje atravè moustik ki enfekte. Men, li ka pwopaje tou, lè ou ap fè sèks (nan vajen, nan bouch, nan dèyè) ak yon moun ki gen viris Zika a.

Si yon fanm ansent e li pran viris Zika a, pitit li a ka fèt ak defòmasyon nan nesans li.



Pou evite viris Zika a e/oubyen yon gwosès ke Zika afekte, pale ak founisè swen lasante sou opsyon planin fanmi.



Pou plis enfòmasyon sou viris Zika a, sèvis kontraseptif, ou planifye yon gwosès an sante, kontakte nou: 888-522-1282



Vizite

www.dph.illinois.gov/topics-services/diseases-and-conditions/zika pou plis enfòmasyon.