



# 2009 Report

## Illinois Pregnancy Risk Assessment Monitoring System (PRAMS)

# **2009 Report**

## **Illinois Pregnancy Risk Assessment Monitoring System (PRAMS)**

**Illinois Department of Public Health  
Illinois Center for Health Statistics  
PRAMS Program  
866-643-7194 (voice and TTY)  
or  
217-785-1064**

### **Acknowledgements**

The Illinois PRAMS program thanks all of the mothers who took the time to participate. The valuable information they provided has contributed to the understanding of why some babies are born healthy and others are not.

Illinois PRAMS thanks the PRAMS team at the U. S. Centers for Disease Control and Prevention for its technical support, review and comments.

This publication was supported by Cooperative Agreement Number 1UO1DP003109-01 from the U.S. Centers for Disease Control and Prevention.

# Table of Contents

<b>Introduction</b> .....	1
<b>Highlights, 2009</b> .....	2
<b>Section 1. Preconception Health</b> .....	3
Vitamin Use During the Month Before Pregnancy .....	4
Health Behaviors During the 12 Months Before Becoming Pregnant .....	5
<b>Section 2. Family Planning.</b> .....	6
Intendedness of Pregnancy .....	7
Unintended Pregnancy by Selected Maternal Characteristics .....	8
Unintended Pregnancy, 1998-2009 .....	9
Reasons for Not Using Birth Control Among Women Not Trying to Get Pregnant.....	10
<b>Section 3. Prenatal Care</b> .....	11
Initiation of Prenatal Care in the First Trimester by Selected Maternal Characteristics .....	12
Problems Getting Prenatal Care Among Women Who Didn't Get Care as Early They Wanted .....	13
Topics Discussed by Health Care Professionals During Prenatal Care Visits .....	14
Discussions About Nutrition and Weight Management During and After Pregnancy.....	15
Knowledge About Folic Acid by Selected Maternal Characteristics .....	16
Oral Health Care During Pregnancy.....	17
WIC Participation.....	18
HIV Testing During Pregnancy.....	19
HIV Testing During Pregnancy by Selected Maternal Characteristics .....	20
HIV Testing During Pregnancy, 1998-2009.....	21
<b>Section 4. Risk Factors</b> .....	22
Problems During Pregnancy.....	23
Stressors Prior to Delivery.....	24
Types of Stressors During the Year Prior to Delivery.....	25
Physical Abuse Before and During Pregnancy.....	26
Prevalence of Smoking Before, During and After Pregnancy.....	27
Prevalence of Smoking During the Last Three Months of Pregnancy by Selected Sociodemographic Characteristics .....	28
Prevalence of Smoking During the Last Three Months of Pregnancy, 1998-2009 .....	29
Prevalence of Drinking Before and During Pregnancy .....	30
Amount of Drinking Before and During Pregnancy .....	31
Prevalence of Drinking During the Last Three Months of Pregnancy by Selected Sociodemographic Characteristics .....	32
Self-Reported Symptoms of Depression and Anxiety After Childbirth.....	33
Depression Diagnosis During and After Pregnancy.....	34
Postpartum Depression Diagnosis by Selected Sociodemographic Characteristics .....	35
<b>Section 5. Infant Care and Health</b> .....	36
Infant Checkups.....	37
Place of Well-baby Checkup .....	38
Infant Sleep Practices .....	39
Sleep Position of Infant .....	40
Prevalence of Laying Infant on Back to Sleep by Selected Maternal Characteristics.....	41

Prevalence of Laying Infant on Back to Sleep, 1998-2009 .....	42
Breastfeeding Initiation and Continuation.....	43
Breastfeeding Initiation by Selected Sociodemographic Characteristics .....	44
Breastfeeding Initiation and Continuation, 1998-2009 .....	45
Reasons for Not Breastfeeding.....	46
How Did You Learn to Install and Use Your Infant Car Seat? .....	47
<b>Section 6. Insurance</b> .....	<b>48</b>
Source of Payment for Prenatal Care and Delivery.....	49
Type of Insurance for Baby.....	50

**Appendix A. Technical Notes**

**Appendix B. Sample Numbers, Response Rates and Weighted Estimates by Selected Characteristics**

**Appendix C. Detailed Tables**

## **Introduction**

The Illinois Pregnancy Risk Assessment Monitoring System (PRAMS) is an ongoing population-based survey of Illinois women who have delivered a live born infant in Illinois. PRAMS is an important part of the Illinois Department of Public Health's surveillance activities and is part of an initiative by the U.S. Centers for Disease Control and Prevention (CDC) to reduce infant mortality and adverse birth outcomes. The unique information collected by the PRAMS project is used by health professionals, administrators, policy makers and researchers to develop and modify programs and policies to improve the health of women and children in Illinois.

PRAMS collects information from mothers about their behaviors and experiences before, during and immediately following the birth of the baby. The 2009 report covers a variety of selected topics broken down into several categories: preconception health, family planning, prenatal care, risk factors, infant care and health, and insurance. Trend data are presented for selected topics. Data are presented graphically in the body of the report with detailed tables containing standard errors and confidence intervals found in Appendix C. Statewide estimates, as well as subgroup analyses by selected sociodemographic categories, are presented. All results are weighted estimates reflective of Illinois resident women who delivered a live birth in Illinois. The data methods used by Illinois PRAMS are standardized CDC protocols used by all participating states (see Appendix A for technical notes).

# Illinois PRAMS Facts For 2009

## Preconception Health

- During the month before pregnancy 30 percent of women took a multivitamin, prenatal vitamin or folic acid vitamin every day of the week. 13 percent took a vitamin between one to six times per week, and 57 percent didn't take a vitamin at all.
- During the year prior to pregnancy:
  - 41 percent of women said they were exercising three or more days of the week and 27 percent were dieting to lose weight.
  - 52 percent of women had their teeth cleaned by a dentist or dental hygienist.
  - 11 percent of women visited a health care worker to be treated or checked for diabetes, depression or anxiety. 12 percent visited a health care worker to be checked for high blood pressure.
  - 28 percent of women talked with a health care worker about family medical history.

## Family Planning

- 44 percent of pregnancies resulting in live births were reported as unintended.
- 50 percent of women not trying to get pregnant were not using birth control at the time of conception.

## Prenatal Care

- 82 percent of women heard or read about the importance of taking folic acid to prevent some birth defects.
- 69 percent of women reported being tested for HIV during their pregnancy, while 14 percent did not know if they were tested.
- 44 percent of women went to a dentist or dental clinic during their pregnancy and 47 percent reported being counseled by a dental or other health care worker about care of teeth and gums.
- During pregnancy, 7 percent of women were told by a health care professional they had depression.
- 75 percent of women reported that a health care worker talked with them about what to do if they felt depressed during their pregnancy or after the baby was born.

## Risk Factors

- 3 percent of women reported being physically abused by their husband or partner in the 12 months before their pregnancy; 2 percent reported abuse during pregnancy.
- 21 percent of women smoked cigarettes during the three months before they became pregnant, and 9 percent smoked during the last three months of pregnancy.
- 53 percent of women reported drinking alcoholic beverages on a weekly basis in the three months before becoming pregnant. That figure dropped to 7 percent during the last three months of pregnancy. During both time periods, most women reported drinking three or fewer drinks per week.
- Since their new baby was born, 8 percent of women reported they often or always felt down, depressed or sad. During the same time period, 9 percent of women were told by a health care professional they had depression.

## Infant Health and Care


- When asked how they learned to install and use an infant car seat, 27 percent of women reported that a health or safety professional showed them. Many women indicated they read the instructions (55%) or they already knew how to install the seat because they have other children (49%).
- 72 percent of women reported they most often put their baby to sleep on his or her back.
- 80 percent of women reported breastfeeding their baby at least once after birth and 67 percent were still breastfeeding one month after delivery.
- Among women who did not breastfeed, the most common reasons for not doing so were that they did not like breastfeeding (46%) or didn't want to breastfeed (29%).

# Section 1

## Preconception Health



### Vitamin Use During the Month Before Pregnancy, 2009




As part of good preconception health care, the CDC recommends that women take 400 mcg of folic acid a day for at least three months before becoming pregnant to reduce the risk of birth defects<sup>1</sup>. PRAMS found that more than half of Illinois women said they did not take a multivitamin, prenatal vitamin, or folic acid vitamin during the month before getting pregnant. Only 30.1 percent of women reported taking a vitamin daily.

Older women were more likely to take vitamins every day of the week when compared with younger women. Black women were significantly less likely to take vitamins daily when compared with other races. Women with more than a high school education were more likely to take vitamins when compared with lesser educated women. Non-Hispanic and married women were more likely to take vitamins than Hispanic and unmarried women. Women whose deliveries were paid by Medicaid were less likely to take vitamins on a daily basis when compared with women whose deliveries were not paid for by Medicaid. See Appendix C, Table 1 for detailed statistics.

---

<sup>1</sup> <http://www.cdc.gov/ndbddd/preconception/QandA.htm> accessed September 9, 2011.

### Health Behaviors During the 12 Months Before Pregnancy, 2009



CDC recommends that women prepare for pregnancy at least three months before conception<sup>2</sup>. The American College of Obstetricians and Gynecologists (ACOG) recommends that women who are planning to become pregnant make an appointment with their doctor to discuss items such as lifestyle, diet, medical history, medications, and diseases that run in the family<sup>3</sup>. Prior to pregnancy, women should strive to attain a healthy weight through proper diet and exercise, maintain good oral health, refrain from smoking and drinking alcohol, talk to a doctor or pharmacist about medications, and make sure any medical conditions are managed and under control.

PRAMS asked women about some of the items related to preconception health during the 12 months before becoming pregnant. Slightly more than half of women said they had their teeth cleaned and 41.2 percent were exercising three or more days of the week. About one in four women had spoken with their doctor about family medical history and about the same percentage were dieting to lose weight. Fewer than 20 percent of women indicated they were taking prescription medication or birth control, or had visited a health care worker to be checked or treated for high blood pressure, depression, anxiety or diabetes.

<sup>2</sup> <http://www.cdc.gov/ndbddd/preconception/QandA.htm> accessed September 9, 2011.


<sup>3</sup> [http://www.acog.org/publications/patient\\_education/bp056.cfm](http://www.acog.org/publications/patient_education/bp056.cfm) accessed September 9, 2011.

## Section 2

### Family Planning




## Intendedness of Pregnancy, 2009



PRAMS asked women how they felt about the timing of becoming pregnant. Women who indicated that they wanted to be pregnant “then” or “sooner” were classified as having an intended pregnancy (55.7%). Those who answered that they wanted to be pregnant “later” or “not then or at any time in the future” were considered to have an unintended pregnancy (44.3%). The Healthy People 2010 Objective (9-1) was to increase intended pregnancies to 70 percent. PRAMS found three out of four unintended births in Illinois were mistimed, meaning that the woman would have preferred being pregnant at a later time. PRAMS also found that among women who were not trying to get pregnant, 50.3 percent were not using contraception at the time of conception. The Healthy People 2010 Objective (9-3) was to increase the proportion of females at risk of unintended pregnancy (and their partners) who use contraception to 100 percent.


### Unintended Pregnancy, 2009 By Selected Maternal Characteristics



When examining subpopulations of women, the percentage of unintended pregnancies varied widely and was very high among certain groups. Teenagers exhibited the highest percentage of unintended pregnancies when compared to other age groups. Black women had much higher percentages of unintended pregnancies when compared to women of other races. Women with a high school education or less had significantly\* more unintended pregnancies when compared to women with more than a high school education. Women whose deliveries were paid for by Medicaid and women who were unmarried had more than double the percentage of unintended pregnancies when compared to women whose deliveries were not paid by Medicaid and married women.


\*Statements throughout this report regarding significance refer to statistical significance as determined through the use of confidence intervals. See Appendix A for more details about interpretation of PRAMS data.

### Unintended Pregnancy, 1998-2009 With 95 Percent Confidence Intervals



The above chart illustrates that during the period of 1998-2009, a decline in unintended pregnancies was not seen in Illinois. The percentage peaked at 46.2 in 2001 and has remained above 40 percent since 1999. Rates among women who were unmarried, young, black, less educated or on Medicaid remained much higher than 40 percent over this time span (see tables 4a-4c, Appendix C).


### Reasons for Not Using Birth Control at Conception Among Women not Trying to Get Pregnant, 2009




PRAMS asked about reasons for not using birth control among women who said they were not trying to get pregnant. A list of reasons was provided and women were asked to check all that applied to their situation. Women checked most often that either they did not mind if they got pregnant or they thought they could not get pregnant.

## Section 2

### Prenatal Care




**Initiation of Prenatal Care in the First Trimester, 2009  
By Selected Maternal Characteristics**




The Healthy People 2010 Objective 16-16 was to increase to at least 90 percent the proportion of all pregnant women who receive prenatal care in the first trimester of pregnancy. Among all Illinois women, 82.3 percent reported having their first prenatal care visit during the first trimester of their pregnancies. Some groups of women met the 90 percent objective while others fell far short of it as is illustrated in the chart above. Characteristics of women who met the objective included: married and delivery not paid for by Medicaid.

## Problems Getting Prenatal Care Among Women Who Didn't Get Care as Early as They Wanted, 2009




PRAMS asked women if they got prenatal care as early in their pregnancy as they wanted. More than 84 percent of women indicated they did get care as early as they wanted while 15.7 percent said they did not. Women who did not get care as early as they wanted were provided a list of reasons for not getting prenatal care early and asked to indicate if each item applied to them. As reflected in the chart above, the reasons most frequently indicated included not being able to get an appointment when wanted, not knowing they were pregnant or not having means to pay for the visit.

## Topics Discussed by Health Care Professionals During Prenatal Care Visits, 2009




PRAMS provided a list of topic areas that may have been discussed at prenatal care visits and asked women to respond to each item with “yes” if the subject was discussed or “no” if the subject was not discussed. Women were asked not to mark “yes” if only reading materials or a video was provided. About 75 percent of women indicated that a health care professional had discussed 10 of the 13 topics. The least discussed topic areas included use of seatbelts during pregnancy and intimate partner physical abuse.

## Discussions About Nutrition and Weight Management During and After Pregnancy, 2009




PRAMS asked women if a doctor, nurse or other health care worker spoke with them about any of the items listed in the chart above. The majority of women were counseled about how much weight to gain, exercise during pregnancy and foods that are good to eat during pregnancy. Women were much less likely to be counseled about programs and resources available either to help gain the right amount of weight during pregnancy or to lose weight after pregnancy.

### Knowledge About Folic Acid, 2009 By Selected Maternal Characteristics



Overall, 81.8 percent of women heard or read that folic acid could prevent some birth defects. However, significant differences were seen among subpopulations of women as highlighted in the chart above. Older or more educated women were more likely to be aware of the benefits of folic acid when compared to younger or less educated women. Black women were less likely to be aware of the benefits when compared with women of other races. Women who were married or whose deliveries were not paid for by Medicaid were more likely to report awareness when compared with unmarried women or women whose deliveries were paid for by Medicaid.

## Oral Health Care During Pregnancy, 2009




Maintaining good oral health during pregnancy is important as researchers have found that women with periodontal disease are at an increased risk for delivering preterm low birthweight babies<sup>4</sup>. Pregnant women should see a dentist or dental hygienist during pregnancy for a preventive cleaning<sup>5</sup>. In 2009, 44 percent of women in Illinois said they visited a dentist or dental clinic during pregnancy. More than 47 percent reported that a health care worker spoke with them about the care of their teeth and gums. Less than one-fourth of women said they needed to see the dentist for a problem. (See table 13, Appendix C, for more details by sociodemographic factors.)

<sup>4</sup> Illinois Department of Public Health. *Women's Oral Health*. n.d. <http://www.idph.state.il.us/HealthWellness/oralhlth/oralwomen.htm> (accessed February 18, 2009).

<sup>5</sup> Illinois Department of Public Health. *Oral Health: A Link to General Health*. 2004.


### WIC Participation, 2009



PRAMS found that 46.8 percent of women reported participating in the WIC\* program during pregnancy, while 54.6 percent participated after delivery. Characteristics of women most likely to participate in the program included being unmarried, young, of black race, of Hispanic ethnicity, or having a high school education or less (see table 14, Appendix C).


\*WIC is the Special Supplemental Nutrition Program for Women, Infants and Children funded by the U. S. Department of Agriculture (USDA) and administered in Illinois by the Illinois Department of Human Services.

## HIV Testing During Pregnancy, 2009



In 2009, 69.5 percent of women indicated they were tested for HIV during pregnancy, 17.0 percent were not tested and 13.5 percent did not know if they were tested.


### HIV Testing During Pregnancy, 2009 By Selected Maternal Characteristics



In 2009, black women were significantly more likely to be tested for HIV when compared to women of other races, and unmarried women were more likely to be tested than married women. Women whose deliveries were paid for by Medicaid were more likely to be tested than women whose deliveries were not paid for by Medicaid. Lesser educated women reported higher percentages of testing when compared with women with more education although the differences were not statistically significant. When examining age groups, the youngest age group reported highest percentages of testing, although the differences between the age groups were not statistically significant.

## HIV Testing During Pregnancy, 1998-2009

With 95 Percent Confidence Intervals




\*Change in question; see appendix A.


When examining HIV testing during the period of 1998 through 2009, PRAMS found the highest reported testing rates occurred between the years 2005 and 2008. In 2009 the overall rate dropped significantly to 69.5 percent from 76 percent in 2008. Throughout the period, black women were more likely to be tested than women of other races. Unmarried women and women whose deliveries were paid for by Medicaid reported higher testing rates than married women and women whose deliveries were not paid for by Medicaid. In general, younger women reported higher rates of testing than older women (see tables 12a-c, Appendix C).

## Section 3

### Risk Factors




### Problems During Pregnancy, 2009



PRAMS asked women if they experienced certain problems during pregnancy by providing a list and asking women to respond “yes” or “no” to each item. From the list provided, the most common problem reported was severe nausea, vomiting or dehydration. The next most common problems reported were kidney/bladder infections, vaginal bleeding, or preterm labor (starting more than three weeks before due date). Fewer than 2 percent of women reported their cervix was sewn shut or they were hurt in a car accident.

### Stressors Prior to Delivery, 2009




High levels of stress during pregnancy can contribute to poor pregnancy outcomes<sup>6</sup>. PRAMS asked about stressful events that occurred during the 12 months prior to delivery and found that 29.3 percent of women reported experiencing no major stressful events while about 41.3 percent reported experiencing one to two stressors. Three or more stressors were reported by 29.4 percent.


---

<sup>6</sup> D'Angelo, D; Williams, L; Morrow, B; Cox, S; Harris, N; Harrison, L; Posner, SF; Richardson Hood, J; and Zapata, L. *Preconception and Interconception Health Status of Women Who Recently Gave Birth to a Live-Born Infant---Pregnancy Risk Assessment Monitoring System (PRAMS), United States, 26 Reporting Areas, 2004*. In: *MMWR Surveillance Summaries, December 14, 2007*; 56 (SS10): 1-35.

## Types of Stressors Prior to Delivery, 2009




PRAMS provided a list of situations that could cause stress and asked women to indicate if any of the situations happened to them during the 12 months before the baby was born. Overall, the most commonly reported event was moving to a new address (27.8%). About one in four women reported having more arguments with a husband/partner. About 23 percent of women reported having a very sick family member or having a lot of bills that could not be paid.




PRAMS asked women about physical abuse by a husband/partner both before and during pregnancy. Nearly 3 percent of women reported abuse before pregnancy, while 2.1 percent reported abuse during pregnancy. Regardless of time period, unmarried women and women whose deliveries were paid for by Medicaid were more likely to report abuse when compared with married women or those whose deliveries were not paid for by Medicaid. Also, reporting of abuse was more prevalent as education of mother decreased (see table 19, Appendix C).

### Prevalence of Smoking Before, During and After Pregnancy, 2009




Healthy People 2010 Objective 16-17c was to reduce cigarette smoking to a prevalence of no more than 1 percent among pregnant women. In Illinois, PRAMS found that during the three months before pregnancy, 20.9 percent of women reported they were smoking. This decreased to 9.2 percent during the last three months of pregnancy. After delivery, reported smoking climbed to 14.6 percent. Significant differences were seen when comparing different groups of women (see chart on next page).

**Prevalence of Smoking  
During the Last Three Months of Pregnancy, 2009  
By Selected Sociodemographic Characteristics**




Women who had given birth to low birthweight babies reported higher percentages of smoking during pregnancy when compared to women who gave birth to normal birthweight babies. Non-Hispanic women were more likely to smoke when compared to Hispanic women. Women in their early 20s reported higher percentages of smoking when compared to older women. Women with a high school education or less were more likely to smoke when compared to women with more than a high school education. Unmarried women and women whose deliveries were paid for by Medicaid reported much higher rates of smoking when compared to married women and women whose deliveries were not paid for by Medicaid. (For more details by sociodemographic characteristics and by time period, see table 20a, Appendix C).

**Prevalence of Smoking  
During the Last Three Months of Pregnancy, 1998-2009**  
With 95 Percent Confidence Intervals




\*Change in question; see appendix A.

From 1998 to 2009, there was an overall gradual decline in the percentage of reported smoking during pregnancy. Gaps remained between groups from 1998 to 2009 with black and white women more likely to smoke than women of other races, and non-Hispanic women more likely to smoke than Hispanic women. Unmarried women and those with a high school education or less were more likely to smoke than married women and those with more than a high school education. Women whose deliveries were paid for by Medicaid were more likely to smoke than women whose deliveries were not paid for by Medicaid (see tables 20a-d, Appendix C).




As part of good preconception health, CDC recommends that women stop drinking before getting pregnant. Healthy People 2010 Objective 16-17a was to increase abstinence from alcohol by pregnant women to at least 94 percent. PRAMS asked women about alcohol use just prior to pregnancy and late into pregnancy. In Illinois, slightly more than half of women reported drinking in the three months before pregnancy while 6.8 percent reported drinking during the last three months of pregnancy.



Women reported drinking more drinks per week prior to pregnancy when compared with during pregnancy. Among women who reported drinking during pregnancy, 95.1 percent drank three or fewer drinks per week.


**Prevalence of Drinking  
During the Last Three Months of Pregnancy, 2009  
By Selected Sociodemographic Characteristics**



\*Fewer than six survey respondents indicated they were drinking during pregnancy

PRAMS found that older women were generally more likely to report any drinking during pregnancy than younger women. Non-Hispanic women and women with more than a high school education were more likely to report drinking when compared with Hispanic women or women with a high school education or less. Women whose deliveries were not paid for by Medicaid reported higher percentages of drinking during pregnancy when compared with women whose deliveries were paid for by Medicaid (see table 21, Appendix C). Women who gave birth to a normal birthweight baby were more likely to report any drinking when compared with women who gave birth to a low birthweight baby.


## Self-Reported Symptoms of Depression and Anxiety After Childbirth, 2009



Women were given a list of feelings and experiences that women sometimes have after childbirth and asked to describe how often (using a scale of Never, Rarely, Sometimes, Often, or Always) they felt or experienced things this way since the new baby was born.


Three of the items PRAMS asked about related to depressive symptoms. These included feeling down, depressed or sad, feeling hopeless, or feeling slowed down. More than a quarter of women reported that they sometimes felt down, depressed, sad or slowed down. More than 13 percent of women reported that they often or always felt slowed down and 8.3 percent of women said they often or always felt down, depressed or sad. More than 8 percent of women said they sometimes felt hopeless, while 4.6 often or always felt hopeless.

Two of the items PRAMS inquired about related to anxiety, including feeling panicky and feeling restless. About one in five women said they sometimes felt restless while 11.2 percent of women said they often or always felt restless. More than 10 percent of women said they sometimes felt panicky while 5.2 percent said they often or always felt panicky.



PRAMS asked women if they were told by a health professional (doctor, nurse or other health care worker) they had depression either during pregnancy or after the baby was born. The chart above illustrates that 7.4 percent of women indicated they had been told they had depression during pregnancy while 9.1 percent indicated they were told they had depression after the baby was born. (See tables 24 and 25 in Appendix C for sociodemographic details.)

## Postpartum Depression Diagnosis, 2009 By Selected Sociodemographic Characteristics




After the baby was born, 9.1 percent of new mothers indicated to PRAMS that they had been diagnosed with depression. The above chart illustrates differences in reported diagnosis of depression among different groups of women. Significant differences in diagnosis were seen when examining ethnicity and type of payment for delivery. Non-Hispanic women were more likely to report being diagnosed with depression when compared with Hispanic women. Women whose deliveries were paid for by Medicaid were more likely to report being diagnosed with depression when compared with women whose deliveries were not paid for by Medicaid.

## **Section 4**

### **Infant Care and Health**




### Infant Checkups, 2009




PRAMS asked women about infant checkups and found that 95.6 percent of babies were seen within the first week after leaving the hospital and nearly all babies (99.0%) had a regular health visit at 2, 4 or 6 months of age.

### Place of Well- baby Checkup, 2009



Women were asked where they usually take their babies for well-baby checkups. About three in five indicated they took their baby to a medical doctor (MD)/health maintenance organization (HMO). 16.9 percent took their baby to a hospital clinic. The remaining women took their babies to community health centers, health department clinics, or other places for care.


## Infant Sleep Practices, 2009



In addition to placing an infant to sleep on his or her back, a number of other infant sleep practices have been recommended in an effort to reduce risk factors that could contribute to Sudden Infant Death Syndrome (SIDS). These are to use a safety approved crib with a firm mattress, to not use bumper pads, and to remove all loose bedding such as pillows, stuffed toys, and blankets from the crib. Also, co-sleeping (both baby and adults or other children sharing the same bed or couch) is not recommended<sup>7</sup>. PRAMS provided a list of items that described sleep practices and asked moms to indicate if each item usually applied to their baby. The chart above shows that a majority of women indicated that their baby usually sleeps in a crib or portable crib. The majority of women also indicated that their baby slept on a firm or hard mattress. Nearly 42 percent of women indicated their baby sleeps with bumper pads, while nearly 23 percent said their baby sleeps with plush blankets. Nearly 18 percent of women indicated that their baby usually sleeps with another person. Slightly more than 10 percent of women indicated their baby usually sleeps with pillows, while fewer than 5 percent sleep with stuffed toys.


<sup>7</sup>[http://www.idph.state.il.us/sids/sids\\_factsheet.htm](http://www.idph.state.il.us/sids/sids_factsheet.htm) accessed November 10, 2011.

### Sleep Position of Infant, 2009




Healthy People 2010 objective 16-13 was to increase the percentage of healthy full-term babies who are put to sleep on their backs to at least 70 percent. While overall, Illinois met this objective starting in 2006, there are disparities among subpopulations of women. Many groups have not yet met the 70 percent objective. These women were more likely to be young, not married, of black race, and of Hispanic ethnicity. They also were more likely to have a high school education or less, and the delivery paid for by Medicaid (see chart on the next page).

**Prevalence of Laying Infant on Back to Sleep, 2009  
By Selected Maternal Characteristics**



PRAMS found generally that older women were more likely to put their babies to sleep on their backs when compared with younger women. Non-black women also were more likely to put babies to sleep on their backs when compared to black women. Non-Hispanic women were more likely to use the back sleep position than Hispanic women. Women with more than a high school education were more likely to put their babies to sleep on their backs when compared to less educated women. Women who were married or whose deliveries were not paid for by Medicaid were more likely to put their babies to sleep on their backs when compared to women who were not married or whose deliveries were paid for by Medicaid.


**Prevalence of Laying Infant on Back to Sleep, 1998-2009**  
With 95 Percent Confidence Intervals



\*Change in question; see Appendix A.


From 1998 to 2009, there was a significant increase in the prevalence of laying infants to sleep on their backs. While most subpopulations of women reported increases in putting infants to sleep on their backs during this time period, gaps remained between groups. Older and more educated women reported higher percentages than younger, less educated women. Non-black women continued to report higher percentages than black women. Women who were married or whose deliveries were not paid for by Medicaid reported higher percentages of laying infants down to sleep on their backs than women who were not married or whose deliveries were paid for by Medicaid. From 2005 through 2007, significant differences were not seen when comparing Hispanic with non-Hispanic ethnicity. Starting in 2008, however, Hispanic women were again significantly less likely than non-Hispanic women to put their babies to sleep on their backs (see tables 30a-30c, Appendix C).

### Breastfeeding Initiation and Continuation, 2009




PRAMS found that 79.9 percent of women reported breastfeeding their babies at least once after birth, while 66.7 percent of women were still breastfeeding at one month after delivery.

**Breastfeeding Initiation, 2009  
By Selected Sociodemographic Characteristics**



When examining subpopulations of women, many significant differences existed in breastfeeding initiation. Older women and those with more than a high school education had higher initiation rates than younger, less educated women. Non-black or Hispanic women were significantly more likely to breastfeed than black or non-Hispanic women. Women who were married or whose deliveries were not paid for by Medicaid had significantly higher rates of initiation when compared with unmarried women or those whose deliveries were paid for by Medicaid. Similar patterns could be seen with continuation at one month (see tables 31 and 32, Appendix C).


### Breastfeeding Initiation and Continuation, 1998-2009



\*Change in questions; see appendix A.


From 1998-2009, breastfeeding initiation and continuation at one month followed similar patterns, peaking in 2009. Throughout the period, gaps remained among groups. Older women and those with more than a high school education were more likely to initiate and continue breastfeeding than their younger, less educated counterparts. Non-black or Hispanic women were significantly more likely to breastfeed than black or non-Hispanic women. Married women or women whose deliveries were not paid for by Medicaid were more likely to initiate and continue breastfeeding when compared to unmarried women or those whose deliveries were paid for by Medicaid.

### Reasons for Not Breastfeeding, 2009



PRAMS provided a list of reasons why women might not initiate breastfeeding and asked women to check all of the reasons that applied to them. The most common reason checked for not breastfeeding was they did not like breastfeeding (45.5%). The second most common reason cited was that women didn't want to breastfeed (29%).

### How Did You Learn to Install and Use Your Infant Car Seat? (2009)




PRAMS asked women if they had an infant car seat for their new baby and nearly all women (99.5%) said “yes.” This figure did not vary significantly across subpopulations of women. Women were then given a list and asked to check all of the ways they learned to install and use the car seat. PRAMS found that women most often reported they read the instructions (55.2%) or already knew how to install the seat because they had other children (49.4%). About one in four women said a health or safety professional showed them how to install the seat, while about one in five women reported that a friend or family member showed them how to install the seat.

## Section 5

### Insurance




### Source of Payment for Prenatal Care and Delivery, 2009



PRAMS asked women how their prenatal care and deliveries were paid. The majority of women said their prenatal care and deliveries were paid for either by Medicaid programs or by health insurance from their job or the job of their husbands, partners or parents. Fewer than 5 percent of women said these services were paid for by military health care, health insurance that was not associated with a job, or other sources.

(Note: Percentages do not add up to 100 as more than one type of payment could have been chosen.)

### Type of Insurance for Baby, 2009



PRAMS asked women if they had health insurance or Medicaid for their new baby and found that 98.1 percent indicated they had some type of coverage. The majority of women indicated that their babies were covered by Medicaid programs or by health insurance from their job or the job of their husbands, partners or parents. Two percent of women said their babies were covered by insurance that was not associated with a job. Fewer than 1 percent (not shown on chart as there were fewer than six respondents) said their babies were covered by military health care or other sources. (Note: Percentages may not add up to 100 as more than one type of insurance could have been chosen.)

# Appendix A

## Technical Notes

### Collection of Data

The data collection for the 2009 Illinois Pregnancy Risk Assessment Monitoring System (PRAMS) was conducted by the Illinois Center for Health Statistics, Illinois Department of Public Health.

Illinois uses the standardized data collection methods developed by the U.S. Centers for Disease Control and Prevention (CDC) and used by all 37 states participating in PRAMS. Data are collected through mailed surveys with telephone follow-up for non-responders. A systematic stratified random sample of approximately 200 mothers is selected monthly from a frame of eligible birth certificates. At two to six months after delivery, each sampled mother is mailed an introductory letter followed up by a 14-page survey. A reminder letter and a second and third mailing of the survey are sent to those who do not respond. PRAMS interviewers telephone mothers who do not respond to any of the mailed surveys to administer the survey by telephone. Surveys are available in both English and Spanish, and telephone interviews were conducted in both languages.

A total of 2,233 women were eligible to participate in the study in 2009; 1,557 women actually participated. Sample numbers, response rates and weighted estimates by selected characteristics are presented in Appendix B.

### Sampling Design

A systematic stratified random sampling design is used to oversample low birthweight (less than 2,500 grams) births. The sample is divided into two strata: low birthweight (less than 2,500 grams) and normal birthweight (2,500 grams or greater). The weighted response rates in 2009 were as follows:

Overall, 70.9 percent

Low birthweight, 65.0

Normal birthweight, 71.3

### Weighting and Interpretation of Results

Statistics in this report are based on weighted data. The weights were developed by CDC to adjust for sample design, non-response patterns and omissions from the sampling frame. The final sampling weight used in the analysis of the survey data is the product of these three elements. Weighting is necessary to give unbiased estimates of population parameters.

Percentages, standard errors and the 95 percent confidence intervals were calculated using Survey Data Analysis (SUDAAN®\*) software. Confidence intervals were used to determine significance of trends. All missing observations were excluded. Missing values equal to or exceeding 10 percent are noted. Estimates for subpopulations in which there were fewer than 32 respondents are not reported, as they may be imprecise and biased. Estimates for response categories with fewer than six respondents are not reported, as they may be unstable.

PRAMS data are representative of Illinois resident women, age 14 years or older, who have given live birth in Illinois. The sampling design is valid at the state level and not intended or developed to represent sub-state geographies or their populations. The data are not applicable to all pregnant women. Women younger than 18 years of age were not asked questions about physical abuse.

\*SUDAAN® is a registered trademark of the Research Triangle Institute.  
© 2005 Research Triangle Institute. All rights reserved

Periodically, the PRAMS survey undergoes a major revision during which questions may be modified, dropped or new questions added. Such revisions occurred in the years 2000, 2004 and 2009. Changes associated with each survey phase are noted throughout this document when multiyear data are presented. Specific wording changes are presented below:

### **HIV Testing**

Phase 3 and 4 Question (1998-2003):

*At any time during your most recent pregnancy or delivery, did you have blood test for HIV (the virus that causes AIDS)?*

*No*  
*Yes*  
*I don't know*

Phase 5 and 6 Question (2004-2009):

*At any time during your most recent pregnancy or delivery, did you have a test for HIV (the virus that causes AIDS)?*

*No*  
*Yes*  
*I don't know*

### **Smoking During Pregnancy**

Phase 3 and 4 Question (1998-2003):

*In the last three months of your pregnancy, how many cigarettes or packs of cigarettes did you smoke on an average day? (A pack has 20 cigarettes.)*

*\_\_\_\_\_ Cigarettes OR \_\_\_\_\_ Packs*  
*Less than 1 cigarette a day*  
*I didn't smoke*  
*I don't know*

Phase 5 and 6 Question (2004-2009):

*In the last three months of your pregnancy, how many cigarettes did you smoke on an average day? (A pack has 20 cigarettes.)*

*41 cigarettes or more*  
*21 to 40 cigarettes*  
*11 to 20 cigarettes*  
*6 to 10 cigarettes*  
*1 to 5 cigarettes*  
*Less than 1 cigarette*  
*None (0 cigarettes)*

### **Breastfeeding Initiation and Duration**

Phase 3 Question (1998, 1999):

*For how many weeks did you breastfeed your new baby?*

*\_\_\_\_\_ Weeks*  
*I didn't breastfeed my baby*  
*I breastfed less than one week*  
*I'm still breastfeeding*

Phase 4 and 5 Questions (2000-2008):

*Did you ever breastfeed or pump breast milk to feed your new baby after delivery?*

*No*

*Yes*

*How many weeks or months did you breastfeed or pump milk to feed your baby?*

*\_\_\_\_\_Weeks **OR** \_\_\_\_\_Months*

*\_\_\_Less than one week*

Phase 6 Questions (2009):

*Did you ever breastfeed or pump breast milk to feed your new baby after delivery, even for a short period of time?*

*No*

*Yes*

*Are you currently breastfeeding or feeding pumped milk to your new baby?*

*No*

*Yes*

*How many weeks or months did you breastfeed or pump milk to feed your baby?*

*\_\_\_\_\_Weeks or \_\_\_\_\_Months*

*\_\_\_ Less than one week*

### **Infant Sleep Position**

Phase 3 Question (1998, 1999):

*How do you put your new baby down to sleep **most** of the time? **Check one answer.***

*On his or her side*

*On his or her back*

*On his or her stomach*

Phase 4 and 5 Question (2000-2008):

*How do you **most often** lay your baby down to sleep now? **Check one answer.***

*On his or her side*

*On his or her back*

*On his or her stomach*

Phase 6 Question (2009):

*In which one position do you **most often** lay your baby down to sleep now? **Check one answer***

*On his or her side*

*On his or her back*

*On his or her stomach*

## Appendix B

### Sample Numbers, Response Rates and Weighted Estimates by Selected Characteristics

#### Illinois PRAMS, 2009

Characteristic	Number Sampled	Number of Respondents	Unweighted Percent Responding	Weighted Estimate*
<b>All</b>	<b>2,233</b>	<b>1,557</b>	<b>69.7</b>	<b>163,018</b>
<b>Age</b>				
< 20 years	247	153	61.9	16,697
20-24 years	458	299	65.3	32,676
25-34 years	1,196	854	71.4	87,965
35+ years	332	251	75.6	25,681
<b>Race</b>				
White	1,652	1,233	74.6	127,235
Black	418	222	53.1	26,394
Other	158	98	62.0	8,974
<b>Hispanic Ethnicity</b>				
Hispanic	550	388	70.5	41,603
Non-Hispanic	1,682	1,168	69.4	121,388
<b>Education</b>				
< High School	420	289	68.8	32,560
High School	604	368	60.9	40,109
> High School	1,180	881	74.7	88,495
<b>Marital Status</b>				
Married	1,311	976	74.4	98,776
Unmarried	922	581	63.0	64,242
<b>Birthweight</b>				
Low (<2,500 grams)	558	362	64.9	11,970
Normal (>=2,500 grams)	1,675	1,195	71.3	151,048

\*For respondents: weighted estimate in PRAMS eligible population of Illinois resident women, age 14 years or older, who gave live birth in Illinois.

## Appendix C

### Detailed Tables

**Table 1**  
Daily Vitamin Use During the Month Before Pregnancy, 2009  
By Selected Sociodemographic Characteristics

	Percent	Standard Error	95% CI *
<b>Total</b>	30.1	1.2	27.7-32.6
<b>Age</b>			
< 20 years	11.2	2.7	6.9-17.7
20-24 years	14.9	2.3	11.0-19.8
25-34 years	33.9	1.7	30.6-37.3
35+ years	48.5	3.4	41.9-55.2
<b>Race</b>			
Black	16.0	2.8	11.3-22.2
White	32.7	1.4	30.0-35.5
Other	31.6	5.1	22.5-42.4
<b>Hispanic Ethnicity</b>			
Hispanic	20.1	2.2	16.2-24.7
Non-Hispanic	33.5	1.5	30.6-36.4
<b>Education</b>			
< High School	15.7	2.2	11.8-20.6
High School	16.7	2.1	13.0-21.1
> High School	41.0	1.8	37.6-44.5
<b>Marital Status</b>			
Married	40.5	1.7	37.2-43.8
Other	14.0	1.6	11.2-17.3
<b>Payment for Delivery</b>			
Medicaid	16.1	1.4	13.5-18.9
Other	46.5	2.0	42.7-50.4
<b>Birthweight</b>			
Low	29.8	2.4	25.4-34.7
Normal	30.1	1.3	27.6-32.8

\* Confidence interval

**Table 2**  
 Health Behaviors During the 12 Months Before Becoming  
 Pregnant, 2009

	Percent	Standard Error	95% CI *
I had my teeth cleaned by a dentist or dental hygienist	51.9	1.4	49.2-54.3
I was exercising three or more days of the week	41.2	1.4	38.5-43.9
I talked to a health care worker about my family medical history	27.6	1.2	25.3-30.1
I was dieting (changing my eating habits) to lose weight	26.6	1.2	24.3-29.1
I was regularly taking prescription medicines other than birth control	17.3	1.0	15.4-19.4
I visited a health care worker to be checked or treated for depression or anxiety	11.2	0.9	9.6-13.0
I visited a health care worker to be checked or treated for diabetes	10.9	0.9	9.3-12.7
I visited a health care worker to be checked or treated for high blood pressure	11.5	0.9	9.9-13.3

**Table 3**  
Intendedness of Pregnancy, 2009

	Percent	Standard Error	95% CI *
Wanted sooner	18.4	1.0	16.5-20.6
Wanted later	32.7	1.3	30.2-35.3
Wanted then	37.3	1.3	34.8-40.0
Didn't want then or at any time in future	11.6	0.9	9.9-13.5

\* Confidence interval

**Table 4a**  
 Unintended Pregnancy, 2008-2009  
 By Selected Sociodemographic Characteristics

	2008			2009		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI*
<b>Total</b>	41.3	1.3	38.8-43.9	44.2	1.4	41.6-47.0
<b>Age</b>						
< 20 years	78.8	3.2	71.8-84.4	74.4	3.9	66.1-81.3
20-24 years	57.1	2.9	51.4-62.7	66.9	2.9	61.0-72.4
25-34 years	32.5	1.7	29.2-35.9	36.0	1.8	32.6-39.6
35+ years	22.8	2.9	17.6-29.0	24.2	3.0	18.9-30.6
<b>Race</b>						
Black	71.2	3.3	64.4-77.3	74.9	3.3	68.0-80.7
White	35.6	1.4	32.9-38.4	39.0	1.5	36.2-42.0
Other	26.1	5.2	17.3-37.4	29.2	5.1	20.3-40.0
<b>Hispanic Ethnicity</b>						
Hispanic	45.2	2.5	40.4-50.1	47.3	2.7	42.0-52.6
Non-Hispanic	39.8	1.6	36.8-42.9	43.2	1.6	40.1-46.3
<b>Education</b>						
< High School	57.7	3.0	51.8-63.4	57.8	3.1	51.6-63.8
High School	53.2	2.6	48.0-58.2	58.9	2.8	53.3-64.2
> High School	29.0	1.7	25.9-32.4	33.0	1.7	29.7-36.4
<b>Marital Status</b>						
Married	23.5	1.4	20.8-26.4	28.3	1.6	25.4-31.5
Other	67.0	2.0	62.9-70.8	68.8	2.1	64.6-72.7
<b>Payment for Delivery</b>						
Medicaid	58.1	1.9	54.4-61.7	59.6	1.9	55.9-63.2
Other	23.6	1.6	20.6-26.9	26.0	1.8	22.7-29.6
<b>Birthweight</b>						
Low	42.1	2.5	37.3-47.1	46.6	2.7	41.4-51.9
Normal	41.3	1.4	38.5-44.1	44.1	1.5	41.2-47.0

\* Confidence interval

**Table 4b**  
 Unintended Pregnancy, 2003-2007  
 By Selected Sociodemographic Characteristics

	2003			2004			2005			2006			2007		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	43.8	1.4	41.1-46.5	41.3	1.2	38.9-43.8	42.2	1.3	39.6-44.8	41.7	1.3	39.1-44.4	42.9	1.3	40.4-45.5
<b>Age</b>															
< 20 years	77.9	3.5	70.2-84.1	75.0	3.7	67.2-81.5	75.7	3.5	68.1-81.9	77.4	3.6	69.5-83.7	80.6	3.3	73.3-86.3
20-24 years	61.9	2.7	56.6-67.0	58.5	2.7	53.1-63.6	58.1	2.9	52.4-63.6	55.1	2.9	49.3-60.7	60.1	2.8	54.5-65.5
25-34 years	31.2	1.8	27.8-34.7	33.5	1.6	30.5-36.7	33.0	1.8	29.7-36.6	35.8	1.7	32.5-39.3	33.9	1.7	30.7-37.3
35+ years	31.7	3.5	25.2-39.1	26.2	2.8	21.1-32.0	27.9	3.1	22.3-34.4	19.7	2.7	14.9-25.6	22.4	2.9	17.3-28.7
<b>Race</b>															
Black	69.1	3.2	62.5-75.0	71.4	2.8	65.5-76.6	69.6	3.1	63.2-75.3	67.5	3.3	60.7-73.7	68.5	3.1	62.1-74.3
White	39.3	1.5	36.4-42.3	34.9	1.3	32.3-37.5	36.3	1.4	33.5-39.2	36.8	1.4	34.0-39.6	37.7	1.4	34.9-40.5
Other	25.8	5.5	16.5-37.9	36.2	5.7	25.8-48.0	31.9	6.2	21.2-45.0	29.7	5.3	20.5-40.9	29.2	4.6	21.0-39.0
<b>Hispanic Ethnicity</b>															
Hispanic	45.2	2.7	40.0-50.6	42.5	2.5	37.7-47.6	44.2	2.6	39.1-49.4	44.0	2.5	39.2-49.0	47.5	2.8	42.1-52.9
Non-Hispanic	43.3	1.6	40.3-46.4	41.0	1.4	38.2-43.8	41.6	1.6	38.6-44.7	40.9	1.6	37.8-44.0	41.7	1.5	38.8-44.6
<b>Education</b>															
< High School	57.4	2.9	51.7-62.9	52.2	2.8	46.7-57.7	53.7	2.9	48.0-59.3	51.5	3.0	45.8-57.3	58.8	3.1	52.7-64.6
High School	56.1	2.6	50.9-61.1	53.4	2.5	48.5-58.3	57.1	2.7	51.8-62.2	51.3	2.7	46.0-56.6	54.7	2.6	49.5-59.8
> High School	30.5	1.8	27.1-34.1	31.4	1.6	28.4-34.6	30.2	1.7	27.0-33.7	32.8	1.7	29.5-36.3	31.5	1.6	28.4-34.7
<b>Marital Status</b>															
Married	29.5	1.6	26.5-32.7	28.1	1.4	25.5-30.9	25.8	1.5	22.9-28.8	26.9	1.5	24.2-29.9	27.1	1.5	24.3-30.1
Other	68.0	2.1	63.7-72.0	65.3	2.0	61.3-69.2	67.6	2.0	63.5-71.5	66.6	2.1	62.3-70.7	67.1	2.0	63.0-70.9
<b>Payment for Delivery</b>															
Medicaid	61.0	2.0	57.1-64.8	58.3	1.8	54.7-61.8	58.1	1.9	54.4-61.7	57.3	1.9	53.5-61.0	59.2	1.8	55.6-62.7
Other	28.6	1.7	25.3-32.0	26.9	1.5	24.1-30.0	25.7	1.7	22.6-29.2	26.3	1.6	23.2-29.7	25.4	1.6	22.3-28.7
<b>Birthweight</b>															
Low	46.1	2.4	41.4-50.9	44.8	2.2	40.6-49.2	48.4	2.4	43.7-53.1	43.3	2.3	38.8-47.9	46.3	2.5	41.5-51.1
Normal	43.6	1.5	40.8-46.5	41.0	1.3	38.5-43.7	41.7	1.4	38.9-44.5	41.6	1.4	38.8-44.4	42.7	1.4	40.0-45.4

\*Confidence interval

**Table 4c**  
Unintended Pregnancy, 1998-2002  
By Selected Sociodemographic Characteristics

	1998			1999			2000			2001			2002		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	40.0	1.2	37.7-42.4	44.5	1.2	42.2-47.0	42.9	1.2	40.6-45.3	46.2	1.2	43.9-48.6	43.0	1.2	40.7-45.4
<b>Age</b>															
< 20 years	73.7	2.9	67.7-79.0	84.4	2.6	78.5-88.8	75.0	3.2	68.3-80.7	77.7	2.9	71.6-82.9	73.7	3.2	67.0-79.5
20-24 years	51.7	2.7	46.5-56.9	58.6	2.5	53.6-63.4	60.2	2.5	55.1-65.0	60.9	2.5	55.9-65.7	61.8	2.5	56.8-66.6
25-34 years	29.0	1.6	26.0-32.2	31.7	1.6	28.7-34.8	32.7	1.6	29.7-35.8	36.0	1.6	32.9-39.3	32.2	1.6	29.1-35.4
35+ years	26.3	3.2	20.6-32.9	28.4	3.1	22.6-34.9	26.3	2.8	21.2-32.2	33.1	2.9	27.6-39.1	27.9	2.9	22.6-33.8
<b>Race</b>															
Black	69.7	2.6	64.4-74.5	76.8	2.4	71.7-81.1	71.5	2.6	66.2-76.3	73.9	2.5	68.6-78.6	70.7	2.7	65.1-75.7
White	32.9	1.4	30.3-35.6	36.6	1.4	33.9-39.3	36.1	1.3	33.6-38.7	39.8	1.3	37.3-42.4	37.4	1.3	34.8-40.0
Other	29.7	8.2	16.4-47.6	34.5	5.3	24.9-45.6	41.7	6.6	29.7-54.9	35.9	5.6	25.8-47.4	35.6	6.0	24.9-47.9
<b>Hispanic Ethnicity</b>															
Hispanic	35.9	2.7	30.8-41.3	43.3	2.8	37.9-48.9	41.7	2.6	36.7-46.7	44.5	2.5	39.6-49.5	48.1	2.5	43.2-53.1
Non-Hispanic	41.0	1.3	38.4-43.6	44.9	1.4	42.2-47.6	43.3	1.4	40.7-46.0	46.8	1.4	44.1-49.5	41.5	1.4	38.8-44.3
<b>Education</b>															
< High School	54.3	2.8	48.8-59.6	60.6	2.7	55.2-65.8	55.1	2.7	49.7-60.3	62.0	2.6	56.8-66.8	59.2	2.7	53.9-64.3
High School	49.7	2.3	45.2-54.1	53.7	2.3	49.1-58.2	55.3	2.2	51.0-59.6	52.6	2.2	48.2-56.9	54.9	2.4	50.2-59.6
> High School	27.6	1.6	24.6-30.8	31.5	1.6	28.4-34.7	29.9	1.6	27.0-33.1	35.3	1.6	32.1-38.6	30.3	1.6	27.3-33.4
<b>Marital Status</b>															
Married	24.7	1.3	22.2-27.5	28.0	1.3	25.4-30.7	29.1	1.3	26.6-31.8	31.7	1.4	29.0-34.5	27.8	1.4	25.2-30.6
Other	70.5	2.0	66.6-74.2	73.3	1.9	69.4-76.8	68.8	2.0	64.7-72.5	71.0	1.8	67.3-74.4	69.5	1.9	65.7-73.1
<b>Payment for Delivery</b>															
Medicaid	59.4	2.1	55.3-63.3	67.6	2.0	63.6-71.3	63.3	1.9	59.4-67.0	64.7	1.8	61.1-68.2	61.1	1.8	57.4-64.6
Other	29.7	1.4	26.9-32.5	31.1	1.4	28.4-33.9	30.2	1.4	27.6-33.0	33.2	1.5	30.4-36.1	29.3	1.5	26.4-32.2
<b>Birthweight</b>															
Low	48.2	2.4	43.5-52.8	53.3	2.9	47.6-58.8	47.5	2.4	42.9-52.1	48.2	2.3	43.8-52.7	46.2	2.3	41.6-50.8
Normal	39.4	1.3	37.0-41.9	43.9	1.5	40.9-46.8	42.6	1.3	40.1-45.1	46.1	1.3	43.6-48.6	42.8	1.3	40.3-45.4

\* Confidence interval

**Table 5**  
 Use of Birth Control at Conception  
 Among Women Not Trying to Get Pregnant, 2009

	Percent	Standard Error	95% CI *
Using	49.8	2.0	45.9-53.6
Not using	50.3	2.0	46.4-54.1

\*Confidence interval

**Table 6**  
 Reasons for Not Using Birth Control at Conception  
 Among Women Not Trying to Get Pregnant, 2009

	Percent	Standard Error	95% CI *
Thought she could not get pregnant	20.8	2.2	16.8-25.4
Did not mind if she got pregnant	40.3	2.6	35.3-45.6
Other reasons	17.5	2.1	13.8-21.9
Had side effects from birth control measures	11.9	1.8	8.8-15.9
Husband or partner did not want to use any birth control	12.6	1.8	9.5-16.7
Had problems getting birth control when she needed	9.5	1.7	6.7-13.2
Thought she or her husband or partner was sterile	4.8	1.1	3.0-7.4

\* Confidence interval

**Table 7**

Trimester When Prenatal Care Was Started, 2009

	Percent	Standard Error	95% CI *
1st trimester	82.3	1.1	80.1-84.3
2nd trimester	15.7	1.0	13.8-17.8
3rd trimester	1.1	0.3	0.7-1.9
No prenatal care	0.9	0.3	0.5-1.6

\* Confidence interval

**Table 8**  
Initiation of Prenatal Care in the First Trimester, 2009  
By Selected Sociodemographic Characteristics

	Percent	Standard Error	95% CI *
<b>Total</b>	82.3	1.1	80.1-84.3
<b>Age</b>			
< 20 years	65.0	4.2	56.4-72.7
20-24 years	73.8	2.8	68.0-78.9
25-34 years	86.8	1.3	84.1-89.1
35+ years	89.0	2.2	83.9-92.6
<b>Race</b>			
Black	68.4	3.5	61.1-74.9
White	84.7	1.1	82.4-86.7
Other	88.1	3.8	78.6-93.7
<b>Hispanic Ethnicity</b>			
Hispanic	74.7	2.4	69.8-79.0
Non-Hispanic	84.9	1.2	82.4-87.1
<b>Education</b>			
< High School	69.0	3.0	62.9-74.5
High School	77.6	2.3	72.7-81.8
> High School	89.1	1.2	86.6-91.2
<b>Marital Status</b>			
Married	90.2	1.1	87.9-92.1
Other	70.2	2.1	65.9-74.1
<b>Payment for Delivery</b>			
Medicaid	73.1	1.7	69.6-76.3
Other	93.1	1.0	90.9-94.9
<b>Birthweight</b>			
Low	80.5	2.2	75.8-84.5
Normal	82.4	1.1	80.1-84.6

\* Confidence interval

**Table 9**  
 Problems Getting Prenatal Care  
 Among women who didn't care as early in pregnancy as  
 they wanted, 2009

	Percent	Standard Error	95% CI *
I couldn't get an appointment when I wanted one	38.9	3.4	32.5-45.6
I didn't have my Medicaid card or All Kids, Moms and Babies Card	30.3	3.2	24.3-37.0
I didn't have enough money or insurance to pay for my visits	27.6	3.1	22.0-34.1
I had no transportation to get to the clinic or doctor's office	15.8	2.6	11.3-21.6
I couldn't take time off from work or school	10.0	2.1	6.5-15.0
The doctor or my health plan would not start as early as I wanted	22.9	2.9	17.7-29.2
I had no one to take care of my children	11.1	2.3	7.4-16.4
Other			
I didn't want anyone else to know I was pregnant	15.6	2.6	11.1-21.4
I had too many other things going on	22.9	3.0	17.5-29.2
I didn't know that I was pregnant	38.3	3.4	31.9-45.2
I didn't want prenatal care	4.6	1.6	2.3-9.0

\*Confidence interval

**Table 10**  
 Topics Discussed by Health Care Professionals  
 During Prenatal Care Visits, 2009

	Percent	Standard Error	95% CI *
Genetic screening	89.8	0.8	88.0-91.3
Safe medications during pregnancy	88.6	0.9	86.8-90.2
What to do if labor starts early	83.5	1.0	81.4-85.4
Asked if mom is drinking alcohol	82.2	1.0	80.0-84.1
Getting tested for HIV (the virus that causes AIDS)	75.6	1.2	73.3-77.9
Breastfeeding the baby	84.4	1.0	82.4-86.2
What to do if I feel depressed during my pregnancy or after the baby is born	74.7	1.2	72.2-76.9
Signs and symptoms of preterm labor	81.4	1.1	79.2-83.4
How drinking alcohol during pregnancy could affect my baby	75.8	1.2	73.4-78.0
How smoking during pregnancy could affect my baby	74.9	1.2	72.5-77.1
How using illegal drugs could affect my baby	67.9	1.3	65.4-70.4
Use of seat belt during pregnancy	56.9	1.4	54.2-59.5
Physical abuse to women by their husbands or partners	54.1	1.4	51.4-56.8

\*Confidence interval

**Table 11**Discussions about Nutrition and Weight Management  
During and After Pregnancy, 2009

	Percent	Standard Error	95% CI *
Foods that are good to eat during pregnancy	82.2	1.0	80.1-84.1
How much weight to gain during pregnancy	81.7	1.1	79.6-83.7
Exercise during pregnancy	78.0	1.1	75.7-80.1
Programs or resources to help me gain the right amount of weight during pregnancy	35.0	1.3	32.5-37.6
Programs or resources to help me lose weight after pregnancy	22.3	1.2	20.1-24.7

**Table 12**  
 Knowledge About Folic Acid, 2009  
 By Selected Sociodemographic Characteristics

	Percent	Standard Error	95% CI *
<b>Total</b>	81.8	1.1	79.6-83.9
<b>Age</b>			
< 20 years	63.6	4.3	54.9-71.4
20-24 years	68.2	2.9	62.2-73.6
25-34 years	87.3	1.2	84.7-89.5
35+ years	92.2	1.9	87.6-95.2
<b>Race</b>			
Black	68.2	3.5	61.0-74.7
White	84.7	1.1	82.4-86.8
Other	80.0	4.5	69.7-87.4
<b>Hispanic Ethnicity</b>			
Hispanic	82.0	2.1	77.5-85.7
Non-Hispanic	81.8	1.3	79.2-84.1
<b>Education</b>			
< High School	72.2	2.9	66.2-77.4
High School	73.3	2.5	68.2-77.8
> High School	89.3	1.2	86.8-91.3
<b>Marital Status</b>			
Married	89.5	1.1	87.2-91.4
Other	70.1	2.1	65.9-74.0
<b>Payment for Delivery</b>			
Medicaid	75.1	1.7	71.8-78.2
Other	89.7	1.2	87.0-91.8
<b>Birthweight</b>			
Low	80.3	2.2	75.6-84.3
Normal	82.0	1.1	79.6-84.1

\*Confidence interval

**Table 13**  
 Oral Health Care During Pregnancy by Selected Sociodemographic Characteristics, 2009

	Needed to See a Dentist for a Problem			Dental/Health Care Worker Spoke About Care of Gums and Teeth			Visited Dentist or Dental Clinic		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	21.8	1.2	19.6-24.1	47.2	1.4	44.5-49.9	44.0	1.4	41.3-46.7
<b>Age</b>									
< 20 years	17.4	3.3	11.8-24.9	51.4	4.5	42.6-60.0	32.9	4.2	25.3-41.5
20-24 years	21.9	2.6	17.3-27.4	34.4	3.0	28.9-40.4	28.6	2.8	23.3-34.4
25-34 years	21.6	1.6	18.7-24.8	49.4	1.9	45.7-53.1	48.0	1.9	44.4-51.7
35+ years	25.1	3.0	19.7-31.4	53.7	3.4	47.0-60.4	57.5	3.4	50.7-64.0
<b>Race</b>									
Black	24.4	3.2	18.6-31.3	37.2	3.7	30.3-44.7	28.1	3.5	21.9-35.4
White	21.8	1.3	19.4-24.4	50.0	1.5	47.0-53.0	48.3	1.5	45.3-51.4
Other	16.1	4.1	9.5-25.8	35.7	5.4	25.9-46.9	27.9	5.1	19.1-38.9
<b>Hispanic Ethnicity</b>									
Hispanic	21.1	2.3	17.0-25.9	43.0	2.7	37.8-48.4	38.0	2.7	32.9-43.4
Non-Hispanic	22.1	1.3	19.6-24.8	48.7	1.6	45.5-51.8	46.0	1.6	42.9-49.1
<b>Education</b>									
< High School	21.2	2.6	16.5-26.9	38.2	3.1	32.2-44.4	33.2	3.1	27.4-39.5
High School	27.4	2.5	22.8-32.7	40.1	2.8	34.8-45.6	32.4	2.7	27.5-37.8
> High School	19.5	1.5	16.8-22.5	54.2	1.8	50.6-57.8	53.2	1.8	49.6-56.8
<b>Marital Status</b>									
Married	20.8	1.4	18.2-23.7	52.2	1.7	48.8-55.6	52.8	1.7	49.3-56.2
Other	23.4	1.9	19.8-27.4	39.6	2.2	35.3-44.0	30.5	2.1	26.6-34.8
<b>Payment for Delivery</b>									
Medicaid	26.3	1.7	23.1-29.7	39.1	1.9	35.5-42.8	32.1	1.8	28.7-35.7
Other	16.6	1.5	13.9-19.7	56.9	2.0	53.0-60.8	57.9	2.0	54.0-61.7
<b>Birthweight</b>									
Low	22.3	2.3	18.2-27.1	40.5	2.7	35.4-45.8	35.1	2.6	30.3-40.2
Normal	21.8	1.2	19.5-24.3	47.8	1.5	44.9-50.7	44.7	1.5	41.9-47.6

\*Confidence interval

**Table 14**  
WIC Participation, 2009  
By Selected Sociodemographic Characteristics

	Participation During Pregnancy			Participation After Delivery		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	46.8	1.4	44.2-49.5	54.6	1.4	51.9-57.3
<b>Age</b>						
< 20 years	85.4	2.9	78.6-90.2	90.3	2.6	83.8-94.4
20-24 years	72.2	2.8	66.4-77.4	85.0	2.2	80.0-88.9
25-34 years	36.9	1.8	33.4-40.5	43.7	1.9	40.1-47.3
35+ years	23.7	2.9	18.4-29.9	31.1	3.3	25.1-37.9
<b>Race</b>						
Black	70.2	3.4	63.1-76.5	87.5	2.6	81.5-91.7
White	43.2	1.5	40.3-46.1	49.4	1.5	46.4-52.4
Other	32.3	5.2	23.0-43.3	38.8	5.5	28.7-49.9
<b>Hispanic Ethnicity</b>						
Hispanic	71.9	2.4	66.9-76.4	84.7	2.0	80.4-88.1
Non-Hispanic	38.3	1.6	35.3-41.4	44.5	1.6	41.4-47.6
<b>Education</b>						
< High School	79.2	2.6	73.7-83.8	94.1	1.5	90.3-96.4
High School	69.7	2.6	64.5-74.5	77.7	2.4	72.8-82.0
> High School	24.5	1.6	21.5-27.7	30.2	1.7	27.0-33.6
<b>Marital Status</b>						
Married	28.5	1.6	25.5-31.7	33.8	1.7	30.6-37.1
Other	75.1	1.9	71.1-78.7	87.7	1.5	84.5-90.3
<b>Payment for Delivery</b>						
Medicaid	78.3	1.6	75.1-81.3	90.5	1.1	88.0-92.5
Other	9.4	1.2	7.3-12.1	13.3	1.4	10.8-16.3
<b>Birthweight</b>						
Low	53.1	2.6	47.9-58.2	64.2	2.6	59.0-69.1
Normal	46.3	1.5	43.5-49.2	53.9	1.5	51.1-56.8

\*Confidence interval

**Table 15a**  
HIV Testing During Pregnancy, 2009  
By Selected Sociodemographic Characteristics

	Tested			Not Tested			Don't Know If Tested		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	69.5	1.3	67.0-71.9	17.0	1.0	15.1-19.1	13.5	0.9	11.8-15.5
<b>Age</b>									
< 20 years	74.1	3.9	65.8-81.0	11.0	2.7	6.7-17.4	15.0	3.2	9.7-22.4
20-24 years	69.9	2.8	64.1-75.1	15.3	2.2	11.5-20.0	14.9	2.2	11.0-19.7
25-34 years	70.3	1.7	66.9-73.4	16.8	1.4	14.3-19.7	12.9	1.2	10.7-15.6
35+ years	63.2	3.3	56.5-69.4	24.0	2.9	18.7-30.2	12.9	2.3	9.0-18.0
<b>Race</b>									
Black	86.1	2.6	80.1-90.5	4.8	1.5	2.6-8.9	9.1	2.3	5.5-14.6
White	67.0	1.4	64.1-69.7	18.6	1.2	16.4-21.1	14.4	1.1	12.4-16.6
Other	58.4	5.5	47.4-68.7	26.7	4.9	18.1-37.3	14.9	4.0	8.6-24.6
<b>Hispanic Ethnicity</b>									
Hispanic	71.3	2.5	66.3-75.9	16.4	2.0	12.9-20.8	12.3	1.8	9.2-16.2
Non-Hispanic	68.8	1.5	65.9-71.6	17.2	1.2	15.0-19.7	14.0	1.1	11.9-16.3
<b>Education</b>									
< High School	73.6	2.8	67.7-78.7	14.4	2.2	10.6-19.3	12.0	2.1	8.4-16.8
High School	71.6	2.5	66.5-76.2	13.0	1.8	9.8-17.1	15.4	2.0	11.9-19.8
> High School	67.3	1.7	63.8-70.5	19.5	1.4	16.9-22.5	13.2	1.2	11.0-15.8
<b>Marital Status</b>									
Married	65.8	1.6	62.6-69.0	19.8	1.4	17.3-22.7	14.3	1.2	12.1-16.9
Other	75.0	1.9	71.0-78.6	12.7	1.5	10.1-15.9	12.3	1.5	9.7-15.5
<b>Payment for Delivery</b>									
Medicaid	72.7	1.7	69.3-75.9	13.5	1.3	11.2-16.2	13.7	1.3	11.4-16.5
Other	65.4	1.9	61.6-69.0	21.3	1.6	18.3-24.6	13.4	1.3	10.9-16.2
<b>Birthweight</b>									
Low	72.5	2.4	67.6-76.9	14.4	1.9	11.1-18.4	13.1	1.8	10.1-17.0
Normal	69.2	1.3	66.5-71.8	17.2	1.1	15.2-19.5	13.6	1.0	11.7-15.6

\*Confidence interval

**Table 15b**  
 Prevalence of HIV Testing During Pregnancy, 2008  
 By Selected Sociodemographic Characteristics

	Tested			Not Tested			Don't Know If Tested		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	76.0	1.1	73.7-78.1	16.0	1.0	14.2-18.0	8.0	0.7	6.7-9.6
<b>Age</b>									
< 20 years	79.8	3.2	72.9-85.3	13.0	2.6	8.7-19.0	7.2	2.1	4.0-12.5
20-24 years	79.9	2.3	75.0-84.1	13.8	2.0	10.3-18.2	6.3	1.4	4.0-9.8
25-34 years	73.9	1.6	70.7-76.9	17.0	1.3	14.6-19.8	9.1	1.0	7.2-11.3
35+ years	75.0	2.9	68.8-80.3	17.6	2.6	13.1-23.2	7.4	1.8	4.6-11.8
<b>Race</b>									
Black	89.8	2.2	84.5-93.4	6.0	1.7	3.4-10.3	4.3	1.5	2.1-8.5
White	73.1	1.3	70.5-75.5	18.0	1.1	15.9-20.3	8.9	0.8	7.4-10.7
Other	72.5	5.2	61.2-81.5	20.3	4.7	12.6-31.1	7.2	3.1	3.0-16.0
<b>Hispanic Ethnicity</b>									
Hispanic	75.3	2.2	70.8-79.3	17.8	1.9	14.4-21.8	6.9	1.3	4.8-9.9
Non-Hispanic	76.2	1.3	73.5-78.7	15.3	1.1	13.3-17.6	8.5	0.9	6.9-10.3
<b>Education</b>									
< High School	80.7	2.3	75.8-84.8	13.6	2.0	10.2-17.9	5.7	1.4	3.6-9.0
High School	78.4	2.2	73.8-82.3	13.6	1.8	10.5-17.5	8.1	1.4	5.7-11.3
> High School	72.8	1.6	69.6-75.8	18.3	1.4	15.7-21.1	8.9	1.0	7.1-11.2
<b>Marital Status</b>									
Married	72.2	1.5	69.2-75.0	18.7	1.3	16.3-21.4	9.1	1.0	7.4-11.2
Other	81.4	1.6	77.9-84.4	12.1	1.4	9.7-15.1	6.5	1.1	4.7-8.9
<b>Payment for Delivery</b>									
Medicaid	80.7	1.5	77.7-83.4	12.6	1.2	10.4-15.1	6.7	0.9	5.1-8.8
Other	70.9	1.7	67.5-74.1	19.7	1.5	16.9-22.7	9.4	1.1	7.5-11.8
<b>Birthweight</b>									
Low	77.7	2.1	73.4-81.5	17.0	1.9	13.6-20.9	5.4	1.1	3.6-7.9
Normal	75.8	1.2	73.4-78.1	15.9	1.0	14.0-18.0	8.2	0.8	6.9-9.9

\*Confidence interval

**Table 15c**

Prevalence of HIV Testing During Pregnancy, 2003-2007

By Selected Sociodemographic Characteristics

	2003			2004**			2005			2006			2007		
	Percent	Standard Error	95% CI*												
<b>Total</b>	66.9	1.3	64.4-69.4	70.0	1.1	67.7-72.2	80.5	1.1	78.4-82.5	77.2	1.1	74.9-79.3	78.4	1.1	76.3-80.4
<b>Age</b>															
< 20 years	75.1	3.7	67.1-81.7	77.5	3.6	69.6-83.8	87.0	2.8	80.5-91.5	82.4	3.3	75.0-88.0	85.6	2.9	79.0-90.3
20-24 years	73.5	2.4	68.5-77.9	75.4	2.3	70.5-79.7	84.7	2.1	80.1-88.3	81.6	2.3	76.7-85.6	83.2	2.1	78.7-87.0
25-34 years	64.0	1.8	60.3-67.5	70.2	1.5	67.2-73.1	77.6	1.5	74.4-80.5	75.7	1.5	72.5-78.6	75.6	1.5	72.6-78.4
35+ years	59.7	3.7	52.3-66.7	57.2	3.1	50.9-63.1	80.4	2.7	74.5-85.2	72.9	3.0	66.6-78.4	76.5	2.9	70.4-81.7
<b>Race</b>															
Black	82.1	2.6	76.4-86.7	84.0	2.3	78.9-88.1	89.7	2.0	85.0-93.0	90.6	2.0	85.8-93.9	91.9	1.8	87.6-94.8
White	63.9	1.5	61.0-66.8	66.8	1.3	64.2-69.3	78.7	1.2	76.1-81.0	74.9	1.3	72.3-77.4	75.2	1.3	72.6-77.6
Other	61.4	6.3	48.6-72.8	70.7	5.3	59.4-79.8	75.6	5.6	63.1-84.8	66.1	5.4	54.8-75.7	76.7	4.2	67.5-84.0
<b>Hispanic Ethnicity</b>															
Hispanic	71.4	2.5	66.4-76.0	76.7	2.2	72.2-80.7	85.2	1.9	81.1-88.6	75.3	2.3	70.6-79.4	79.0	2.2	74.3-83.0
Non-Hispanic	65.4	1.5	62.4-68.3	68.0	1.3	65.3-70.5	79.1	1.3	76.5-81.4	77.9	1.3	75.2-80.3	78.3	1.2	75.8-80.5
<b>Education</b>															
< High School	75.7	2.5	70.6-80.2	78.7	2.3	73.9-82.8	83.9	2.1	79.4-87.7	74.7	2.6	69.2-79.5	82.7	2.3	77.7-86.8
High School	71.1	2.4	66.2-75.5	73.4	2.2	68.9-77.5	80.5	2.1	76.0-84.3	79.2	2.2	74.6-83.1	80.5	2.1	76.1-84.2
> High School	60.6	1.9	56.8-64.2	65.1	1.6	61.9-68.2	79.0	1.5	75.9-81.8	77.3	1.5	74.2-80.1	75.7	1.5	72.7-78.5
<b>Marital Status</b>															
Married	61.2	1.7	57.9-64.4	63.9	1.5	60.9-66.7	76.9	1.5	73.9-79.6	73.8	1.5	70.9-76.6	74.6	1.4	71.8-77.3
Other	76.6	1.9	72.7-80.2	81.1	1.7	77.7-84.2	86.1	1.5	83.0-88.8	82.8	1.7	79.3-85.9	84.1	1.5	80.9-86.9
<b>Payment for Delivery</b>															
Medicaid	74.6	1.7	71.0-77.8	77.6	1.5	74.5-80.5	83.2	1.4	80.2-85.8	80.7	1.5	77.5-83.5	81.2	1.4	78.2-83.9
Other	60.1	1.8	56.4-63.6	63.5	1.6	60.3-66.6	77.8	1.6	74.5-80.8	73.6	1.6	70.3-76.7	75.4	1.6	72.3-78.4
<b>Birthweight</b>															
Low	68.4	2.3	63.9-72.6	72.7	1.9	68.8-76.3	81.3	1.9	77.3-84.7	79.2	1.9	75.2-82.7	82.9	1.8	79.0-86.1
Normal	66.8	1.4	64.1-69.4	69.8	1.2	67.3-72.1	80.5	1.1	78.2-82.6	77.0	1.2	74.6-79.3	78.1	1.1	75.8-80.2

\*Confidence interval \*\* Question changed in 2004

**Table15d**  
Prevalence of HIV Testing During Pregnancy, 1998-2002  
By Selected Sociodemographic Characteristics

	1998			1999			2000			2001			2002		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	64.4	1.2	62.1-66.6	72.4	1.1	70.3-74.4	70.7	1.1	68.5-72.8	71.4	1.1	69.2-73.4	73.1	1.1	70.9-75.2
<b>Age</b>															
< 20 years	78.6	2.7	73.0-83.4	79.5	2.8	73.4-84.5	81.0	2.8	74.9-85.9	80.6	2.7	74.7-85.3	76.9	3.1	70.3-82.4
20-24 years	69.1	2.4	64.2-73.6	78.4	2.0	74.2-82.0	74.3	2.3	69.6-78.4	76.3	2.2	71.8-80.3	81.0	2.0	76.7-84.6
25-34 years	60.4	1.7	57.1-63.7	68.7	1.5	65.7-71.7	67.9	1.5	64.8-70.8	68.9	1.5	65.8-71.9	72.3	1.5	69.2-75.2
35+ years	56.7	3.4	50.0-63.2	68.3	3.0	62.1-73.9	66.6	2.9	60.6-72.1	64.7	2.9	58.8-70.2	60.4	3.1	54.1-66.3
<b>Race</b>															
Black	76.8	2.3	72.0-81.0	84.7	2.0	80.3-88.2	80.9	2.2	76.2-84.8	87.8	1.8	83.7-91.0	87.9	2.0	83.5-91.3
White	61.5	1.4	58.8-64.1	69.2	1.3	66.7-71.6	68.2	1.3	65.6-70.6	67.3	1.3	64.8-69.8	69.9	1.3	67.4-72.4
Other	56.1	8.0	40.3-70.7	72.4	4.9	61.9-80.8	71.2	6.1	58.1-81.5	69.7	5.2	58.5-78.9	69.8	5.6	57.8-79.6
<b>Hispanic Ethnicity</b>															
Hispanic	65.1	2.6	59.8-70.0	78.3	2.3	73.5-82.4	73.3	2.3	68.6-77.5	76.6	2.2	72.0-80.5	79.1	2.0	74.8-82.8
Non-Hispanic	64.2	1.3	61.6-66.7	70.9	1.2	68.5-73.2	69.9	1.2	67.4-72.3	69.8	1.2	67.4-72.2	71.1	1.3	68.5-73.5
<b>Education</b>															
< High School	72.9	2.4	67.9-77.3	78.1	2.2	73.6-82.0	78.3	2.2	73.6-82.3	76.4	2.2	71.8-80.5	80.6	2.1	76.1-84.4
High School	65.0	2.1	60.8-69.0	74.3	1.9	70.3-77.9	72.6	2.0	68.5-76.2	73.9	1.9	69.9-77.5	75.0	2.1	70.6-78.9
> High School	60.4	1.7	57.0-63.6	68.6	1.6	65.5-71.6	66.0	1.6	62.8-69.0	67.2	1.6	64.0-70.3	69.4	1.6	66.2-72.4
<b>Marital Status</b>															
Married	58.1	1.5	55.2-61.0	67.1	1.4	64.4-69.7	64.7	1.4	62.0-67.4	66.2	1.4	63.4-68.9	68.3	1.4	65.5-71.1
Other	76.9	1.8	73.2-80.2	81.5	1.6	78.1-84.4	81.6	1.6	78.2-84.6	80.2	1.6	76.9-83.1	81.3	1.6	77.9-84.3
<b>Payment for Delivery</b>															
Medicaid	72.4	1.9	68.6-75.9	79.1	1.6	75.7-82.1	80.2	1.6	76.9-83.1	78.8	1.5	75.6-81.7	79.6	1.5	76.4-82.4
Other	60.3	1.5	57.4-63.2	68.5	1.4	65.8-71.2	64.9	1.4	62.1-67.6	66.1	1.5	63.2-68.9	68.1	1.5	65.1-71.0
<b>Birthweight</b>															
Low	71.8	2.1	67.5-75.8	72.7	2.5	67.5-77.3	73.4	2.1	69.2-77.3	73.0	2.0	68.9-76.8	77.7	1.9	73.7-81.2
Normal	63.8	1.2	61.4-66.2	72.9	1.3	70.3-75.3	70.5	1.2	68.1-72.7	71.2	1.2	68.9-73.4	72.7	1.2	70.4-75.0

\*Confidence interval

**Table 16**  
Problems During Pregnancy, 2009

	Percent	Standard Error	95% CI *
Severe nausea, vomiting or dehydration	24.6	1.2	22.4-27.0
Kidney or bladder (urinary tract) infection	20.4	1.1	18.3-22.7
Labor pains more than three weeks before my baby was due (preterm or early labor)	17.7	1.0	15.8-19.7
Vaginal bleeding	18.3	1.1	16.3-20.5
High blood pressure, hypertension (including pregnancy-induced hypertension (PIH), preeclampsia, or toxemia)	12.9	0.9	11.2-14.7
Problems with placenta	4.7	0.5	3.8-5.9
Water broke more than three weeks before my baby was due (premature rupture of membranes [PROM])	5.3	0.5	4.4-6.4
Hurt in car accident	0.8	0.3	0.4-1.5
Had a blood transfusion	2.0	0.4	1.4-2.9
Cervix sewn shut	1.3	0.3	0.9-2.1

\*Confidence interval

**Table 17**  
 Number of Stressors During the Year Prior to Delivery, 2009  
 By Selected Sociodemographic Characteristics

	No Stressors			1-2 Stressors			3-5 Stressors			More than 5 Stressors		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	29.3	1.2	27.0-31.8	41.3	1.3	38.7-43.9	23.7	1.2	21.5-26.1	5.7	0.7	4.6-7.2
<b>Age</b>												
< 20 years	18.2	3.5	12.4-26.0	43.5	4.4	35.3-52.2	27.2	3.9	20.3-35.4	11.1	2.8	6.6-17.9
20-24 years	15.8	2.2	11.8-20.7	37.5	3.0	31.8-43.6	37.7	3.1	31.9-43.8	9.1	1.8	6.2-13.2
25-34 years	33.3	1.7	30.0-36.7	43.0	1.8	39.5-46.6	19.1	1.5	16.4-22.1	4.6	0.8	3.3-6.5
35+ years	40.0	3.3	33.6-46.6	38.4	3.3	32.2-45.1	19.8	2.8	15.0-25.8	--	--	--
<b>Race</b>												
Black	21.0	3.0	15.6-27.6	30.6	3.4	24.4-37.7	35.7	3.6	28.9-43.1	12.7	2.5	8.5-18.5
White	30.3	1.4	27.6-33.1	43.5	1.5	40.6-46.5	21.8	1.3	19.4-24.3	4.5	0.6	3.4-5.9
Other	38.9	5.4	29.0-49.8	42.3	5.5	32.1-53.3	15.5	4.0	9.2-25.0	--	--	--
<b>Hispanic Ethnicity</b>												
Hispanic	23.8	2.3	19.6-28.6	45.6	2.7	40.4-50.9	25.9	2.4	21.5-30.8	4.7	1.2	2.9-7.6
Non-Hispanic	31.2	1.5	28.4-34.1	39.8	1.5	36.8-42.9	23.0	1.4	20.4-25.8	6.1	0.8	4.7-7.8
<b>Education</b>												
< High School	22.7	2.6	17.9-28.3	41.2	3.1	35.2-47.5	27.5	2.9	22.2-33.4	8.7	1.8	5.7-12.9
High School	21.5	2.3	17.3-26.4	39.0	2.7	33.8-44.4	31.3	2.6	26.4-36.6	8.2	1.6	5.7-11.9
> High School	35.2	1.7	31.9-38.6	43.0	1.8	39.5-46.5	18.8	1.5	16.1-21.8	3.1	0.6	2.0-4.6
<b>Marital Status</b>												
Married	36.9	1.7	33.7-40.2	44.8	1.7	41.5-48.2	16.3	1.3	13.9-19.0	2.0	0.5	1.3-3.3
Other	17.6	1.7	14.5-21.2	35.7	2.2	31.6-40.0	35.3	2.2	31.2-39.6	11.4	1.5	8.9-14.6
<b>Payment for Delivery</b>												
Medicaid	20.3	1.5	17.5-23.4	39.2	1.9	35.6-42.9	32.0	1.8	28.6-35.6	8.6	1.1	6.7-10.9
Other	40.1	1.9	36.3-43.9	43.7	2.0	39.9-47.6	13.9	1.4	11.4-16.8	2.4	0.7	1.3-4.1
<b>Birthweight</b>												
Low	27.3	2.3	23.0-32.1	38.9	2.6	34.0-44.0	25.5	2.4	21.1-30.3	8.3	1.5	5.8-11.9
Normal	29.5	1.3	27.0-32.1	41.4	1.4	38.7-44.3	23.6	1.3	21.2-26.1	5.5	0.7	4.3-7.1

\*Confidence interval  
 -- Fewer than six respondents

**Table 18**

Types of Stressors Encountered During the Year Prior to Delivery, 2009

	Percent	Standard Error	95% CI *
Moved to a new address	27.8	1.2	25.4-30.3
Arguments with husband or partner more than usual	25.3	1.2	23.0-27.7
Family member was sick and had to go to the hospital	23.3	1.2	21.1-25.7
Lots of bills that couldn't be paid	23.5	1.2	21.3-26.0
Someone very close died	15.3	1.0	13.5-17.4
Husband or partner lost job	16.0	1.0	14.1-18.1
Lost job	13.6	1.0	11.8-15.5
Separated or divorced from husband or partner	10.7	0.9	9.1-12.5
Someone close had problem with drinking or drugs	10.2	0.9	8.7-12.0
Husband or partner did not want pregnancy	7.6	0.8	6.2-9.2
Husband or partner or I went to jail	3.9	0.6	3.0-5.2
Physical fight	3.0	0.5	2.2-4.2
Homeless	8.4	0.8	7.0-10.0

\*Confidence interval

**Table 19**  
Physical Abuse by Husband or Partner Before and During Pregnancy, 2009  
By Selected Sociodemographic Characteristics

	12 Months Before Pregnancy			During Most Recent Pregnancy		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	2.8	0.5	2.0-3.9	2.1	0.4	1.4-3.2
<b>Age</b>						
< 20 years***	8.7	2.8	4.6-16.0	8.9	2.9	4.6-16.5
20-24 years	5.6	1.6	3.3-9.6	3.3	1.2	1.6-6.8
25-34 years	1.7	0.5	1.0-2.9	1.3	0.4	0.7-2.6
35+ years	--	--	--	--	--	--
<b>Race</b>						
Black	5.1	1.8	2.5-10.2	4.0	0.7	1.7-9.0
White	2.5	0.5	1.7-3.6	1.9	0.4	1.2-3.0
Other	--	--	--	--	--	--
<b>Hispanic Ethnicity</b>						
Hispanic	3.7	1.0	2.1-6.4	3.2	1.0	1.8-5.8
Non-Hispanic	2.5	0.5	1.7-3.8	1.8	0.5	1.1-2.9
<b>Education</b>						
< High School	5.2	1.6	2.9-9.3	6.3	1.8	3.6-10.7
High School	4.0	1.1	2.3-6.7	2.0	0.8	0.9-4.2
> High School	1.3	0.4	0.7-2.4	0.6	0.3	0.3-1.5
<b>Marital Status</b>						
Married	1.0	0.3	0.5-1.9	0.7	0.3	0.3-1.5
Other	5.8	1.1	4.0-8.4	4.6	1.0	3.0-7.1
<b>Payment for Delivery</b>						
Medicaid	4.6	0.8	3.2-6.5	3.6	0.8	2.4-5.4
Other	0.8	0.3	0.3-1.8	--	--	--
<b>Birthweight</b>						
Low	5.5	1.3	3.4-8.6	3.0	0.9	1.6-5.4
Normal	2.6	0.5	1.8-3.8	2.1	0.5	1.3-3.2

\*Confidence interval

\*\*\*Mothers younger than 18 years old were not asked abuse questions.

--Fewer than six respondents

**Table 20a**  
 Prevalence of Smoking Before, During and After Pregnancy, 2009  
 By Selected Sociodemographic Characteristics

	Three Months Before Pregnancy			Last Three Months Of Pregnancy			Now		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	20.9	1.1	18.8-23.2	9.2	0.8	7.7-10.8	14.6	1.0	12.8-16.5
<b>Age</b>									
< 20 years	27.0	3.9	20.1-35.2	10.8	2.7	6.5-17.3	20.6	3.5	14.5-28.3
20-24 years	30.1	2.9	24.8-36.1	12.4	2.0	8.9-16.9	24.2	2.7	19.4-29.8
25-34 years	20.0	1.5	17.3-23.1	9.6	1.1	7.7-12.0	13.1	1.2	10.8-15.7
35+ years	8.6	1.9	5.5-13.2	2.6	1.0	1.2-5.6	3.5	1.2	1.8-6.8
<b>Race</b>									
Black	19.4	3.0	14.2-25.9	8.2	2.0	5.0-13.1	15.2	2.7	10.6-21.2
White	22.3	1.3	19.9-24.9	9.8	0.9	8.2-11.7	15.1	1.1	13.1-17.3
Other	6.7	2.8	2.9-14.7	--	--	--	--	--	--
<b>Hispanic Ethnicity</b>									
Hispanic	9.0	1.6	6.4-12.6	2.0	0.8	1.0-4.2	4.4	1.1	2.7-7.1
Non-Hispanic	25.0	1.4	22.4-27.8	11.6	1.0	9.8-13.7	18.0	1.2	15.8-20.5
<b>Education</b>									
< High School	18.9	2.5	14.5-24.2	8.2	1.7	5.4-12.2	15.9	2.3	11.9-21.0
High School	31.1	2.6	26.3-36.4	18.1	2.1	14.3-22.6	25.5	2.4	21.0-30.6
> High School	17.0	1.4	14.5-19.8	5.7	0.8	4.3-7.5	9.1	1.0	7.3-11.3
<b>Marital Status</b>									
Married	15.3	1.2	13.0-17.9	5.8	0.8	4.5-7.6	8.6	1.0	6.9-10.7
Other	29.6	2.0	25.8-33.8	14.3	1.6	11.6-17.6	23.7	1.9	20.2-27.6
<b>Payment for Delivery</b>									
Medicaid	25.5	1.7	22.4-28.9	12.5	1.2	10.2-15.1	19.7	1.5	17.0-22.8
Other	15.6	1.4	13.0-18.6	5.3	0.9	3.9-7.3	8.5	1.1	6.6-10.9
<b>Birthweight</b>									
Low	26.1	2.4	21.8-31.1	13.1	1.9	9.9-17.2	20.6	2.2	16.6-25.2
Normal	20.5	1.2	18.3-22.9	8.9	0.8	7.4-10.6	14.1	1.0	12.2-16.2

\* Confidence interval

-- Fewer than six respondents

**Table 20b**  
 Prevalence of Smoking Before, During and After Pregnancy, 2008  
 By Selected Sociodemographic Characteristics

	Three Months Before Pregnancy			Last Three Months Of Pregnancy			Now		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	19.2	1.1	17.2-21.4	9.6	0.8	8.2-11.3	14.2	0.9	12.5-16.1
<b>Age</b>									
< 20 years	27.2	3.6	20.8-34.7	15.5	2.9	10.6-22.0	22.3	3.3	16.5-29.5
20-24 years	28.7	2.7	23.7-34.2	14.8	2.1	11.1-19.3	25.5	2.6	20.7-30.9
25-34 years	16.4	1.4	13.9-19.2	8.0	1.0	6.2-10.1	10.8	1.1	8.8-13.2
35+ years	10.1	2.0	6.8-14.8	3.9	1.3	2.1-7.3	4.4	1.3	2.4-7.9
<b>Race</b>									
Black	19.9	2.9	14.8-26.3	11.8	2.3	8.0-17.1	17.5	2.8	12.8-23.6
White	19.8	1.2	17.6-22.2	9.6	0.9	8.0-11.4	13.9	1.0	12.0-16.0
Other	--	--	--	--	--	--	--	--	--
<b>Hispanic Ethnicity</b>									
Hispanic	9.3	1.5	6.8-12.7	2.2	0.7	1.1-4.1	5.9	1.2	3.9-8.7
Non-Hispanic	22.8	1.3	20.3-25.4	12.3	1.0	10.4-14.5	17.2	1.2	15.0-19.6
<b>Education</b>									
< High School	22.2	2.5	17.7-27.5	15.1	2.2	11.3-19.8	19.1	2.4	14.9-24.2
High School	29.8	2.4	25.2-34.7	16.2	1.9	12.8-20.4	24.6	2.3	20.4-29.3
> High School	12.5	1.2	10.4-15.1	4.5	0.7	3.2-6.2	7.0	0.9	5.4-9.0
<b>Marital Status</b>									
Married	10.9	1.0	9.0-13.1	4.6	0.7	3.4-6.1	6.8	0.8	5.4-8.6
Other	31.2	2.0	27.4-35.3	16.9	1.6	14.0-20.2	24.8	1.9	21.4-28.6
<b>Payment for Delivery</b>									
Medicaid	25.2	1.6	22.1-28.6	14.5	1.3	12.1-17.3	20.8	1.5	18.0-24.0
Other	12.9	1.3	10.6-15.5	4.5	0.7	3.2-6.2	7.2	1.0	5.6-9.3
<b>Birthweight</b>									
Low	21.3	2.1	17.5-25.6	14.4	1.8	11.2-18.3	18.3	2.0	14.7-22.5
Normal	19.1	1.1	17.0-21.4	9.3	0.8	7.8-11.0	13.9	1.0	12.0-15.9

\* Confidence interval

-- Fewer than six respondents

**Table20c**Prevalence of Smoking During the Last Three Months of Pregnancy,  
2003- 2007

By Selected Sociodemographic Characteristics

	2003			2004**			2005			2006			2007		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI*									
<b>Total</b>	12.6	0.9	11-14.5	12.1	0.8	10.6-13.8	10.4	0.8	8.9-12.1	11.4	0.9	9.8-13.2	10.7	0.8	9.2-12.4
<b>Age</b>															
< 20 years	17.9	3.3	12.2-25.4	15.7	3.1	10.5-22.8	13.9	2.9	9.2-20.5	16.7	3.2	11.3-24.0	8.8	2.4	5.2-14.6
20-24 years	18.0	2.1	14.2-22.5	20.5	2.2	16.5-25.1	15.0	2.1	11.4-19.5	18.6	2.3	14.5-23.6	16.3	2.1	12.6-20.9
25-34 years	9.1	1.1	7.1-11.5	9.4	1.0	7.6-11.5	8.6	1.0	6.8-10.8	9.0	1.1	7.1-11.4	9.6	1.0	7.8-11.9
35+ years	12.7	2.5	8.6-18.3	7.9	1.6	5.2-11.8	7.6	1.8	4.8-11.9	6.0	1.6	3.5-10.2	7.9	1.8	5.0-12.1
<b>Race</b>															
Black	13.7	2.4	9.7-19.0	13.3	2.2	9.6-18.2	8.5	1.8	5.5-12.9	21.2	3.0	16.0-27.6	15.7	2.4	11.6-21.0
White	13.1	1.0	11.2-15.2	12.3	0.9	10.6-14.2	11.4	0.9	9.7-13.4	9.7	0.9	8.2-11.6	10.4	0.9	8.8-12.2
Other	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
<b>Hispanic Ethnicity</b>															
Hispanic	2.2	0.8	1.1-4.5	4.6	1.2	2.8-7.5	2.4	0.8	1.2-4.7	1.7	0.6	0.9-3.4	1.8	0.7	0.8-4.0
Non-Hispanic	16.1	1.2	13.9-18.5	14.3	1.0	12.4-16.3	13.1	1.0	11.2-15.2	14.9	1.1	12.8-17.3	13.3	1.0	11.5-15.4
<b>Education</b>															
< High School	17.5	2.2	13.6-22.2	21.3	2.4	17.1-26.3	12.1	1.9	8.9-16.3	17.7	2.3	13.6-22.8	18.2	2.3	14.0-23.2
High School	19.8	2.1	16.0-24.2	18.4	1.9	14.9-22.4	15.0	1.9	11.7-19.1	17.6	2.1	13.9-22.0	16.5	2.0	13.0-20.7
> High School	6.5	1.0	4.9-8.7	5.9	0.8	4.5-7.6	7.4	1.0	5.8-9.5	5.9	0.9	4.4-7.8	5.7	0.8	4.4-7.5
<b>Marital Status</b>															
Married	6.4	0.8	4.9-8.2	7.4	0.8	5.9-9.1	6.0	0.8	4.6-7.8	5.0	0.7	3.8-6.6	6.2	0.8	4.8-7.9
Other	23.5	1.9	19.9-27.5	20.6	1.7	17.5-24.2	17.2	1.6	14.3-20.6	22.1	1.9	18.6-26.1	17.5	1.6	14.6-20.9
<b>Payment for Delivery</b>															
Medicaid	18.3	1.6	15.4-21.5	20.1	1.5	17.3-23.2	14.2	1.3	11.9-17.0	18.0	1.5	15.2-21.2	15.1	1.3	12.7-17.8
Other	7.5	1.0	5.8-9.6	5.4	0.7	4.1-7.0	6.4	0.9	4.8-8.5	4.7	0.8	3.4-6.4	6.0	0.9	4.5-7.9
<b>Birthweight</b>															
Low	16.1	1.8	12.9-19.9	14.5	1.5	11.7-17.8	16.6	1.9	13.2-20.6	18.0	1.8	14.8-21.8	16.9	1.9	13.5-20.8
Normal	12.4	1.0	10.6-14.4	11.9	0.9	10.3-13.7	9.9	0.9	8.3-11.7	10.9	0.9	9.2-12.8	10.2	0.9	8.7-12.0

\*Confidence interval

\*\*Question Changed in 2004

--Fewer than six respondents

**Table 20d**  
 Prevalence of Smoking During Last Three Months of Pregnancy, 1998-2002  
 By Selected Sociodemographic Characteristics

	1998			1999			2000			2001			2002		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	13.8	0.9	12.2-15.6	12.8	0.8	11.3-14.5	12.5	0.8	11.0-14.2	12.6	0.8	11.1-14.2	11.4	0.8	10.0-13.0
<b>Age</b>															
< 20 years	18.8	2.6	14.3-24.3	18.6	2.8	13.8-24.6	12.6	2.4	8.7-18.0	15.3	2.5	11.1-20.8	11.9	2.4	8.0-17.4
20-24 years	16.6	2.0	13.0-20.9	14.0	1.7	11.0-17.8	18.3	2.0	14.6-22.6	15.1	1.9	11.8-19.1	16.0	1.9	12.6-20.1
25-34 years	11.7	1.2	9.6-14.2	10.7	1.0	8.9-12.8	9.9	1.0	8.1-12.0	10.5	1.0	8.6-12.6	9.1	1.0	7.3-11.2
35+ years	11.7	2.2	8.0-16.8	13.0	2.2	9.3-18.0	12.6	2.1	9.1-17.2	13.9	2.1	10.2-18.6	12.1	2.1	8.6-16.8
<b>Race</b>															
Black	10.9	1.7	8.0-14.8	9.8	1.6	7.0-13.4	13.0	2.0	9.7-17.4	14.4	2.0	10.9-18.8	8.3	1.6	5.7-12.0
White	14.8	1.0	12.9-16.9	14.3	1.0	12.5-16.3	12.7	0.9	11.0-14.6	12.7	0.9	11.1-14.6	12.6	0.9	10.9-14.5
Other	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
<b>Hispanic Ethnicity</b>															
Hispanic	2.6	0.9	1.4-5.0	4.5	1.2	2.6-7.4	3.8	1.1	2.2-6.5	1.8	0.7	0.8-3.7	3.2	0.9	1.9-5.4
Non-Hispanic	16.5	1.1	14.6-18.7	15.0	1.0	13.3-17.0	15.1	1.0	13.3-17.1	15.8	1.0	14.0-17.9	14.0	1.0	12.2-16.1
<b>Education</b>															
< High School	17.2	2.0	13.6-21.6	18.2	2.0	14.6-22.5	15.8	2.0	12.3-20.2	18.4	2.0	14.8-22.6	13.1	1.8	9.9-17.2
High School	20.3	1.9	16.9-24.2	18.5	1.7	15.4-22.1	18.0	1.7	14.9-21.6	17.3	1.7	14.3-20.8	18.0	1.9	14.7-22.0
> High School	7.9	1.0	6.2-10.0	7.2	0.9	5.7-9.1	7.7	0.9	6.2-9.7	7.5	0.9	6.0-9.5	7.6	0.9	6.0-9.5
<b>Marital Status</b>															
Married	9.9	0.9	8.3-11.9	10.1	0.9	8.6-12.0	9.0	0.8	7.5-10.7	8.0	0.8	6.6-9.7	8.5	0.9	7.0-10.4
Other	21.5	1.8	18.2-25.1	17.5	1.6	14.6-20.7	19.1	1.7	16.0-22.5	20.5	1.6	17.5-23.8	16.5	1.5	13.7-19.7
<b>Payment for Delivery</b>															
Medicaid	21.5	1.7	18.3-25.0	17.1	1.5	14.4-20.2	17.9	1.5	15.1-21.1	19.3	1.5	16.6-22.4	15.0	1.4	12.5-17.8
Other	9.6	0.9	7.9-11.6	10.2	0.9	8.6-12.1	9.1	0.9	7.6-11.0	7.9	0.8	6.4-9.7	8.6	0.9	7.0-10.6
<b>Birthweight</b>															
Low	20.6	1.9	17.1-24.5	21.0	2.4	16.8-26.0	19.2	1.9	15.8-23.2	18.1	1.8	14.8-21.9	16.2	1.7	13.1-19.9
Normal	13.3	0.9	11.6-15.2	12.0	1.0	10.2-14.0	12.0	0.9	10.4-13.8	12.2	0.8	10.6-13.9	11.1	0.8	9.5-12.8

\*Confidence interval

--Fewer than six respondents

**Table 21**  
 Prevalence of Drinking, 2009  
 By Selected Sociodemographic Characteristics

	Three Months Before Pregnancy			Last Three Months Of Pregnancy		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	53.3	1.4	50.6-56.0	6.8	0.7	5.6-8.3
<b>Age</b>						
< 20 years	33.6	4.2	26.0-42.2	--	--	--
20-24 years	48.9	3.1	42.8-55.1	4.8	1.4	2.7-8.3
25-34 years	59.2	1.8	55.6-62.8	7.8	1.0	6.1-10.0
35+ years	51.5	3.4	44.8-58.2	9.9	2.1	6.5-14.8
<b>Race</b>						
Black	43.0	3.8	35.8-50.4	4.2	1.6	2.0-8.6
White	57.4	1.5	54.4-60.4	7.4	0.8	6.0-9.2
Other	24.2	4.7	16.2-34.5	--	--	--
<b>Hispanic Ethnicity</b>						
Hispanic	28.0	2.5	23.5-33.1	3.5	1.0	2.0-6.0
Non- Hispanic	61.7	1.6	58.6-64.8	7.9	0.9	6.4-9.7
<b>Education</b>						
< High School	24.9	2.8	19.9-30.8	3.5	1.2	1.8-6.9
High School	44.3	2.8	38.8-49.9	3.6	1.1	2.0-6.4
> High School	67.2	1.7	63.8-70.5	9.5	1.1	7.6-11.9
<b>Marital Status</b>						
Married	56.6	1.7	53.2-59.9	7.9	0.9	6.2-9.9
Other	48.2	2.3	43.8-52.7	5.1	1.0	3.5-7.5
<b>Payment for Delivery</b>						
Medicaid	40.9	1.9	37.3-44.7	4.4	0.8	3.1-6.2
Other	67.8	1.9	64.0-71.3	9.6	1.2	7.6-12.2
<b>Birthweight</b>						
Low	50.8	2.7	45.5-56.1	2.9	0.9	1.6-5.2
Normal	53.5	1.5	50.6-56.4	7.1	0.7	5.8-8.7

\*Confidence interval  
 --Fewer than six respondents

**Table 22**

Number of Drinks Per Week Among Mothers Who Reported Drinking, 2009

	Three Months Before Pregnancy			Last Three Months Of Pregnancy		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
< 1 drink per week	53.0	1.9	49.3-56.6	83.0	3.9	73.9-89.3
1-3 drinks per week	30.5	1.7	27.3-34.0	12.1	3.4	6.9-20.5
4 or more drinks per week	16.5	1.4	13.9-19.4	--	--	--

\*Confidence interval

--Fewer than six respondents

**Table 23**

Question description: Women were given a list of feelings and experiences that women sometimes have after childbirth and asked to describe how often (using a scale of “Never” to “Always”) they felt or experienced things this way since their new baby was born.

	Never			Rarely			Sometimes			Often			Always		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
I felt down, depressed, or sad	36.1	1.3	33.5-38.7	28.2	1.2	25.8-30.6	27.6	1.2	25.2-30.0	6.6	0.7	5.3-8.0	1.7	0.4	1.1-2.5
I felt hopeless	72.1	1.2	69.6-74.4	15.0	1.0	13.1-17.0	8.4	0.8	7.0-10.0	3.5	0.5	2.6-4.7	1.1	0.3	0.7-1.8
I felt slowed down	40.9	1.4	38.3-43.6	18.9	1.1	16.9-21.1	26.7	1.2	24.3-29.1	10.9	0.9	9.3-12.7	2.6	0.4	1.9-3.6
I felt panicky	65.3	1.3	62.7-67.8	17.6	1.0	15.7-19.8	11.9	0.9	10.2-13.8	4.3	0.6	3.3-5.5	0.9	0.3	0.5-1.6
I felt restless	48.2	1.4	45.5-51.0	21.6	1.1	19.5-23.9	19.0	1.1	16.9-21.2	8.3	0.8	6.9-9.9	2.9	0.5	2.1-3.9

**Table 24**  
 Depression Diagnosis During Pregnancy, 2009  
 By Selected Sociodemographic Characteristics

	Percent	Standard Error	95% CI *
<b>Total</b>	7.4	0.7	6.1-9.0
<b>Age</b>			
< 20 years	8.7	2.5	4.9-14.8
20-24 years	8.3	1.8	5.4-12.5
25-34 years	6.6	0.9	5.0-8.7
35+ years	8.4	1.9	5.3-13.0
<b>Race</b>			
Black	11.0	2.4	7.1-16.8
White	6.8	0.8	5.5-8.5
Other	--	--	--
<b>Hispanic Ethnicity</b>			
Hispanic	4.8	1.1	3.0-7.5
Non- Hispanic	8.4	0.9	6.8-10.3
<b>Education</b>			
< High School	10.1	1.9	6.9-14.5
High School	8.3	1.6	5.7-11.9
> High School	6.3	0.9	4.7-8.3
<b>Marital Status</b>			
Married	6.3	0.9	4.8-8.2
Other	9.2	1.3	7.0-12.2
<b>Payment for Delivery</b>			
Medicaid	9.0	1.1	7.1-11.4
Other	5.5	0.9	4.0-7.6
<b>Birthweight</b>			
Low	11.5	1.7	8.6-15.3
Normal	7.1	0.8	5.7-8.8

\*Confidence interval  
 --Fewer than six respondents

**Table 25**  
 Postpartum Depression Diagnosis, 2009  
 By Selected Sociodemographic Characteristics

	Percent	Standard Error	95% CI *
<b>Total</b>	9.1	0.8	7.6-10.7
<b>Age</b>			
< 20 years	9.3	2.5	5.4-15.5
20-24 years	10.5	1.9	7.3-14.9
25-34 years	8.4	1.0	6.6-10.6
35+ years	9.6	2.0	6.3-14.3
<b>Race</b>			
Black	8.8	2.1	5.4-14.0
White	9.2	0.9	7.7-11.1
Other	7.9	3.0	3.7-16.1
<b>Hispanic Ethnicity</b>			
Hispanic	5.4	1.2	3.4-8.3
Non- Hispanic	10.3	1.0	8.6-12.4
<b>Education</b>			
< High School	11.1	2.0	7.8-15.6
High School	10.4	1.7	7.5-14.2
> High School	7.9	1.0	6.2-10.1
<b>Marital Status</b>			
Married	8.0	0.9	6.3-10.1
Other	10.7	1.4	8.3-13.7
<b>Payment for Delivery</b>			
Medicaid	11.2	1.2	9.0-13.7
Other	6.6	1.0	5.0-8.8
<b>Birthweight</b>			
Low	11.6	1.7	8.7-15.4
Normal	8.9	0.8	7.4-10.7

\*Confidence interval  
 --Fewer than six respondents

**Table 26**  
Infant Checkups, 2009

	Percent	Standard Error	95% CI *
Infant seen for one week checkup	95.6	0.6	94.3-96.6
Well-baby checkup at 2, 4 or 6 months	99.0	0.3	98.3-99.5

\*Confidence interval

**Table 27**  
Place of Well- baby Checkup, 2009  
By Selected Sociodemographic Characteristics

	Hospital Clinic			Health Department Clinic			MD/HMO			Comm. Health Center			Other		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	16.9	1.1	15.0-19.1	10.5	0.9	8.9-12.3	62.2	1.4	59.4-64.8	9.2	0.8	7.7-10.9	1.3	0.3	0.8-2.1
<b>Age</b>															
< 20 years	21.5	3.8	15.0-29.9	30.1	4.2	22.5-39.0	26.6	4.0	19.6-35.0	20.1	3.6	14.0-28.1	--	--	--
20-24 years	20.5	2.6	16.0-26.0	13.0	2.2	9.4-17.9	47.6	3.2	41.4-53.8	16.3	2.4	12.1-21.6	2.6	1.0	1.2-5.5
25-34 years	16.2	1.4	13.7-19.2	6.7	1.0	5.0-8.8	69.9	1.8	66.4-73.2	6.0	0.9	4.4-8.2	1.1	0.4	0.6-2.3
35+ years	11.9	2.3	8.0-17.3	7.7	1.9	4.8-12.2	76.5	3.0	70.1-81.9	3.9	1.4	1.9-8.0	--	--	--
<b>Race</b>															
Black	29.4	3.6	22.9-36.9	18.0	3.0	12.9-24.6	34.7	3.7	27.8-42.3	17.0	3.0	11.9-23.7	--	--	--
White	13.9	1.1	12.0-16.2	9.0	0.9	7.4-10.9	67.4	1.5	64.5-70.2	8.2	0.9	6.6-10.0	1.5	0.4	0.9-2.5
Other	22.9	4.7	15.1-33.3	10.4	3.7	5.1-20.2	64.9	5.4	53.8-74.6	--	--	--	--	--	--
<b>Hispanic Ethnicity</b>															
Hispanic	24.1	2.4	19.7-29.1	21.9	2.3	17.7-26.8	34.2	2.6	29.3-39.6	17.0	2.1	13.2-21.6	2.8	0.9	1.5-5.3
Non-Hispanic	14.6	1.2	12.4-17.0	6.7	0.8	5.2-8.5	71.4	1.5	68.4-74.2	6.6	0.9	5.1-8.5	0.8	0.3	0.4-1.7
<b>Education</b>															
< High School	21.1	2.7	16.4-26.9	26.0	2.9	20.7-32.1	29.2	3.0	23.7-35.4	19.2	2.6	14.6-24.9	4.5	1.3	2.5-7.9
High School	22.4	2.4	18.0-27.4	13.2	1.9	9.8-17.5	50.6	2.9	45.0-56.2	13.2	2.0	9.8-17.5	--	--	--
> High School	12.7	1.3	10.4-15.4	4.0	0.7	2.8-5.8	79.1	1.5	75.9-82.0	3.7	0.7	2.5-5.5	--	--	--
<b>Marital Status</b>															
Married	13.9	1.2	11.7-16.5	6.4	0.9	4.8-8.3	75.2	1.5	72.1-78.1	3.7	0.7	2.5-5.2	0.9	0.3	0.4-1.8
Other	21.7	1.9	18.1-25.7	16.9	1.7	13.8-20.6	41.6	2.3	37.2-46.1	17.8	1.8	14.6-21.6	2.0	0.6	1.1-3.8
<b>Payment for Delivery</b>															
Medicaid	22.3	1.6	19.3-25.6	17.6	1.5	14.9-20.7	41.8	1.9	38.2-45.6	16.0	1.5	13.3-19.1	2.3	0.6	1.4-3.7
Other	10.6	1.3	8.4-13.4	2.2	0.6	1.2-3.8	85.8	1.4	82.7-88.4	1.3	0.5	0.6-2.5	--	--	--
<b>Birthweight</b>															
Low	19.5	2.3	15.4-24.4	10.9	1.8	7.8-15.0	61.3	2.8	55.8-66.6	7.3	1.5	4.8-11.0	--	--	--
Normal	16.8	1.1	14.7-19.1	10.4	0.9	8.7-12.4	62.2	1.5	59.3-65.0	9.3	0.9	7.7-11.2	1.3	0.3	0.8-2.2

\*Confidence interval  
-- Fewer than six respondents

**Table 28**  
 Infant Sleep Practices, 2009

	Percent	Standard Error	95% CI *
My new baby sleeps in a crib or portable crib	85.9	1.0	83.9-87.8
My new baby sleeps on a firm or hard mattress	87.2	1.0	85.2-88.9
My new baby sleeps with pillows	10.4	0.9	8.8-12.2
My new baby sleeps with bumper pads	41.6	1.4	38.9-44.3
My new baby sleeps with plush blankets	22.6	1.2	20.4-25.0
My new baby sleeps with stuffed toys	4.2	0.6	3.2-5.5
My new baby sleeps with another person	17.9	1.1	15.8-20.1

\*Confidence interval

**Table 29**  
Sleep Position of Infant, 2009

	Percent	Standard Error	95% CI *
Back	71.6	1.3	69.1-74.0
Side	14.5	1.0	12.7-16.6
Stomach	9.1	0.8	7.6-10.7
Other positions	4.8	0.6	3.7-6.1

\*Confidence interval

**Table 30a**  
 Prevalence of Laying Infant on Back to Sleep, 2008-2009  
 By Selected Sociodemographic Characteristics

	2008			2009		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI*
<b>Total</b>	70.0	1.2	67.5-72.3	71.6	1.3	69.1-74.0
<b>Age</b>						
< 20 years	58.0	4.1	49.7-65.8	58.0	4.5	49.1-66.5
20-24 years	63.4	2.9	57.7-68.8	63.6	3.1	57.4-69.4
25-34 years	73.1	1.6	69.8-76.2	75.5	1.6	72.2-78.5
35+ years	76.5	2.9	70.4-81.7	77.1	2.9	70.9-82.3
<b>Race</b>						
Black	59.8	3.7	52.4-66.7	56.2	3.9	48.6-63.6
White	72.0	1.3	69.3-74.5	74.1	1.4	71.4-76.7
Other	72.1	5.3	60.8-81.3	78.5	4.5	68.3-86.0
<b>Hispanic Ethnicity</b>						
Hispanic	63.5	2.4	58.7-68.1	66.8	2.6	61.6-71.7
Non- Hispanic	72.3	1.4	69.5-75.1	73.3	1.4	70.4-76.0
<b>Education</b>						
< High School	59.1	3.0	53.1-64.9	61.1	3.2	54.7-67.2
High School	66.2	2.5	61.1-70.9	63.7	2.7	58.2-68.9
> High School	75.9	1.6	72.7-78.8	78.9	1.5	75.8-81.7
<b>Marital Status</b>						
Married	73.9	1.5	70.9-76.7	77.4	1.5	74.4-80.1
Other	64.2	2.1	60.0-68.2	62.7	2.2	58.2-66.9
<b>Payment for Delivery</b>						
Medicaid	63.8	1.8	60.1-67.4	65.7	1.8	62.0-69.2
Other	76.3	1.6	73.0-79.3	78.8	1.6	75.4-81.8
<b>Birthweight</b>						
Low	65.6	2.5	60.6-70.3	72.8	2.5	67.5-77.5
Normal	70.3	1.3	67.6-72.8	71.6	1.3	68.9-74.1

**Table 30b**  
Prevalence of Laying Infant on Back to Sleep, 2003- 2007  
By Selected Sociodemographic Characteristics

	2003			2004			2005			2006			2007		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	63.7	1.3	61.1-66.3	65.7	1.2	63.3-68.0	68.3	1.3	65.7-70.7	70.5	1.3	68.0-72.9	70.7	1.2	68.2-73.0
<b>Age</b>															
< 20 years	47.0	4.4	38.6-55.6	57.5	4.3	48.9-65.7	53.2	4.2	44.9-61.3	63.4	4.3	54.7-71.3	58.7	4.2	50.3-66.6
20-24 years	59.6	2.8	54.1-64.9	56.2	2.8	50.7-61.6	64.4	2.8	58.6-69.7	66.5	2.9	60.7-71.9	61.6	2.8	55.9-67.0
25-34 years	67.1	1.8	63.5-70.6	67.7	1.6	64.6-70.7	72.4	1.7	68.9-75.5	71.7	1.7	68.4-74.9	73.4	1.6	70.2-76.4
35+ years	70.9	3.5	63.6-77.3	76.0	2.7	70.4-80.9	70.3	3.2	63.7-76.1	76.6	2.9	70.5-81.8	82.8	2.6	77.2-87.3
<b>Race</b>															
Black	46.1	3.5	39.4-53.0	48.1	3.2	41.9-54.4	49.4	3.5	42.7-56.2	54.3	3.7	47.1-61.4	53.6	3.4	47.0-60.1
White	66.8	1.5	63.8-69.6	69.3	1.3	66.7-71.8	72.4	1.4	69.6-74.9	73.6	1.3	70.9-76.1	74.2	1.3	71.6-76.7
Other	76.7	5.4	64.7-85.5	70.7	5.5	59.0-80.1	73.9	5.7	61.3-83.5	78.0	4.7	67.4-85.8	79.4	4.1	70.2-86.4
<b>Hispanic Ethnicity</b>															
Hispanic	57.3	2.7	51.9-62.5	56.8	2.6	51.7-61.7	66.1	2.5	61.1-70.8	65.7	2.4	60.8-70.3	68.4	2.6	63.1-73.3
Non- Hispanic	65.9	1.5	62.8-68.8	68.4	1.4	65.7-71.0	69.0	1.5	66.0-71.8	72.3	1.5	69.4-75.1	71.3	1.4	68.5-73.9
<b>Education</b>															
< High School	57.1	2.9	51.3-62.7	50.9	2.8	45.4-56.5	60.7	2.9	54.9-66.1	60.9	2.9	55.0-66.4	60.2	3.1	54.0-66.1
High School	58.7	2.6	53.5-63.7	60.1	2.5	55.1-64.9	65.9	2.6	60.7-70.8	64.0	2.7	58.7-69.1	64.1	2.6	58.9-69.0
> High School	69.6	1.8	65.9-73.0	73.9	1.5	70.9-76.7	73.0	1.7	69.6-76.2	77.3	1.6	74.1-80.2	76.9	1.5	73.9-79.7
<b>Marital Status</b>															
Married	67.9	1.6	64.6-71.0	71.0	1.4	68.1-73.6	73.7	1.5	70.6-76.6	74.8	1.4	71.8-77.5	76.9	1.4	74.1-79.6
Other	56.5	2.3	51.9-60.9	55.9	2.2	51.6-60.1	59.7	2.2	55.4-64.0	63.4	2.3	58.9-67.7	61.1	2.1	56.8-65.1
<b>Payment for Delivery</b>															
Medicaid	57.6	2.0	53.7-61.5	56.8	1.9	53.1-60.5	61.6	1.9	57.8-65.2	63.5	1.9	59.7-67.2	63.0	1.8	59.3-66.4
Other	69.2	1.8	65.7-72.6	73.1	1.5	70.1-75.9	75.4	1.7	72.0-78.5	77.3	1.6	74.1-80.2	79.0	1.5	75.9-81.8
<b>Birthweight</b>															
Low	61.4	2.5	56.4-66.2	65.2	2.2	60.8-69.3	69.3	2.3	64.6-73.7	70.3	2.2	65.8-74.5	71.1	2.4	66.2-75.5
Normal	63.9	1.4	61.0-66.6	65.7	1.3	63.2-68.2	68.2	1.4	65.5-70.8	70.5	1.3	67.9-73.1	70.6	1.3	68.0-73.1

\*Confidence interval

**Table 30c**  
Prevalence of Laying Infant on Back to Sleep, 1998-2002  
By Selected Sociodemographic Characteristics

	1998			1999			2000**			2001			2002		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	53.8	1.3	51.4-56.2	56.5	1.2	54.1-58.9	60.9	1.2	58.6-63.2	63.6	1.2	61.3-65.9	63.6	1.2	61.2-65.9
<b>Age</b>															
< 20 years	40.8	3.4	34.4-47.5	42.4	3.5	35.6-49.4	54.9	3.7	47.7-62.0	53.4	3.5	46.4-60.2	50.2	3.7	43.0-57.4
20-24 years	50.3	2.7	45.1-55.5	53.1	2.6	48.0-58.1	56.1	2.6	50.9-61.2	58.4	2.6	53.2-63.3	56.3	2.6	51.2-61.3
25-34 years	56.7	1.8	53.2-60.0	59.4	1.7	56.1-62.6	63.1	1.6	59.8-66.2	66.3	1.6	63.1-69.4	66.1	1.6	62.8-69.2
35+ years	62.3	3.3	55.7-68.6	66.2	3.2	59.7-72.1	65.7	3.0	59.6-71.4	70.4	2.9	64.5-75.7	76.3	2.7	70.5-81.2
<b>Race</b>															
Black	40.2	2.8	34.9-45.8	43.4	2.9	37.8-49.1	45.1	3.0	39.4-51.0	44.7	2.9	39.0-50.5	38.8	3.0	33.2-44.7
White	57.3	1.4	54.5-60.0	59.2	1.4	56.5-61.9	64.6	1.3	62.0-67.1	68.1	1.3	65.6-70.6	69.0	1.3	66.4-71.5
Other	48.7	8.2	33.3-64.4	68.2	5.2	57.2-77.4	60.0	6.6	46.6-72.0	67.2	5.6	55.5-77.1	62.4	6.0	50.2-73.2
<b>Hispanic Ethnicity</b>															
Hispanic	51.6	2.8	46.2-57.0	51.2	2.8	45.7-56.7	56.9	2.6	51.7-61.9	59.1	2.5	54.1-64.0	58.0	2.5	53.0-62.9
Non- Hispanic	54.3	1.4	51.6-57.0	57.9	1.4	55.3-60.6	62.2	1.3	59.5-64.7	64.9	1.3	62.3-67.5	65.2	1.4	62.5-67.9
<b>Education</b>															
< High School	41.7	2.8	36.3-47.3	44.9	2.7	39.7-50.3	52.3	2.8	46.8-57.7	53.0	2.7	47.7-58.3	47.1	2.7	41.8-52.4
High School	50.1	2.3	45.6-54.6	53.1	2.3	48.5-57.6	57.0	2.3	52.5-61.3	62.3	2.2	58.0-66.5	58.7	2.4	54.0-63.4
> High School	61.6	1.7	58.2-64.9	63.8	1.7	60.5-67.0	67.4	1.6	64.2-70.4	68.5	1.6	65.3-71.6	73.0	1.5	70.0-75.9
<b>Marital Status</b>															
Married	58.2	1.5	55.3-61.2	61.0	1.5	58.1-63.8	65.1	1.4	62.3-67.8	69.2	1.4	66.4-71.9	69.5	1.4	66.7-72.2
Other	44.7	2.2	40.5-49.0	48.8	2.1	44.6-53.0	53.0	2.2	48.8-57.3	53.9	2.0	49.9-57.9	53.1	2.1	49.0-57.2
<b>Payment for Delivery</b>															
Medicaid	45.7	2.1	41.5-49.9	48.9	2.1	44.8-53.0	53.2	2.1	49.2-57.2	53.3	1.9	49.5-57.0	53.7	1.9	50.0-57.4
Other	58.1	1.5	55.0-61.0	61.2	1.5	58.2-64.0	65.7	1.4	62.9-68.5	70.9	1.4	68.0-73.6	71.1	1.5	68.2-73.9
<b>Birthweight</b>															
Low	49.5	2.5	44.6-54.5	49.4	3.0	43.6-55.3	52.9	2.5	48.0-57.7	55.8	2.4	51.1-60.4	56.3	2.4	51.5-61.0
Normal	54.1	1.3	51.5-56.7	56.8	1.5	53.8-59.6	61.5	1.3	59.0-63.9	64.1	1.2	61.7-66.5	64.1	1.3	61.5-66.5

\*Confidence interval

\*\*Question changed in 2000

**Table 31a**  
Breastfeeding Initiation, 2008-2009  
By Selected Sociodemographic Characteristics

	2008			2009		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI*
<b>Total</b>	77.8	1.1	75.5-79.9	79.9	1.1	77.6-82.0
<b>Age</b>						
< 20 years	67.9	3.9	59.8-75.1	69.5	4.1	60.9-77.0
20-24 years	70.4	2.8	64.7-75.5	72.8	2.8	66.9-78.0
25-34 years	80.8	1.5	77.7-83.5	83.7	1.4	80.8-86.2
35+ years	84.8	2.5	79.2-89.1	82.4	2.7	76.5-87.0
<b>Race</b>						
Black	56.0	3.7	48.6-63.1	64.0	3.7	56.5-70.9
White	81.6	1.1	79.2-83.7	81.9	1.2	79.4-84.1
Other	94.1	2.7	85.9-97.7	95.4	2.4	87.6-98.4
<b>Hispanic Ethnicity</b>						
Hispanic	86.6	1.7	82.9-89.6	86.4	1.8	82.4-89.7
Non- Hispanic	74.4	1.4	71.5-77.1	77.6	1.4	74.8-80.1
<b>Education</b>						
< High School	68.6	2.9	62.6-74.0	72.4	2.9	66.3-77.7
High School	67.7	2.5	62.6-72.4	69.3	2.6	64.0-74.2
> High School	86.2	1.3	83.5-88.5	86.8	1.3	84.2-89.1
<b>Marital Status</b>						
Married	86.7	1.1	84.4-88.8	86.8	1.2	84.3-89.0
Other	64.9	2.1	60.6-68.9	69.0	2.1	64.7-73.0
<b>Payment for Delivery</b>						
Medicaid	69.8	1.8	66.2-73.2	73.4	1.7	69.9-76.6
Other	86.3	1.3	83.6-88.7	87.4	1.3	84.6-89.7
<b>Birthweight</b>						
Low	75.1	2.3	70.3-79.3	76.0	2.4	70.9-80.4
Normal	78.0	1.2	75.5-80.3	80.1	1.2	77.7-82.4

**Table 31b**  
Breastfeeding Initiation, 2003-2007  
By Selected Sociodemographic Characteristics

	2003			2004			2005			2006			2007		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	72.0	1.3	69.5-74.4	75.0	1.1	72.8-77.1	76.2	1.2	73.8-78.4	76.7	1.2	74.3-78.9	74.8	1.2	72.5-77.1
<b>Age</b>															
< 20 years	55.3	4.4	46.7-63.7	55.6	4.3	47.0-63.9	55.7	4.2	47.4-63.8	58.2	4.4	49.3-66.5	57.3	4.2	48.9-65.2
20-24 years	63.9	2.7	58.5-69.0	64.6	2.7	59.2-69.7	70.8	2.7	65.3-75.8	71.4	2.8	65.7-76.5	68.7	2.7	63.1-73.8
25-34 years	78.0	1.6	74.7-81.0	79.2	1.4	76.4-81.7	80.7	1.5	77.6-83.4	80.2	1.5	77.2-83.0	79.2	1.5	76.2-81.9
35+ years	76.2	3.3	69.2-82.0	85.2	2.2	80.3-89.0	82.3	2.7	76.5-86.9	83.9	2.5	78.3-88.3	80.8	2.8	74.9-85.6
<b>Race</b>															
Black	52.6	3.5	45.7-59.3	56.6	3.2	50.2-62.7	57.8	3.4	51.0-64.4	55.8	3.7	48.5-62.8	52.5	3.4	45.9-59.0
White	75.4	1.3	72.7-78.0	78.2	1.2	75.9-80.4	79.7	1.2	77.2-82.0	80.3	1.2	77.9-82.5	78.4	1.2	75.9-80.7
Other	84.9	4.7	73.2-92.0	91.8	3.3	82.7-96.3	90.4	3.9	79.6-95.8	91.1	3.3	82.0-95.8	97.3	1.6	91.6-99.2
<b>Hispanic Ethnicity</b>															
Hispanic	81.1	2.2	76.5-85.0	83.6	1.9	79.5-87.0	84.1	1.9	80.0-87.5	85.1	1.8	81.2-88.4	86.3	1.9	82.1-89.6
Non- Hispanic	68.9	1.5	66.0-71.8	72.4	1.3	69.8-74.9	73.4	1.4	70.5-76.1	73.6	1.4	70.7-76.3	71.5	1.4	68.7-74.1
<b>Education</b>															
< High School	63.0	2.9	57.3-68.4	64.5	2.7	59.0-69.6	66.1	2.8	60.4-71.4	68.7	2.9	62.8-74.0	65.8	3.0	59.7-71.4
High School	65.5	2.5	60.5-70.3	64.9	2.4	60.0-69.5	67.7	2.6	62.5-72.6	65.6	2.6	60.3-70.5	62.8	2.6	57.6-67.8
> High School	79.6	1.6	76.3-82.6	83.3	1.3	80.7-85.6	84.3	1.4	81.5-86.8	84.8	1.3	82.0-87.3	83.2	1.3	80.5-85.7
<b>Marital Status</b>															
Married	80.0	1.4	77.1-82.5	82.6	1.2	80.2-84.7	83.5	1.3	80.9-85.9	83.6	1.2	81.0-85.9	83.5	1.2	81.0-85.8
Other	58.0	2.3	53.5-62.4	60.9	2.1	56.7-64.9	64.5	2.1	60.2-68.6	65.0	2.3	60.5-69.3	61.6	2.1	57.3-65.6
<b>Payment for Delivery</b>															
Medicaid	65.8	1.9	61.9-69.5	68.4	1.8	64.8-71.7	68.0	1.8	64.4-71.5	69.4	1.9	65.7-73.0	66.3	1.8	62.7-69.7
Other	77.6	1.6	74.4-80.6	80.5	1.3	77.8-83.0	84.7	1.4	81.7-87.2	83.9	1.4	81.1-86.5	84.1	1.4	81.3-86.6
<b>Birthweight</b>															
Low	70.1	2.4	65.3-74.6	65.7	2.2	61.3-69.8	72.4	2.3	67.7-76.7	74.9	2.1	70.6-78.7	76.9	2.2	72.3-80.9
Normal	72.1	1.3	69.5-74.7	75.7	1.2	73.3-77.9	76.4	1.3	73.9-78.8	76.8	1.3	74.2-79.2	74.7	1.2	72.2-77.0

\*Confidence interval

**Table 31c**  
Breastfeeding Initiation, 1998-2002  
By Selected Sociodemographic Characteristics

	1998			1999			2000**			2001			2002		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	66.5	1.2	64.2-68.7	62.6	1.2	60.2-64.9	69.0	1.0	66.8-71.2	68.6	1.1	66.3-70.8	73.8	1.1	71.6-75.9
<b>Age</b>															
< 20 years	46.0	3.4	39.5-52.6	43.8	3.5	37.1-50.8	53.9	3.7	46.6-61.0	48.1	3.6	41.3-55.1	53.6	3.7	46.3-60.7
20-24 years	60.4	2.6	55.2-65.3	57.9	2.5	52.9-62.8	61.1	2.6	55.8-66.0	61.0	2.6	55.9-65.9	67.6	2.5	62.6-72.2
25-34 years	73.0	1.6	69.9-76.0	68.0	1.6	64.9-71.0	74.2	1.5	71.1-77.0	73.2	1.5	70.2-76.1	79.0	1.4	76.2-81.7
35+ years	72.7	3.0	66.4-78.3	68.9	3.1	62.6-74.6	75.1	2.7	69.4-80.1	80.6	2.4	75.4-84.9	80.9	2.5	75.5-85.4
<b>Race</b>															
Black	43.0	2.8	37.6-48.6	32.8	2.7	27.8-38.3	50.1	3.0	44.3-55.9	47.5	2.9	41.8-53.3	48.5	3.0	42.6-54.4
White	71.6	1.3	69.0-74.0	68.9	1.3	66.3-71.4	72.5	1.2	70.1-74.9	72.7	1.2	70.2-75.0	78.5	1.2	76.1-80.6
Other	87.2	4.8	74.5-94.1	86.1	3.8	76.8-92.1	90.2	4.1	78.7-95.9	92.0	3.3	82.8-96.5	93.1	3.3	83.0-97.4
<b>Hispanic Ethnicity</b>															
Hispanic	77.5	2.3	72.7-81.7	75.8	2.4	70.7-80.3	77.4	2.2	72.8-81.5	78.1	2.2	73.5-82.0	84.8	1.8	80.8-88.0
Non-Hispanic	63.8	1.3	61.2-66.3	59.2	1.3	56.6-61.8	66.5	1.3	63.9-69.1	65.8	1.3	63.2-68.3	70.3	1.3	67.7-72.8
<b>Education</b>															
< High School	56.3	2.8	50.9-61.6	50.9	2.7	45.5-56.2	59.7	2.8	54.1-65.0	55.3	2.7	50.0-60.6	66.5	2.6	61.3-71.4
High School	54.3	2.3	49.9-58.6	51.4	2.3	46.9-55.9	56.5	2.3	52.0-60.8	59.5	2.2	55.1-63.8	62.7	2.4	58.0-67.2
> High School	78.3	1.5	75.3-81.0	73.8	1.5	70.7-76.6	80.1	1.4	77.3-82.6	79.8	1.4	77.0-82.4	82.5	1.3	79.8-84.9
<b>Marital Status</b>															
Married	76.2	1.3	73.6-78.7	72.8	1.3	70.1-75.3	78.4	1.2	75.9-80.6	78.3	1.3	75.7-80.6	81.5	1.2	79.0-83.7
Other	46.6	2.1	42.4-50.8	45.0	2.1	40.9-49.1	51.5	2.2	47.2-55.8	52.0	2.0	48.0-56.0	60.2	2.1	56.1-64.2
<b>Payment for Delivery</b>															
Medicaid	53.1	2.1	49.1-57.1	50.7	2.1	46.6-54.7	57.9	2.0	53.9-61.9	55.8	1.9	52.0-59.5	65.6	1.8	61.9-69.0
Other	73.5	1.4	70.7-76.1	69.7	1.4	67.0-72.4	75.8	1.3	73.1-78.2	77.4	1.3	74.7-79.8	80.2	1.3	77.6-82.7
<b>Birthweight</b>															
Low	54.6	2.5	49.7-59.5	55.1	3.0	49.3-60.8	62.7	2.4	57.8-67.3	64.7	2.3	60.1-69.1	68.4	2.3	63.8-72.7
Normal	67.3	1.2	64.9-69.6	64.4	1.4	61.6-67.2	69.5	1.2	67.0-71.8	68.9	1.2	66.5-71.2	74.2	1.2	71.8-76.4

\*Confidence interval

\*\*Question changed in 2000

**Table 32a**  
Breastfeeding at Four Weeks After Delivery, 2008-2009  
By Selected Sociodemographic Characteristics

	2008			2009		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI*
<b>Total</b>	65.8	1.3	63.3-68.3	66.7	1.3	64.1-69.2
<b>Age</b>						
< 20 years	40.1	4.1	32.4-48.4	42.9	4.5	34.5-51.8
20-24 years	56.9	3.0	51.0-62.6	55.8	3.1	49.6-61.8
25-34 years	71.9	1.7	68.5-75.0	73.6	1.7	70.2-76.7
35+ years	76.1	3.0	69.8-81.5	72.4	3.1	65.9-78.1
<b>Race</b>						
Black	41.9	3.7	34.9-49.4	47.8	3.9	40.3-55.4
White	69.9	1.4	67.2-72.5	69.1	1.4	66.3-71.8
Other	85.0	4.2	74.7-91.6	86.6	3.9	77.1-92.6
<b>Hispanic Ethnicity</b>						
Hispanic	73.2	2.2	68.6-77.3	69.8	2.5	64.7-74.5
Non- Hispanic	63.1	1.6	60.0-66.1	65.6	1.5	62.6-68.6
<b>Education</b>						
< High School	53.1	3.1	47.1-59.1	57.4	3.2	51.0-63.5
High School	54.6	2.7	49.3-59.7	51.5	2.8	46.0-57.0
> High School	76.3	1.6	73.1-79.3	76.5	1.6	73.3-79.4
<b>Marital Status</b>						
Married	77.9	1.4	75.0-80.5	77.5	1.5	74.6-80.3
Other	48.6	2.2	44.3-52.9	49.9	2.3	45.5-54.4
<b>Payment for Delivery</b>						
Medicaid	55.2	1.9	51.5-59.0	57.2	1.9	53.4-60.9
Other	77.4	1.6	74.2-80.4	78.0	1.7	74.6-81.1
<b>Birthweight</b>						
Low	62.4	2.5	57.3-67.2	63.7	2.7	58.3-68.8
Normal	66.1	1.4	63.4-68.7	67.0	1.4	64.2-69.6

\*Confidence interval

**Table 32b**  
Breastfeeding Four Weeks After Delivery, 2003-2007  
By Selected Sociodemographic Characteristics

	2003			2004			2005			2006			2007		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	58.4	1.4	55.7-61.1	62.4	1.2	60.0-64.8	61.8	1.3	59.2-64.4	64.4	1.3	61.8-66.9	62.4	1.3	59.8-64.9
<b>Age</b>															
< 20 years	31.4	4.0	24.1-39.8	36.0	4.2	28.2-44.6	32.2	3.9	25.0-40.3	42.1	4.4	33.7-50.9	40.7	4.1	32.9-49.0
20-24 years	48.0	2.8	42.5-53.5	49.8	2.8	44.3-55.2	52.5	3.0	46.7-58.3	52.9	3.1	46.9-58.8	51.5	2.9	45.8-57.1
25-34 years	66.5	1.8	62.9-70.0	67.2	1.6	64.1-70.3	68.6	1.8	65.0-71.9	69.3	1.7	65.8-72.5	69.2	1.7	65.9-72.3
35+ years	66.6	3.6	59.2-73.3	77.4	2.6	71.8-82.1	72.1	3.1	65.6-77.8	77.9	2.9	71.8-83.0	69.6	3.2	63.0-75.5
<b>Race</b>															
Black	39.3	3.4	32.8-46.1	39.9	3.2	33.8-46.2	43.5	3.5	36.9-50.4	46.0	3.7	39.0-53.2	44.2	3.4	37.7-50.8
White	61.4	1.5	58.4-64.3	66.1	1.3	63.5-68.6	65.4	1.5	62.5-68.1	66.8	1.4	64.0-69.5	65.1	1.4	62.3-67.9
Other	78.4	5.4	66.1-87.1	86.2	4.1	76.1-92.4	75.8	5.7	63.1-85.2	89.6	3.6	80.1-94.8	83.9	3.8	74.9-90.1
<b>Hispanic Ethnicity</b>															
Hispanic	69.0	2.6	63.8-73.8	70.2	2.4	65.3-74.6	68.5	2.5	63.4-73.2	72.0	2.3	67.3-76.3	69.4	2.6	64.2-74.2
Non-Hispanic	54.8	1.6	51.6-57.9	60.1	1.4	57.3-62.8	59.5	1.6	56.4-62.6	61.6	1.6	58.5-64.6	60.3	1.5	57.4-63.2
<b>Education</b>															
< High School	51.8	2.9	46.1-57.5	50.0	2.8	44.5-55.6	49.5	3.0	43.7-55.3	54.5	3.0	48.5-60.3	49.2	3.2	43.0-55.4
High School	47.6	2.7	42.4-52.8	50.8	2.6	45.8-55.8	49.8	2.8	44.4-55.2	50.5	2.8	45.1-55.8	49.0	2.7	43.8-54.2
> High School	67.3	1.9	63.6-70.8	72.2	1.5	69.1-75.0	72.6	1.7	69.1-75.7	74.4	1.6	71.1-77.4	72.7	1.6	69.5-75.6
<b>Marital Status</b>															
Married	68.5	1.6	65.3-71.6	72.3	1.4	69.6-75.0	71.4	1.6	68.2-74.4	71.8	1.5	68.8-74.7	72.9	1.5	69.9-75.6
Other	40.8	2.3	36.4-45.3	43.8	2.2	39.6-48.0	46.8	2.2	42.4-51.2	51.7	2.4	47.1-56.3	46.6	2.2	42.4-50.8
<b>Payment for Delivery</b>															
Medicaid	51.1	2.0	47.1-55.1	54.3	1.9	50.6-57.9	50.6	2.0	46.8-54.4	55.2	2.0	51.3-59.1	52.4	1.9	48.7-56.0
Other	64.9	1.8	61.3-68.4	69.1	1.6	66.0-72.1	73.4	1.7	69.9-76.6	73.6	1.7	70.3-76.7	73.2	1.6	69.9-76.3
<b>Birthweight</b>															
Low	57.0	2.6	51.9-61.9	52.5	2.3	48.1-57.0	57.6	2.5	52.6-62.4	57.3	2.4	52.5-61.9	64.3	2.5	59.3-69.1
Normal	58.5	1.5	55.6-61.3	63.1	1.3	60.5-65.6	62.1	1.4	59.3-64.9	64.9	1.4	62.1-67.6	62.2	1.4	59.5-64.9

\* Confidence interval

**Table 32c**  
Breastfeeding At Four Weeks After Delivery, 1998-2002  
By Selected Sociodemographic Characteristics

	1998			1999			2000**			2001			2002		
	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *	Percent	Standard Error	95% CI *
<b>Total</b>	55.0	1.2	52.6-57.4	49.2	1.2	46.8-51.5	56.0	1.2	53.6-58.4	56.7	1.2	54.3-59.1	60.2	1.2	57.8-62.5
<b>Age</b>															
< 20 years	34.9	3.3	28.8-41.5	27.4	3.1	21.7-34.0	31.5	3.5	25.2-38.7	34.1	3.4	27.8-41.0	31.3	3.4	25.1-38.4
20-24 years	45.2	2.6	40.1-50.4	41.2	2.5	36.4-46.2	46.2	2.6	41.1-51.4	45.5	2.6	40.4-50.6	48.6	2.6	43.5-53.7
25-34 years	62.3	1.7	58.9-65.6	56.5	1.7	53.2-59.8	62.6	1.6	59.4-65.8	62.8	1.7	59.5-66.0	68.2	1.6	65.0-71.3
35+ years	64.3	3.3	57.7-70.5	57.1	3.3	50.6-63.3	67.7	3.0	61.5-73.2	71.1	2.8	65.2-76.3	72.6	2.9	66.6-77.9
<b>Race</b>															
Black	34.3	2.7	29.2-39.8	23.3	2.4	18.9-28.3	35.0	2.9	29.7-40.8	35.6	2.9	30.2-41.4	34.2	2.9	28.8-40.0
White	59.6	1.4	56.9-62.3	54.4	1.4	51.7-57.1	59.9	1.4	57.2-62.5	60.4	1.3	57.7-63.0	64.7	1.3	62.0-67.3
Other	69.7	8.0	52.4-82.9	74.1	4.9	63.4-82.6	81.3	5.5	68.3-89.8	87.1	4.1	76.7-93.2	85.4	4.4	74.6-92.1
<b>Hispanic Ethnicity</b>															
Hispanic	62.5	2.7	57.2-67.6	60.5	2.8	54.9-65.7	61.3	2.6	56.2-66.2	64.2	2.5	59.2-68.9	68.4	2.4	63.6-72.8
Non-Hispanic	53.2	1.4	50.5-55.8	46.3	1.3	43.7-48.9	54.4	1.4	51.7-57.1	54.5	1.4	51.8-57.2	57.5	1.4	54.7-60.3
<b>Education</b>															
< High School	46.1	2.7	40.7-51.5	35.5	2.6	30.6-40.7	46.1	2.8	40.7-51.6	44.2	2.7	39.0-49.6	49.9	2.7	44.6-55.2
High School	41.3	2.2	37.1-45.7	38.1	2.2	33.9-42.5	41.4	2.2	37.1-45.9	45.7	2.3	41.3-50.1	46.4	2.4	41.6-51.2
> High School	67.3	1.6	64.0-70.4	60.9	1.7	57.6-64.1	68.7	1.6	65.5-71.7	68.7	1.6	65.4-71.7	71.3	1.6	68.2-74.2
<b>Marital Status</b>															
Married	65.2	1.5	62.3-68.0	59.9	1.4	57.0-62.7	67.5	1.4	64.7-70.1	67.8	1.4	65.0-70.5	70.6	1.4	67.8-73.3
Other	34.2	2.1	30.3-38.3	30.7	2.0	27.0-34.6	34.6	2.1	30.7-38.8	37.4	2.0	33.6-41.4	41.7	2.1	37.7-45.8
<b>Payment for Delivery</b>															
Medicaid	40.8	2.0	36.9-44.9	35.1	2.0	31.4-39.1	42.0	2.0	38.0-46.0	41.8	1.9	38.1-45.6	47.9	1.9	44.2-51.6
Other	62.5	1.5	59.5-65.4	57.6	1.5	54.7-60.5	64.7	1.5	61.8-67.5	67.0	1.5	64.0-69.8	69.7	1.5	66.6-72.5
<b>Birthweight</b>															
Low	40.8	2.4	36.2-45.6	43.0	2.9	37.4-48.7	50.9	2.5	46.0-55.8	52.6	2.4	47.9-57.2	55.6	2.4	50.8-60.3
Normal	56.0	1.3	53.4-58.5	50.5	1.5	47.6-53.4	56.4	1.3	53.8-58.9	57.0	1.3	54.5-59.5	60.5	1.3	57.9-63.0

\*Confidence interval

\*\*Question changed in 2000

**Table 33**

Reasons for Not Breastfeeding, 2009

	Percent	Standard Error	95% CI *
I did not like breastfeeding	45.5	3.1	39.6-51.6
I had other children to take care of	15.9	2.2	12.1-20.8
I went back to school or work	12.3	2.0	8.9-16.9
Other reasons	13.6	2.0	10.1-18.1
I was sick or on medicine	14.2	2.1	10.7-18.8
I had too many household duties	9.2	1.8	6.2-13.3
I wanted my body back to myself	3.4	1.1	1.8-6.4
I was embarrassed to breastfeed	6.9	1.5	4.5-10.6
Baby was sick	3.4	1.0	1.9-6.0
I tried but it was too hard	9.9	1.8	6.9-14.0
I didn't want to	29.0	2.8	23.9-34.7

\*Confidence interval

**Table 34**

How did you learn to install and use your infant car seat? 2009

	Percent	Standard Error	95% CI *
I read the instructions	55.2	1.4	52.5-57.9
A health or safety professional showed me	27.4	1.2	25.1-29.9
A friend or family member showed me	21.8	1.1	19.6-24.1
I figured it out myself	14.3	1.0	12.5-16.3
Some other way	5.3	0.6	4.2-6.6
I already knew how to install it because I have other children	49.4	1.4	46.7-52.1

\*Confidence interval

**Table 35**

Source of Payment for Prenatal Care, 2009

	Percent	Standard Error	95% CI *
Health insurance from your job or the job of husband, partner or parents	47.0	1.4	44.4-49.7
Health insurance that you or someone else pays for (not from a job)	2.1	0.4	1.4-3.0
Medicaid, All Kids, Moms and Babies	53.2	1.4	50.5-55.9
TRICARE or other military health care	--	--	--
Other sources	0.5	0.2	0.3-1.1

\*Confidence interval

-- Fewer than six respondents

**Table 36**

Source of Payment for Delivery, 2009

	Percent	Standard Error	95% CI *
Health insurance from your job or the job of husband, partner or parents	45.3	1.4	42.7-48.0
Health insurance that you or someone else pays for (not from a job)	1.7	0.3	1.1-2.5
Medicaid, All Kids, Moms and Babies	54.2	1.4	51.5-56.8
TRICARE or other military health care	0.7	0.2	0.4-1.4
Other sources	0.7	0.2	0.4-1.3

\*Confidence interval

**Table 37**

Prevalence of Health Insurance or Medicaid for Baby, 2009

	Percent	Standard Error	95% CI *
Yes	98.1	0.4	97.2-98.7
No	1.9	0.4	1.3-2.8

\*Confidence interval

**Table 38**

Type of Insurance Plan for Baby, 2009

	Percent	Standard Error	95% CI *
Health insurance from your job or the job of husband, partner or parents	42.5	1.4	39.9-45.2
Health insurance that you or someone else pays for (not from a job)	2.0	0.4	1.3-2.8
Medicaid, All Kids, Moms and Babies	55.9	1.4	53.1-58.5
TRICARE or other military health care	--	--	--
Other sources	--	--	--

\*Confidence interval

-- Fewer than six respondents